

1 **Isolation, Characterization and Antibiotic Susceptibility of**
2 ***Mycoplasma hominis* and *Ureaplasma urealyticum* from Infertile**
3 **Women in Lagos, Nigeria**

(Authors need to note that research topic contradicts methodology content, samples were collected from pregnant patients).

4 **ABSTRACT**

5 **Background:**

6 *Mycoplasma hominis* and *Ureaplasma urealyticum* are important aetiologic agents of cervicitis,
7 vaginitis, postpartum sepsis, reproductive infections and infertility.

8 **Aim:**

9 **Patients and Methods:**

10 A total of 90 High vaginal Swab (HVS) and first void urine specimens were collected from
11 infertile
12 women attendees of two fertility clinics while 45 HVS and first void urine specimens were also collected
13 from pregnant women attending antenatal clinics at Lagos University Teaching Hospital and 68 Nigerian
14 Army Reference Hospital (68 NARHY) Lagos. All the specimens were inoculated into *Mycoplasma* broth
15 and subsequently Blood Agar plates, incubated appropriately and identified. Antibiotic susceptibility
16 tests were carried out on the 52 isolates. Polymerase chain reaction (PCR) was used to detect the
17 organisms in all the collected specimens. (Presentation not detail enough: authors should state the
18 number of subjects that took part in the research or number of specimen collected altogether before
19 subdividing it into subgroups. Also, authors should include statistical package used for sample analysis).

20 **Results:**

21 Of the 90 HVS specimens collected from infertile women, 9 (10.0%) were positive for *M.*
22 *hominis*, while 21 (23.3%) were positive for *U. urealyticum*. For the pregnant women using HVS
23 specimens, 6 (13.3%) were positive for *M. hominis* while 5 (11.1%) were positive for *U. urealyticum*. The
24 first void urine specimens gave lower values in both the infertile and pregnant women. Prevalence of *U.*
25 *urealyticum* was higher in infertile women than in pregnant women ($p < 0.05$). The PCR technique gave
26 higher values of 78.5% and 71.1% using HVS specimens for the infertile and pregnant women
27 respectively for *Mycoplasma/Urealyticum* species. The antibiotic susceptibility test showed that all the
28 isolates of *M. hominis* (n=18) were sensitive to Tetracycline (%) and Ciprofloxacin (%) while all the isolates
29 of *U.*
30 *urealyticum* (n=34) were sensitive to Tetracycline (%) and Erythromycin (%). (Authors should separate the
31 PCR isolate base on sample source).

32 **Conclusion:**

33 The significantly higher prevalence of *U. urealyticum* infection in infertile women (%) compared
34 to
35 the lower prevalence in pregnant women (%) may suggest that *U. urealyticum* can be incriminated in
36 infertility. HVS specimen is preferred over urine specimens for the detection of *Mycoplasma* and

- 30 *Ureaplasma*. Application of the PCR method, where affordable, is recommended for rapid and sensitive
31 detection of *Mycoplasma and Ureaplasma* in HVS specimens. Tetracycline is the antibiotic of choice,
32 unless contraindicated, for the treatment of the infections. (Sample size is too small for authors to
conclude antibiotics of choice).
- 33 Keywords: Prevalence, Infertility, Pregnancy, Nigeria, *Mycoplasma hominis*, *Ureaplasma urealyticum*.

34

INTRODUCTION

35 Infertility is considered when couples have been trying to achieve pregnancy with frequent
36 sexual intercourse for at least a year without success¹. Documented data revealed that approximately
37 72.4 million couples are infertile¹. The causes of 25% of the cases of infertility are still unknown².
38 Majority of infertile females have inflammatory changes of the oviduct or the surrounding peritoneum
39 and most of these alterations are caused by infections³.

40 Generally, *Mycoplasma hominis* and *Ureaplasma urealyticum* have been isolated from genital
41 mucosal surfaces, vagina and cervical parts of females^{4,5}. They have been isolated from genital
42 infections in both males and females⁶. *U. urealyticum* is a major cause of non-chlamydial and non-
43 gonococcal urethritis, chorioamnionitis, acute prostatitis, vaginitis, cervicitis, preterm delivery and sepsis
44^{6,7}. *M. hominis* is often associated with vaginitis, cervicitis, postpartum sepsis, pyelonephritis, preterm
45 labour and premature birth^{8,9}. Bacterial vaginosis is strongly implicated in female infertility, and
46 screening and treatment of bacterial vaginosis during the course of infertility management increased the
47 rate of pregnancy¹⁰. The isolation of these organisms in the diagnostic laboratories is cumbersome and
48 takes several days to achieve. The identification from clinical specimens using Nucleic Acid Amplification
49 Test is very expensive for routine purposes. We therefore decided to determine the prevalence of
50 *Mycoplasma* infections among infertile and pregnant women in Lagos Metropolis, Nigeria; and to
51 determine the susceptibility patterns of the isolates to some of the commonly prescribed antibiotics.

(Authors should search for more related journals and add to literature review).

52

PATIENTS MATERIALS AND METHODS

53 **Ethical Issues:** The study was approved by the Ethical Committees of Lagos University Teaching Hospital
54 (LUTH) and the 68 Nigerian Army Reference Hospital, Yaba (68NARHY). All the specimens (Urine and
55 HVS) were from married infertile women attending the Gynecology Clinics as part of a work-up for
56 fertility investigations after failing to conceive for at least one year of unprotected sexual intercourse.
57 Also included in the study were expectant mothers who were normal antenatal clinic attendees for
58 routine medical attention. Subjects with any clinical symptom of sexually transmitted disease were
59 excluded from the study. (This should be part of methodology not ethical issue; inclusion and exclusion
criteria).

60 **Sample Collection:** From July 2012 to September 2012, a total of 270 HVS and urine specimens were
61 collected from the study centres in females aged between 22 to 45 years. One hundred and forty (140)
62 HVS/Urine specimens were collected from infertile women in LUTH and 130 HVS/Urine specimens were
63 from infertile women in 68 NARHY. Ninety specimens (HVS/Urine) were also collected from pregnant
64 women from both study centres. The HVS specimens were collected by clinicians using disposable
65 speculum while the subjects were informed on how to collect the urine specimens devoid of
66 contamination (Authors need to clarify sample size. Also, authors need to state number of samples from
each source).

67 ***Mycoplasma hominis* and *Ureaplasma urealyticum* isolation and identification:**

68 The specimens were inoculated into the *Mycoplasma* transport/growth medium without delay. Clearly
69 labelled specimens were treated by inoculating 0.1ml into 5ml of prepared *Mycoplasma* broth at the site

70 of specimen collection and transported to the laboratory. The remaining urine specimens (about 10ml

71 each) were stored at -70°C. Both the urine and HVS specimens were incubated for up to 48h at 37°C in
 72 5% CO₂ for *Ureaplasma* and up to 5 days for Mycoplasmas. These were examined daily for turbidity as
 73 evidence of growth. Subsequently, the broth cultures were subcultured onto solid blood agar media and
 74 incubated at 37°C in 5%CO₂ for 24 to 48 hours for *Ureaplasma* and up to 5 days for Mycoplasmas. Urea
 75 for urease activities and L-arginine for arginine utilization were used for biochemical tests. Strains of *M.*
 76 *hominis* ATCC 23114 and *U. urealyticum* ATCC 33175 were used as positive controls. Colonies presenting
 77 a fried egg appearance suggested the presence of *M. hominis*, while colonies that were brown and tiny
 78 indicated the presence of *U. urealyticum*. *U. urealyticum* also hydrolysed urea while *M. hominis*
 79 metabolised the L. arginine. (Procedure not referenced)

80 **Polymerase Chain Reaction Confirmation of *U. urealyticum* and *M. hominis***

81 A PCR technique was used to detect *U. urealyticum* and *M. hominis*. Two millilitres of urine and
 82 HVS broth specimens was used Ref: standard strains of *U. urealyticum* (ATCC 33175); *M. hominis* (ATCC
 83 23114) and *Clostridium difficile* (ATCC BA-2155) were used. (State the PCR procedure and reference it)

84 **Antimicrobial Susceptibility Test on the Isolates**

85 Antimicrobial susceptibility test was carried out on the isolates using the modified Kirby-Bauer method
 86 with Tetracycline (30µg), Gentamycin (10µg), Erythromycin (15µg), Streptomycin (30µg) and
 87 Ciprofloxacin (5µg). (Not referenced and include statistical package used for sample analysis).

88 **RESULTS**

89 A total of 90 High Vaginal Swabs from infertile women and 45 HVS specimens from pregnant
 90 women were analysed using culture methods and PCR for *M. hominis* and *U. urealyticum*. Out of the 90
 91 HVS specimens from the infertile women, 9 (10.0%) were positive for *M. hominis* while 21 (23.3%) were
 92 positive for *U. urealyticum*. The urine specimens from both the infertile and pregnant women gave
 93 lower prevalences of 3.3% for *M. hominis*, 6.7% for *U. urealyticum* and 0.0% for *M. hominis*, 4.4% for *U.*
 94 *urealyticum* respectively (Table 1). HVS specimens yielded more isolates than the urine
 specimens. (Authors did not state numbers of urine sample collected from infertile and pregnant woman).

95 $\chi^2= 72.174$; $P<0.05$.

96 Table 2 shows the distribution of *Mycoplasma/Ureaplasma* organisms in infertile and pregnant
 97 women in the study area using *Ureaplasma* using PCR. Out of the 45 HVS specimens from the pregnant
 98 women, 32 (71.1%) were *Mycoplasma/Ureaplasma*. The 78 analysed urine specimens from the infertile
 99 women gave a prevalence of 58.9% of *Mycoplasma/Ureaplasma* infections. For the pregnant women,
 100 out of the screened 45 urine specimens 12 (26.6%) were positive for *Mycoplasma/Ureaplasma*. (Authors
 need to communicate research results with clarity; separate *Mycoplasma hominis* from *Ureaplasma*
urealyticum in each sample studied).

101 The antibiotic susceptibility patterns of the *M. hominis* and *U. urealyticum* is shown in Table 3.
 102 *M. hominis* showed 100% sensitivity to Tetracycline and Ciprofloxacin while *U. urealyticum* showed
 103 100% sensitivity to Tetracycline and Erythromycin.

104

105 **Table 1:** Distribution of *M. hominis* and *U. urealyticum* in Infertile (n=90) and Pregnant (n=45)
 106 Women in Lagos, Nigeria.

| Source | Organism | No. of Isolates | Percentage |
|------------------------|-----------------------|-----------------|------------|
| Infertile Women | | | |
| HVS | <i>M. hominis</i> | 9 | 10.0 |
| | <i>U. urealyticum</i> | 21 | 23.3 |
| Urine | <i>M. hominis</i> | 3 | 3.30 |
| | <i>U. urealyticum</i> | 6 | 6.70 |
| Pregnant Women | | | |
| HVS | <i>M. hominis</i> | 6 | 13.30 |
| | <i>U. urealyticum</i> | 5 | 11.10 |
| Urine | <i>M. hominis</i> | 0 | 0.00 |
| | <i>U. urealyticum</i> | 2 | 4.40 |

107
 (Authors should communicate research results as presented on table 1).

108 **Table 2:** Distribution of *Mycoplasma/Ureaplasma* Species in Infertile and Pregnant Women in
 109 Lagos, Nigeria using PCR

| Source | Organism | No. of Isolates | Percentage |
|------------------------|------------------------------|-----------------|------------|
| Infertile Women | | | |
| HVS (n=65) | <i>Mycoplasma/Ureaplasma</i> | 51 | 78.5 |
| Urine (n=78) | <i>Mycoplasma/Ureaplasma</i> | 46 | 58.9 |
| Pregnant Women | | | |
| HVS (n=45) | <i>Mycoplasma/Ureaplasma</i> | 32 | 71.1 |
| Urine (n=45) | <i>Mycoplasma/Ureaplasma</i> | 12 | 26.6 |

110
 (Authors should separate isolates; it is different organism form same source)

111 **Table 3:** Antimicrobial Susceptibility Patterns of *M. hominis* and *U. urealyticum* Isolated from
 112 Infertile and Pregnant Women in Lagos, Nigeria.

| Source | Organism | No. of Isolates | Antimicrobial Agent Activities | | | | |
|--------|-----------------------|-----------------|--------------------------------|--------|----------|----------|--------|
| | | | % Sensitive | | | | |
| | | | Te | CN | E | CPX | S |
| HVS | <i>M. hominis</i> | 15 | 15(100) | 0(0.0) | 0(0.0) | 15(100) | 0(0.0) |
| | <i>U. urealyticum</i> | 26 | 26(100) | 0(0.0) | 26(100) | 0(0.0) | 0(0.0) |
| Urine | <i>M. hominis</i> | 3 | 3(100) | 0(0.0) | 0(0.0) | 3(100) | 0(0.0) |
| | <i>U. urealyticum</i> | 8 | 8(100) | 0(0.0) | 8(100) | 0(0.0) | 0(0.0) |
| Total | | 52 | 52(100) | 0(0.0) | 34(65.4) | 18(34.6) | 0(0.0) |

113
 114 **Key:**

115 Te = Tetracycline (30µg)

116 CN=Gentamycin (10µg)

117 E=Erythromycin (15µg)

118 CPX=Ciprofloxacin (5µg)

119 S=Streptomycin (30µg)

120 DISCUSSION

121 The results of our study showed that *M. hominis* and *U. urealyticum* infections are prevalent
122 among the infertile and pregnant women in Lagos Metropolis, Nigeria. In this study both the HVS and
123 Urine specimens were examined using culture and PCR techniques. The culture technique gave a
124 prevalence, using the HVS specimens, of *M. hominis* (10.0%) and 13.3% for infertile and pregnant
125 women respectively. The urine specimens yielded lower values for *M. hominis* with a prevalence of 3.3%
126 and 0.0% for infertile and pregnant women respectively. A prevalence of 6.7% for infertile women and
127 4.4% for pregnant women were recorded for *U. urealyticum* infection. The PCR technique expectedly
128 gave higher prevalence of 78.5% and 71.1% using HVS specimens for *Mycoplasma/Ureaplasma* in
129 infertile and pregnant women respectively. The urine specimens gave lower values of 58.9% and 26.6%
130 for *Mycoplasma/Ureaplasma* infection in infertile and pregnant women respectively. The urine
131 specimens gave lower values of 58.9% and 26.6% for *Mycoplasma/Ureaplasma* infection in infertile and
132 pregnant women respectively. One possible explanation for the higher prevalences of these infections in
133 infertile women is hormonal disorders which can lead to reduced levels of immunity and increased
134 bacterial colonization and survival in the vaginal epithelium¹².

135 In addition to the considerably high prevalence of *M. hominis* and *U. urealyticum* infections in
136 our study, several published reports revealed various prevalence rates. Agbakoba *et al*¹² reported a
137 (italicize *et al*) prevalence of 35.7% in women and of reproductive age in Ibadan. The prevalence rate of
138 *M. hominis* and *U. urealyticum* in HVS specimens ranged from zero to 3.8%^{14, 15}. The prevalence of *U.*
139 *urealyticum* infection was reported from 20% in South Africa to 41.9% and 51.5% in Italy and Africa
140 respectively¹⁴. In contrast, the prevalence was lower in some other studies^{16, 17}. The varied prevalence
141 rates all over the
142 world may be due to the use of only culture methods or PCR. In addition there are many factors that will
143 affect the culture results such as the use of transport medium, the duration between the collection of
144 the sample and the inoculation, and the duration of incubation. The Nucleic Acid Amplification tests
gave higher results due to its sensitivity and it is also a rapid test, although it is more expensive.

145 In our study, it was observed that *M. hominis* and *U. urealyticum* were detected more from HVS
146 than urine specimens. This is in agreement with Taylor-Robinson¹⁷ who reported that the numbers of
147 isolates from urine specimens were usually 10 fold less than in swabs.

148 In vitro antibiotic susceptibility test showed that all the isolates of *M. hominis* and *U.*
149 *urealyticum* were sensitive to Tetracycline(%). The implication of this finding is that where it is not
150 possible to detect the causative agent of non-gonococcal urethritis, it is safer to prescribe Tetracycline
instead

151 of Erythromycin as being practiced in some centres. (Base on sample size, it is not good enough to
 152 conclude. More work need to be done to establish authors claim) *U. urealyticum* isolates in addition to
 Tetracycline

were all sensitive to Erythromycin, while all *M. hominis* isolates were also sensitive to Ciprofloxacin.

153 In conclusion therefore, we have indicated that *M. hominis* and *U. urealyticum* infections are
 154 prevalent in both the infertile and pregnant women in Lagos metropolis. Efforts, where feasible, should
 155 be made to detect the presence of these organisms during the work-up of cases of infertility and in
 156 pregnancy to prevent the adverse outcomes. We therefore recommend the precise monitoring of fertile
 157 and infertile females for the presence of *M. hominis* and *U. urealyticum* and treatment of positive cases
 158 to prevent diseases and possibly infertility.

(Authors should include research limitation)

(All et al. should be expunge and add names of all authors)

159 REFERENCES

- 160 1. Boivin, J; Bunting L; Collins JA et al. International estimates of infertility prevalence and
 161 treatment-seeking: Potential need and demand for infertility medical care. *Hum Reprod.*
 162 2007;22(16): 1506-1512. Doi:10.1093/hum re/dem046. [PubMed][Cross Ref]
- 163 2. Daar AS, Merali Z. Infertility and social suffering: the case of ART in developing countries. In:
 164 Vayena E, Rowe P, Griffin D, editors. Report of a meeting on "Medical, ethical and social aspects
 165 of assisted reproduction." Geneva: WHO, 2001. Pp.16-21.
- 166 3. Wiesenfeld HC, Hillier SL, Meyn LA et al. Subclinical pelvic inflammatory disease and infertility.
 167 *Obstet Gynecol* (Not italicize). 2012;120(1):37-43. Doi:10.1097/AOG.Ob 013e31825a6bc9.
 168 Ref]
- 169 4. Capoccia R, Greub G, Baud D. *Ureaplasma urealyticum*, *Mycoplasma hominis* and adverse
 170 pregnancy outcomes. *Curr Opin Infect Dis.* 2013; 26(3):231-240.doi:10. 1097/QCO.
 171 Ob013e328360db58. [PubMed][Cross Ref]
- 172 5. Wang QY, Li RH, Zheng LQ et al. Prevalence and antimicrobial susceptibility of *Ureaplasma*
 173 *urealyticum* and *Mycoplasma hominis* in female outpatients, 2009-2013. *J Microbiol, Immunol*
 174 *Infection*(Not italicize). 2009-2013. [PubMed].
- 175 6. Al-Sweih NA, Al-Fadli AH, Omu AE et al. Prevalence of *Chlamydia trachomatis*, *Mycoplasma*
 176 *hominis*, *Mycoplasma genitalium* and *Ureaplasma urealyticum* infections and seminal quality in
 177 infertile and fertile men in Kuwait. *J Androl.*2012;33(6):1323-1329.doi:10. 2164/jandrol.
 178 111.013821. [PubMed] [Cross Ref].
- 179 7. Zhu C, Liu J, Ling Y et al. Prevalence and antimicrobial susceptibility of *Ureaplasma urealyticum*
 180 and *Mycoplasma hominis* in Chinese women with genital infectious diseases. *Indian J Dermatol,*
 181 *Venereology, Leprology.* 2012;78(3):406-407 [PubMed].
- 182 8. Aydin Y, Atis A, Ocer F et al. Association of cervical infection of *Chlamydia trachomatis*,

- 183 Ureaplasma urealyticum and Mycoplasma hominis with peritoneum colonization in pregnancy. J
184 **Obstet Gynecol.** 2010, 30(8):809-812.doi:10.3109/01443615.2010. 519063. [PubMed][Cross Ref]

- 185 9. Campos GB, Lobao TN, Selis NN [et al.](#) Prevalence of Mycoplasma genitalium and Mycoplasma
186 hominis in urogenital tract of Brazilian Women. [BMC Infect Dis](#) 2015;15(1):60. [PMC free
187 article][PubMed]
- 188 10. Salah RM, Allam AM, Magdy AM [et al.](#) Bacterial vaginosis and infertility: cause or association?
189 [Eur J. Obstet Gynecol Reprod Biol.](#) 2013;167(1): 59-63.doi:10.1016/j. ejogrb. 2012.10.031. [Pub
190 Med] [Cross Ref]
- 191 11. Clinical and Laboratory Standards Institute: Methods for antimicrobial dilution and disk
192 susceptibility testing of infrequently isolated or fastidious bacteria, approved guideline M45-A
193 Wayne, Pa, 2005, CLSI.
- 194 12. Kornats'ka AH. Local humoral immunity in women with combined forms of infertility. *Lik sprava.*
195 1998(4): 82-84 [PubMed].
- 196 13. Agbakoba NR, Adetosoye AI, Adewole IF. Presence of Mycoplasma and Ureaplasma species in
197 the vagina of women of reproductive age. [West Afr J Med.](#) 26(1).
- 198 14. Leli C, Mencacci A, Bombaci JC [et al.](#) Prevalence and antimicrobial susceptibility of Ureaplasma
199 urealyticum and Mycoplasma hominis in population of Italian and immigrant outpatients. [Infez](#)
200 [Med](#) 2012; 20(2):82-87 [PubMed]
- 201 15. Elias M, Grzesko J, Siejkowski R [et al.](#) The presence of Mycoplasma hominis and Ureaplasma
202 urealyticum in the cervical of uterus. [Ginekol Pol.](#) 2005; 76(1):28-32. [PubMed].
- 203 16. Blanchard A, Yanez A, Dybvig K [et al.](#) Evaluation of intraspecies genetic variation within 16S
204 rRNA gene of Mycoplasma hominis and detection by polymerase chain reaction. [J Clin Microbiol.](#)
205 1993;31(5):1358-1361.
- 206 17. Taylor-Robinson, D. Genital Mycoplasma infection. [Clin Lab Med](#) 9:501-523.