

Original Research Article

MICROORGANISMS ASSOCIATED WITH THE PRODUCTION OF A NIGERIAN FERMENTED BEVERAGE, 'AGADAGIDI'

ABSTRACT

Aim: The microbial types, occurrence, loads and interactions were studied during the production of a Nigerian fermented beverage, 'Agadagidi', from overripe plantains.

Place and Duration of Study: Federal University of Technology Akure, Ondo State, Nigeria. March- July, 2012.

Methodology: Isolation, enumeration and identification of bacteria and fungi were carried out by using standard pour plate, morphological, biochemical and physiological characterization methods. Antagonistic and mutualistic interactions among the microorganisms were investigated using agar well diffusion method.

Results: *Bacillus subtilis*, *B. megaterium*, *Staphylococcus aureus*, *Escherichia coli*, *Enterococcus* species, *Lactobacillus plantarum*, *Pediococcus acidilactici*, *L. fermentum*, *Leuconostoc mesenteroides*, *Aspergillus flavus*, *A. niger*, *A. fumigatus*, *Penicillium notatum*, *Trichoderma viride*, *Saccharomyces cerevisiae*, *Candida utilis* and *Geotrichum* species were identified during the production of the beverage. The loads of total bacteria (TB), Lactic Acid Bacteria (LAB), enterobacteriaceae and fungi of the uncrushed plantain pulp were $2.6 \times 10^7 \pm 0.05$ cfu/ml, $6.7 \times 10^6 \pm 0.05$ cfu/ml, $3.8 \times 10^5 \pm 0.05$ cfu/ml and $2.0 \times 10^7 \pm 0.05$ cfu/ml. At 0 hour of fermentation, the loads of total bacteria, fungi and enterobacteriaceae increased. Then after, the total bacteria, enterobacteriaceae and fungi counts decreased to $2.0 \times 10^5 \pm 0.11$ cfu/ml, $1.3 \times 10^5 \pm 0.11$ cfu/ml and $1.03 \times 10^5 \pm 0.05$ cfu/ml respectively. In contrast, the LAB cell number increased to $8.6 \times 10^7 \pm 0.1$ cfu/ml at the end (48 hours) of fermentation. The level of the microbial occurrence was 25 to 100% with *B. subtilis*, *L. plantarum*, *L. mesenteroides*, *S. cerevisiae* and *C. utilis* occurring as the highest. *B. megaterium*, *E. spp.*, *A. niger* and *T. viride* occurred least. There was positive co-existence between Yeast and LAB. The yeasts and LAB exhibited antagonism against other bacteria.

Conclusion: The data obtained in this work has shown some functional microflora and their relationship during the production of "Agadagidi". This information can contribute to a better understanding of the "Agadagidi" production process for a consistent quality beverage.

Keywords: Microbial community; “Agadagidi”; Fermented beverage; Microbial loads; Antagonism.

1. INTRODUCTION

Plantain (*Musa sapientum* var. *Paradisiacal* Linn) is one of the most important staple food crops for millions of people both in developed and developing countries. It is one of the foods commonly consumed in the West Africa sub-region, Northern America, Mexico and the Caribbean. In Nigeria, its consumption cuts across the indigenous groups and the numerous socio-economic classes because of the ease of preparation and consumption [1]. In West and central Africa, more than 10 million tons are produced annually and are traded locally [2, 3].

“Agadagidi,” a cloudy effervescent sweet-sour taste typical African traditional alcoholic beverage is made from overripe bananas and plantains through fermentation. It is common in south-western part of Nigeria [4].

The techniques used in the production of wines from tropical fruits are similar to those of grape wine production which include pressing out the juice, fermenting, maturing and bottling [5]. Fermentation leads to changes in appearance of food which is characterized by different reactions of microorganisms [6]. Food fermentation represents one of the oldest known uses of biotechnology. Fermented foods and beverages form a significant proportion of all diets worldwide; they are typically about one-third of the foods consumed globally. Fermentation of foods covering a wide range of microbial and enzymatic processing of foods and ingredients is used to achieve desirable characteristics such as prolonged shelf-life, improved safety, attractive flavour, nutritional enrichment and promotion of health [7, 8]. The fermentation of overripe plantain to produce “Agadagidi” is a waste prevention processing of plantain. Plantain is a perishable crop which has much less value when it is overripe; hence it is used for wine production [9-11].

At present, there is no adequate information on the spectrum of microorganisms and microbial interaction associated with the production of overripe plantain pulp to yield ‘Agadagidi’. **i.e. from the raw material to the finished product.** Microbial information on the production of “Agadagidi” will contribute to the development of starter cultures with predictable characteristics, for use in small-scale and commercial production of stable and consistent quality ‘Agadagidi’. Therefore, this research was proposed to reveal the microbial community and confirmation of mutualism or commensalism and antagonistic interaction during the production of “Agadagidi”.

2.0 MATERIALS AND METHODS

2.1 Traditional Preparation of “Agadagidi” sample

The production of “Agadagidi” was done in the laboratory based on the local or indigenous method. The plantains used were bought from the king’s market, Akure, Ondo State. The Overripe plantain pulps peels were washed in tap water to remove debris and dirt. The plantain was peeled and the pulps were crushed in portable tap water at a ratio of 1:5 (w/v) in a sterilized container, covered and left to ferment for 2 days at ambient temperature of $27\pm 2^{\circ}\text{C}$. Samples were withdrawn from peeled, uncrushed pulps and at 0 h (This is immediately after water is added into the crushed plantain pulp), 24 h and 48 h of fermentation. The fermented liquid was filtered with a sterilized muslin cloth to remove the plantain mashes. The liquid then served as “Agadagidi”.

2.2 Enumeration and Isolation of Microorganisms

Serial dilutions (1:10 v/v) were made with the samples collected from peeled, uncrushed pulps and at 0 h, 24 h and 48 h of fermentation and pour plated on nutrient agar for mesophilic bacteria, Man Rogosa Sharpe agar at pH 5.5 for lactic acid bacteria (LAB), Malt Extract Agar (supplemented with streptomycin sulphate) for yeasts, Eosin Methylene Blue for the isolation of the member enterobacteriaceae. Incubation was carried out at 37°C for total bacteria and LAB and 27°C for fungi. Colonies and spore forming units formed on the media were counted and subcultured. The Bacteria isolate were observed using microscopy, Gram staining, sugar fermentation test, biochemical tests such as urease test, catalase test, citrate utilization test and indole test. While Fungal identification was done using the fungi conventional identification method according to the methods of [12-15].

2.3 Determination of Positive and Negative Microbial Interactions between the Isolates

Mutualism/commensalism and antagonism between the microbial isolates were determined using Muller Hilton agar and Agar Well Assay method with slight modification as described by [16]. Pre-poured Muller Hillton agar (MHA) in separate Petri dishes containing various bacterial and yeast’s cells were bored using a sterile cork borer of 5 mm diameter and 1 mL of each test isolate was transferred into each well, incubated for 24 hours at 37°C . The agar was examined for zones of inhibition which were measured in millimetres. Creation of inhibitory zone indicated antagonism and absence of zone of inhibition signified no inhibition.

2.4 Statistical analysis

Data obtained were subjected to Analysis of Variance (ANOVA) and separation of means was done with Duncan's New Multiple Range Test at 95% confidence level using SPSS 16.0 version.

3.0 RESULTS

The microbial composition of 'Agadagidi' is made up of nine bacteria and eight fungi. The bacteria consist of two species of *Bacillus* and *Lactobacillus*, one species of *Staphylococcus*, *Enterococcus*, *Escherichia coli*, and *Leuconostoc*. *Pediococcus acidilactici* was also isolated. Among the fungi, three species of *Aspergillus*, one species of *Trichoderma* and *Penicillium* were the moulds isolated. *Geotrichum*, *Saccharomyces* and *Candida utilis* were the yeasts obtained (Table 1).

Table 1 also revealed the types of microbes associated with the "Agadagidi" production. *Bacillus subtilis*, *Leuconostoc mesenteroides*, *Lactobacillus plantarum*, *Saccharomyces cerevisiae* and *Candida utilis* had the highest (100%) occurrence throughout the stages of production. *Bacillus megaterium*, *Enterococcus* spp., *A. niger* and *T. viride* occurred lesser (25%) than other microorganisms.

Table 1: The occurrence of bacteria and fungi during the production of "Agadagidi"

Microorganism	Plantain Pulp (Uncrushed)		Fermentation: (hr)		Level of Occurrence (%)
	0	24	48		
<i>Bacillus subtilis</i>	+	+	+	+	100
<i>Bacillus megaterium</i>	+	-	-	-	25
<i>Staphylococcus aureus</i>	+	+	-	-	50
<i>Escherichia coli</i>	+	+	-	-	50
<i>Enterococcus</i> species	-	+	-	-	25
<i>Lactobacillus plantarum</i>	+	+	+	+	100
<i>Lactobacillus fermentum</i>	+	+	+	-	75
<i>Leuconostoc mesenteroides</i>	+	+	+	+	100
<i>Pediococcus acidilactici</i>	-	+	+	+	75
<i>Aspergillus flavus</i>	+	+	-	-	50
<i>Aspergillus niger</i>	+	+	-	-	50
<i>Aspergillus fumigatus</i>	+	+	-	-	50
<i>Penicillium notatum</i>	+	+	-	-	50

<i>Trichoderma viride</i>	+	-	-	-	25
<i>Sacharomyces cerevisiae</i>	+	+	+	+	100
<i>Candida utilis</i>	+	+	+	+	100
<i>Geotrichum</i> species	+	+	+	-	75

Legend:

+ Present

- Absent

During the production of “Agadagidi”, the load of bacteria, fungi and the members of enterobacteriaceae followed the same trend. Their populations increased at 24 hr and decreased at 48 hours of fermentation respectively (Fig.1, 2 and 3). The pattern of lactic acid bacteria load differs with decrease load at 24 hours and increase at 48 hours (Fig. 4).

The total bacterial load of the overripe plantain pulp was $2.6 \times 10^7 \pm 0.05$ cfu/ml which increased before fermentation i.e. after adding water to the overripe plantain pulp (0 hr) to $5.2 \times 10^7 \pm 0.05$ cfu/ml (Figure 1). This load decreased significantly (<0.05) to $2.0 \times 10^5 \pm 0.11$ cfu/ml at 48 hours of fermentation. Fungal load of $2.0 \times 10^7 \pm 0.05$ cfu/ml was obtained on the ripe plantain pulp which increased and decreased at 0 hr and 24 hr of fermentation to $2.2 \times 10^7 \pm 0.05$ cfu/ml and $4.3 \times 10^5 \pm 0.05$ cfu/ml respectively (Fig. 2). There was a continuous decrease till the end (48 hr) of fermentation, having a load of $1.03 \times 10^5 \pm 0.05$ cfu/ml. Figure 3 shows enterobacteriaceae load during the preparation of “Agadagidi.” The load was 2.8×10^7 cfu/ml at 0 hr of fermentation. There was a drastic reduction in the enterobacteriaceae load to $1.3 \times 10^5 \pm 0.11$ cfu/ml at 48 hr of fermentation.




Fig. 1: Total bacterial load during production of "Agadagidi".

Each value represents the mean value (log Cf u/mL), standard deviation (SD) from three trials and standard error.




Fig. 2: Fungal load during the production of "Agadagidi"

Each value represents the mean value (log Cf u/mL), standard deviation (SD) from three trials and standard error.

Lactic acid bacteria had a load of $6.7 \times 10^6 \pm 0.05$ cfu/ml on the overripe plantain pulp (Fig. 4). The load reduced after 24 hr of fermentation to $4.2 \times 10^5 \pm 0.05$ cfu/ml and then after increased to $8.6 \times 10^7 \pm 0.1$ cfu/ml at 48 hr.




Fig. 3: Enterobacteriaceae load during the production of “Agadagidi”.

Each value represents the mean value (log Cf u/mL), standard deviation (SD) from three trials and standard error.




Fig. 4: Lactic acid bacteria load during the production of “Agadagidi”.

Each value represents the mean value (log Cf u/mL), standard deviation (SD) from three trials and standard error.

162
 163TTable 2: Interaction between Lactic acid bacteria, yeasts and some of the bacteria isolated during
 164 the production of “Agadagidi”
 165

Microorganism (LAB and yeast)	<i>Enterococcus</i> spp.	<i>Lactobacillus</i> <i>plantarum</i>	<i>Bacillus</i> <i>subtilis</i>	<i>Lactobacillus</i> <i>Fermentum</i>	<i>Escherichia</i> <i>coli</i>	<i>Staphylococcus</i> <i>aureus</i>	<i>Bacillus</i> <i>megaterium</i>	<i>Leuconostoc</i> <i>mesenteroides</i>	<i>Saccharomyces</i> <i>cerevisiae</i>	<i>Candida</i> <i>utilis</i>
<i>Leuconostoc</i> <i>Mesenteroides</i>	+	-	+	-	+	+	+	-	-	-
<i>Lactobacillus</i> <i>fermentum</i>	+	-	+	-	+	+	+	-	-	-
<i>Lactobacillus</i> <i>plantarum</i>	+	-	+	-	+	+	+	-	-	-
<i>Lactococcus</i> <i>Lactis</i>	+	-	+	-	+	+	+	-	-	-
<i>Saccharomyces</i> <i>cerevisiae</i>	+	-	+	-	+	+	+	-	-	-
<i>Candida utilis</i>	+	-	+	-	+	+	+	-	-	-

166 **Legend:**

167 + = Positive interaction (No Zone of inhibition found)

168 - = Negative interaction (zone of inhibition)

169 Table 2 shows the interactions which occurred among the microorganisms *in vitro*. Lactic
 170 acid bacteria (LAB) and yeasts inhibited the growth of other bacteria which were all spoilage
 171 and pathogenic microorganisms. The interaction between the yeasts and the lactic acid bacteria
 172 were positive as zone of inhibition was not created when they were co-cultured.

173 **DISCUSSION**

174 Many species of microorganisms were isolated during the production of “Agadagidi.” The
 175 production stages comprises of both pre-fermentation and during fermentation. The cultural and
 176 biochemical properties exhibited by the microorganisms were similar to those described by [12-
 177 14] which in turn gave the names of the microorganisms in table 1. The 100% occurrence of
 178 *Leuconostocmesenteroides*, *L. plantarum* and *Bacillus subtilis* isolated could be as result of their
 179 dominancy and the ability to withstand acidic condition associated with the fermentation
 180 stages. *Leuconostocmesenteroides* and *L. plantarum* are heterofermentative, hence they were
 181 highly resistant to acid. [17] confirmed that their dominancy is determined by the sensitivities of
 182 microorganisms to the acidic conditions that develop during the fermentation. The presence of *B.*
 183 *subtilis* during the production of “Agadagidi” may be due to contamination through

their endospores from dust, air and peels. Bacilli are spore forming bacteria, able to withstand harsh conditions which are widely distributed in nature and in many cases with a pH as low as 3.9 [18, 19]

The appearance of the two yeasts, *Saccharomyces cerevisiae* and *Candida utilis* throughout the stages of the production of 'Agadagidi' can be attributed to the environment and the fruits itself. These yeasts are naturally associated with ripened fruits and they are known to be responsible for alcoholic fermentation [20-21]. This findings is similar to the report of [22] who identified yeasts species in "Tchapalo" production and observed highest frequency with *S. cerevisiae* (87.36%), followed by *Candida tropicalis* (5.45%) and *Meyerozyma caribbica* (2.71%). The presence of moulds on the uncrushed ripened plantain pulp, after adding water may be as a result of improper handling of the ripe plantain. The peel possibly contained moulds which might have been transferred to the plantain pulp. The elimination of moulds stated during fermentation may be as a result of high loads and activities of LAB during fermentation. Bacteria have been shown to suppress the growth of mould during fermentation [23]. [24] who worked on yeasts and moulds associated with "Ogi" similarly observed moulds on the surface of raw maize grains for "Ogi" production. The increase in the load of bacteria, fungi and enterobacteriaceae in mashed overripe plantain pulp (0 hr fermentation) may be as a result of the potable tap water added. Potable water passes through pipes that may not have been treated or sanitized for a long period of time and as a result microorganisms present in the pipes might have increased the microbial loads. [25, 26] confirmed the presence of coliform bacteria in "Kununzaki" where tap water was the source of water used. The decrease in total bacterial and fungal loads after 24 hr of fermentation may be as a result of the increase in population of the LAB that must have formed acid thereby reducing pH (acidity) which seems to be detrimental to the other bacteria. This data agrees with the work of [27, 28] who explained that LAB produce many organic acids such as lactic, acetic and propionic acids as end products during fermentation which provide an acidic environment unfavourable for the growth of many pathogenic and spoilage microorganisms.

The increment LAB population after 24 hr may be due to their ability to proliferate in the presence of the metabolic product synthesized by the yeasts. This observation is similar to that of [29] who worked on "Akamu". They observed that LAB occurred early in the fermentation, increasing rapidly from 1.6×10^7 cfu/g to 7.1×10^8 cfu/g after 72 hr. The drop in population of enterobacteriaceae during fermentation was as a result of the low pH and acid production by some of the bacteria which could destroy the cells of these microbes. Similar observation was

obtained in the [30] work, who reported decrease in the population of enterobacteriaceae as the pH dropped was due to the effect of lactic acid bacteria present in 'bushera' and 'kirario'

LAB and yeasts evaluated in this study showed antagonistic interaction against other isolated bacteria, which mean that these microbes might have produced inhibitory metabolites. This agree with the report of [31, 32] who documented that the ability of antimicrobial compounds produced by LAB enabled them to exert strong antagonistic activity against food contaminating microorganisms. None of the LAB isolates inhibited any of the yeast cultures. The association between the yeasts and LAB may therefore be mutualism. Several authors have reported similar co-existence and positive interactions between yeasts and lactic acid bacteria in different African fermented foods [24, 33].

The stimulatory effect of yeasts on lactic acid bacteria during fermentation has been attributed to the provision of some compounds such as soluble nitrogenous compounds, B-vitamins, CO₂, pyruvate, propionate, acetate and succinate [34]. It has also been shown that yeasts multiplication is associated with an increase in acid formation in fermented products.

CONCLUSION

This study has provided valuable information on the types of microbial communities and their interactions during the production of "Agadagidi". *Lactobacillus plantarum*, *Leuconostoc mesenteroides*, *B. subtilis*, *S.cerevisiae* and *C.utilis* were found to be predominant in the production of the beverage. This research also established positive and negative interactions between the isolated microorganisms. This will contribute to the development of strains with predictable characteristics for use in small and large scale production.

RECOMMENDATION

It is therefore recommended that there should be proper handling of the overripe plantain pulp and tap water should also be boiled before use so as to reduce or eliminate undesirable microorganisms.

REFERENCES

1. Solanke AO and Falade KO. A review of the uses and methods of processing banana and plantain (*Musa* spp.) into storable food products. *Journal of Agricultural Research and Development*. 2010; **9**(2): 87-96.
2. Latham, MC. La nutrition dans les pays en development, In Info Musa Africa :Abstract from International Conference on plantain and banana for Africa. *International Journal on Banana and Plantain*. 2001; 16(1):5 -12.
3. Nse-Nelson FA, Oke UR, Adindu JU. Analysis of plantain marketing in Ikwuano local government area of Abia State, Nigeria. *Nigerian Journal of Agriculture, Food and Environment*. 2016; **12**(1): 85-89.

- 253 4. Omojasola PF, Davies OF, Kayode RM. The effect of chemical preservatives, pasteurization
254 and refrigeration on the shelf life of “Agadagidi” a fermented plantain drink. *Research Journal of*
255 *Microbiology*. 2012; **7**(3): 145-157.
- 256 5. Okafor N. Modern Industrial Microbiology and Biotechnology. Science Publishers, Enfield, N. Y,
257 U. S. A. 2007; pp. 530.
- 258 6. Ray M, KuntalGhosh S, Singh K, Chandra M. Folk to functional: An explorative overview of
259 rice-based fermented foods and beverages in India. *Journal Ethnic Foods*. 2016; **3**: 5-18.
- 260 7. Holzapfel WH. Appropriate starter culture technologies for small-scale fermentation in
261 developing countries. *International Journal of Food Microbiology*. 2002; **75**: 197–212.
- 262 8. Tamang JP. Health Benefits of Fermented Foods and Beverages. CRC Press, Taylor and
263 Francis Group, New York. 2015; pp. 1-110.
- 264 9. Sanni AI, Oso BA. The production of “Agadagidi”, a Nigerian fermented beverage. *Die*
265 *Nahrung*, 1988; **32**: 319-326.
- 266 10. Chia CL, Huggins CA. Bananas. In: Community Fact Sheet fruit. Hawaii. CTAHR edition.
267 2003; pp. 23-29.
- 268 11. Areola JK, Yussuf IO. The processing and nutritional analysis of plantain wine. *International*
269 *Journal of Business and Educational Policies*. 2011; **7**(2): 238-245.
- 270 12. Deak T Beuchat LR. Yeasts associated with fruit juice concentrate. *Journal of Food*
271 *Protection*. 1993; **56**: 777-782.
- 272 13. Sanni AI, Lonner C, Marklinder L, Johansson ML, Molin G. Starter cultures for the production
273 of “Ogi”, a fermented infant food from maize and sorghum. *International Journal of Food*
274 *Chemistry, Microbiology and Technology*. 1994; **16**: 29-33.
- 275 14. Brenner J, Kreig V, Stanley T. Bergey's Manual of Systematic Bacteriology. The
276 Proteobacteria. 2nd Edition. Published by Springer, New York. 2005; pp. 587-848.
- 277 15. Fawole MO, Oso, BA. Characterization of Bacteria: Laboratory Manual in Microbiology. 5th
278 Edition. New Spectrum Books Publisher, Ibadan, Nigeria. 2007; pp. 24-33.
- 279 16. Oriola O, Boboye B, Adetuyi F. Bacterial and fungal communities associated with the
280 production of a Nigerian fermented beverage, ‘Otika’. 2017. *Journal in Advances in*
281 *Microbiology*, Under Review.
- 282 17. Ampe F, Sirvent A, Zakhia, N. Dynamics of the microbial community responsible for traditional
283 sour cassava starch fermentation studied by denaturing gradient gel electrophoresis and
284 quantitative rRNA hybridization. *International Journal of Food Microbiology*. 2001; **65**: 45– 54.
- 285 18. Hanlin JH Spoilage of acidic products by *Bacillus* species. *Dairy Foods Environment*. 1998,
286 **18**: 655-659.
- 287 19. Alo MN, Anyim C, Igwe JC, Elom M, Uchenna DS. Antibacterial activity of water, ethanol and
288 methanol extracts of *Ocimum gratissimum*, *Vernonia amygdalina* and *Aframomum melegueta*.
289 *Advances in Applied Science Research*. 2012; **3**(2): 844-848.
- 290 20. Sanni AI, Onilude AA, Ibidapo OT. Biochemical composition of infant weaning food fabricated
291 from fermented blends of cereal and soybean. *Food Chemistry*. 1999; **65**: 35-39.
- 292 21. Jay, J. M., Loessner, M. J. and Golden, D. A. (2005). Modern Food Microbiology. 7th edition.
293 Springer Science and Business Media, New York, U. S. A. pp. 101- 125.
- 294 22. N'guessan FK, Brou K, Jacques N, Casaregola S and Dje, KM. Identification of yeasts during
295 alcoholic fermentation of “Tchapalo”, a traditional sorghum beer from Côte d'Ivoire. *Anton*
296 *Leeuwenhoek International Journal of General and Molecular Microbiology*. 2011; **99**: 855-864.

- 297 23. Dalié DK, Deschamps AM, Richard FF. Lactic acid bacteria – Potential for control of mould
298 growth and mycotoxins: A review. *Food Control*.2010; **21**: 370–380.
- 299 24. Omemu AM, OyewoleOB, Bankole MO, Akintokun AK. Yeasts and moulds associated with
300 “Ogi” - a cereal based weaning food during storage. *Research Journal of Microbiology*. 2007;
301 **2**(2) 141-148.
- 302 25. Adegoke GA, Asaye O,Shridhar MK. Microbiological and physico-chemicalparameters of
303 water used by some brewery, bakery and soft drink plant in Oyo state, Nigeria. *Journal of*
304 *Science and Technology*. 1993;**3**: 92-95.
- 305 26. Amusa NA. Microbiological and nutritional quality of hawked “Kunun” (a sorghum based non-
306 alcoholic beverage) widely consumed in Nigeria. *Pakistan Journal of Nutrition*, 2007;**8**: 20-25.
- 307 27. Babatunde DA,Oladejo PO. Identification of Lactic Acid Bacteria isolated from Nigerian Foods:
308 Medical importance and comparison of their bacteriocin activities. *Journal ofNatural Sciences*
309 *Research*, 2014;**4**(23): 246-252.
- 310 28. Teshome G. Review on lactic acid bacteria function in milk fermentation and preservation.
311 *African Journal of Food Science*. 2015;**9**(4): 170-175.
- 312 29. Ogbonnaya N, Chukwu, BC. (2012). Studies on “Akamu”, a traditional fermented maize food.
313 *Review on Children Nutrition*, **39**: 180-184.
- 314 30. Muyanja, CM, NarvhusJA, Treimo, J, Langsrud T. Isolation, characterization and identification
315 of lactic acid bacteria from “Bushera”: A Ugandan traditional fermented beverage. *International*
316 *Journal of Food Microbiology*. 2003;**80**: 201-210.
- 317 31. De Martinis ECP, Publio MRP, Santarosa PR, Freitas FZ. Antilisterial activity of lactic acid
318 bacteria isolated from vacuum packaged Brazilian meat and meat products. *Brazil Journal of*
319 *Microbiology*, 2001; **32**: 32-37.
- 320 32. Ogunbanwo ST, Sanni AI, Onilude AA. Characterization of bacteriocin produced by
321 *Lactobacillus plantarum* F1 and *Lactobacillus brevis* OG1. *Journal of Applied Microbiology*.
322 2003;**99**: 77-84.
- 323 33. Gulitz A, Stadie J, Ehrmann M, Vogel R. Comparative phylobiomic analysis of bacteria
324 community of water kefir by 16S rRNA gene amplicon sequencing and ARDRA analysis.
325 *Journal of Applied Microbiology*. 2013;**114**: 1082-1091.
- 326 34. Stadie J, Gulitz A, Ehrmann M, Vogel R. Metabolic activity and symbiotic interaction of lactic
327 acid bacteria and yeasts isolated from water kefir. *Food Microbiology*. 2013;**35**: 92-98.
- 328