

Original Research Article

Diversity and Antimicrobial Activity of Hydrobionts Associated Microorganisms from the Sea of Japan with the Occurrence of Tropodithietic Acid Producing Bacteria.

ABSTRACT

The study was undertaken to survey microorganisms associated with colonial fouling hydrozoans and red alga *Polysiphonia* sp. collected from the Sea of Japan seashore and to screen them for antimicrobial effects. On the basis of 16S rRNA gene sequences the isolates were assigned to 21 genera of the *Alphaproteobacteria*, *Gammaproteobacteria*, *Actinobacteria* as the dominant followed by *Firmicutes*, *Bacteroidetes* and *Betaproteobacteria*; the most shared 98-99% sequence similarity to recognized species recovered from marine sources. Hydrozoan's and red alga microbial associations were different in their taxonomic compositions at the generic level. Members of the genera *Shewanella*, *Labrenzia* and *Streptomyces* occurred in both specimens. Antimicrobial screening revealed 46 strains capable to inhibit growth of two to seven indicator cultures. Active hydrozoan's strains were phylogenetically close to *Phaeobacter inhibens*. Active isolates from red alga were similar to *Pseudovibrio ascidiaceicola*, and in addition to *Paenibacillus xylanexedens* and *Bacillus murimartini*. *Streptomyces* strains with antimicrobial activity were found in both specimens. Strains *Phaeobacter* sp. H8 and *Pseudovibrio* sp. rh17 were selected to be examined for the production of tropodithietic acid, TDA, which is the known antimicrobial metabolite produced by *Phaeobacter* and *Pseudovibrio* bacteria. On the basis of spectral analyses both strains were found to produce TDA, which could be responsible for their antimicrobial activity. Our findings demonstrated that hydrozoans and red alga inhabiting the Peter the Great Bay of the Sea of Japan harbor diverse microbial communities with a high proportion of bacteria capable of antimicrobials production, including TDA-producing strains which are conceivable to be of importance for chemical protection of their host-hydrobionts and bacterial relationships in marine shallow environments.

Keywords: Bacterial diversity; hydrobionts; antimicrobial activity; *Phaeobacter*; *Pseudovibrio*; tropodithietic acid.

1. INTRODUCTION

There has been increasing interest in the study of microbial communities that are associated with invertebrates and plants dwelling in marine shallow habitats. Coastal ecosystems are known to be of biological significance, harboring a lot of hydrobionts species and providing favourable conditions for marine biota nursery and bioremediation activities. Marine invertebrates and plants serve as unique natural accumulators of specific microbial communities. External envelopes and internal tissues of hydrobionts enriched by nutrient compounds and adhesive substances, polysaccharides, proteins, glycolipids can be considered to provide auspicious conditions for colonization, attachment and activity of associated microorganisms. Microbial associants of marine biota have proven to be a rich source of biologically active substances with antimicrobial, cytotoxic, or antineoplastic activities which have biotechnological and pharmaceutical application [1, 2, 3, 4]. At the same time, our knowledge of the diversity of marine hydrobionts associated bacteria and their biological activity, interrelations between bacteria and host organism is still limited. It is becoming important to study microbial diversity from unexplored or known marine resources, and to search potential microorganisms producing bioactive metabolites. It should be noted that some bacteria play an increasingly significant role as applied to the biological control of aquacultures. This study is focused on the cultivable heterotrophic bacteria isolated from colonial hydrozoan's and red alga *Polysiphonia* sp. specimens collected from the Sea of Japan seashore where being common and abundant they form the fouling on the rocks, mussels, clams and other underwater substrates. The study was aimed to evaluate potential of associated bacteria for the production of biologically active metabolites.

2. MATERIAL AND METHODS

2.1. Isolation and Cultivation of Microorganisms

Fifty-five strains were isolated from colonial unidentified hydrozoan's (class *Hydrozoa*) and eighty strains were isolated from red alga *Polysiphonia* sp. (*Polysiphonia* Greville, 1823, family *Rhodomelaceae*) specimens, which were collected from two different shallow locations of the Peter the Great Bay, the Sea of Japan, Russia, in October 2011 and September 2015, respectively. Isolation and phenotypic characterization of microorganisms were carried out as described previously [5]. The strains were stored

at –80°C in Marine Broth 2216 supplemented with 30% glycerol and deposited to the Collection of Marine Microorganisms, KMM, G.B. Elyakov Pacific Institute of Bioorganic Chemistry, Vladivostok, Russia.

2.2. PCR Amplification of 16S rRNA Genes and Sequencing

Single colonies of isolates were added to 50 µl of autoclaved ultra purified water, and incubated for 15 min at 98 °C. The mixture was centrifuged at 13000 g for 3 min, and 3 µl of the crude extracts was used in the PCR. Partial 16S rRNA gene sequences were amplified using the universal primers 27f (5'-AGAGTTTGATCMTGGCTCAG-3') and 15r (5'-AAGGAGGTGATCCARCCGCA-3'), generating a PCR product of c. 1300-1400 bp. The following PCR conditions were applied: initial denaturation at 95 °C for 15 min, followed by 30 cycles of 94 °C for 1 min, 55 °C for 1 min and 72 °C for 2 min, and a final elongation step of 72 °C for 10 min. The reaction mixture (50 µl) contained DreamTaq™ Green PCR Master Mix (2X) 25 µl (Fermentas), 27f forward primer 5 µl (5 µmol l⁻¹) (Evrogen, Russia), 15r reverse primer 5 µl (5 µmol l⁻¹) (Evrogen, Russia), 12 µl molecular biology grade water and 3 µl DNA. The PCR products were analysed by agarose gel electrophoresis. The purified PCR fragments were sequenced using the ABI PRISM 3130xl Genetic Analyzer (Applied Biosystems) and by the Big Dye v. 3.1 sequencing kit (Applied Biosystems).

2.3. Phylogenetic Analysis

Phylogenetic and molecular evolutionary analyses were conducted using Molecular Evolutionary Genetics Analysis (MEGA, version 6.0.) [6]. Twenty five nucleotide sequences generated in this study have been deposited in the DDBJ/EMBL/GenBank database under the accession numbers from LC230094 to LC230118 as indicated in the phylogenetic trees (Fig. 1, 2).

2.4. Extraction and Isolation of Antimicrobial Compounds

Two strains H8 and rh17 were grown in Marine Broth 2216 with shaking for 30 h. Bacterial cells were removed by centrifugation and the pH of the supernatants was adjusted to 3.5. Equal volume ethyl acetate with 1% formic acid were added twice to the cell-free supernatant (CFS) and incubated shaking at 200 rpm for 30 min followed by centrifugation. Subsequently, the upper layer was removed, washed with H₂O, dried over Na₂SO₄ and vacuum concentrated at 40 °C until dry. The extracts obtained were

examined for the presence of antimicrobial activities at a concentration of 1 mg/ml by an agar diffusion method as described previously [7].

2.5. Instrumental Analysis

UV spectra were recorded on a UV spectrophotometer CECIL, CE 7250, 7000 series (England) in methanol. Liquid chromatography-mass spectrometry, including high resolution, (LC-HR-MS) were recorded on an HPLC Agilent 1200 Series coupled to a maXis impact II (Bruker) spectrometer equipped with electrospray source. HPLC analysis was conducted using a Zorbax Eclipse XDB C₁₈ column (1.0 mm ID, 150 mm, 3.5 µm). The system was operated at a flow rate of 0.1 mL/min with solvents A/B. The gradient was started at 10% B and increased to 95% B within 25.0 min, then to 100% in 1.0 min, keeping this for 10.0 min, returning to 10% B in 2.0 min, and equilibrating for the next sample in 3.0 min (total runtime 41.0 min).

2.6. Antimicrobial Assay

The inhibitory activity against indicator bacteria, *Escherichia coli* K-12 CL588, *Enterococcus faecium* CIP 104105, *Staphylococcus aureus* CIP 65.8^T, *Staphylococcus epidermidis* CIP 81.55^T, *Bacillus subtilis* CIP 52.65^T, *Xanthomonas* sp. pv. *badrii* LMG 546, *Candida albicans* KMM 455 obtained from the respective Culture Collections was tested by an agar diffusion method as described previously [5]. All tests were carried out in three independent experiments.

3. RESULTS AND DISCUSSION

3.1. Diversity of Hydrozoa and Red Alga *Polysiphonia* sp. Associated Bacterial Isolates

Fifty-five strains were isolated from colonial hydrozoan's (class *Hydrozoa*) specimen and investigated phenotypically, and selected strains were studied phylogenetically. On the basis of 16S rRNA gene sequence analysis the hydrozoan's isolates were affiliated to twelve genera *Pseudoalteromonas*, *Shewanella*, (класс *Gammaproteobacteria*); *Phaeobacter*, *Sphingorhabdus*, *Aurantimonas*, *Labrenzia* (класс *Alphaproteobacteria*); *Hydrogenophaga* (класс *Betaproteobacteria*); *Streptomyces*, *Demequina*, *Cellulomonas* (тип *Actinobacteria*); *Kriegella*, *Maribacter* (phylum *Bacteroidetes*) (Fig. 1).




Fig. 1. A neighbor-joining phylogenetic tree based on the 16S rRNA gene sequences available

from the GenBank/EMBL/DDBJ databases (accession numbers are given in parentheses) showing

relationship of *Hydrozoa* associated isolates and related bacteria. Bootstrap values based on 1000 replications are given as percentages at the branching points. Numbers indicate percentages greater than 50%. Bar, 0.02 substitutions per nucleotide position.

Eighty strains were isolated from red alga *Polysiphonia* sp. specimens. All isolates were investigated phenotypically, and selected strains were studied phylogenetically. Bacteria associated with red alga *Polysiphonia* sp. were assigned to twelve genera *Pseudovibrio*, *Labrenzia*, *Sphingomonas* (class *Alphaproteobacteria*); *Cobetia*, *Shewanella*, *Vibrio* (class *Gammaproteobacteria*); *Microbacterium*, *Dietzia*, *Agrococcus*, *Streptomyces* (phylum *Actinobacteria*); *Bacillus*, *Paenibacillus* (phylum *Firmicutes*) (Fig. 2).

Phylogenetic analysis revealed most isolates from both specimens shared 98-99% sequence similarity to recognized species, including those recovered previously from marine sources. Bacterial compositions associated with hydrozoans and red algae were diverse and different to each other at the generic level. Strains of three genera *Shewanella*, *Labrenzia* and *Streptomyces* occurred in both specimens (Fig. 1, 2), and some of *Shewanella* and *Labrenzia* strains belonged to the same phyletic groups whereas *Streptomyces* strains clustered with different phylogenetic lineages of the genus *Streptomyces* separately for each of hydrobionts. Hydrozoan associated bacterial community comprised *Pseudoalteromonas* and *Shewanella* members as abundant and phylogenetically diverse at the species level whereas the isolates within each of the genera *Phaeobacter*, *Sphingorhabdus*, *Aurantimonas*, *Labrenzia* were represented by the homogeneous 16S rRNA gene sequence types. *Hydrogenophaga* (класс *Betaproteobacteria*) and *Bacteroidetes* members were found among bacteria isolated from the hydrozoan's specimen.




Fig. 2. A neighbor-joining phylogenetic tree based on the 16S rRNA gene sequences available from the GenBank/EMBL/DDBJ databases (accession numbers are given in parentheses) showing relationship of isolates associated with red alga *Polysiphonia* sp. and related bacteria. Bootstrap

values based on 1000 replications are given as percentages at the branching points. Numbers indicate percentages greater than 50%. Bar, 0.02 substitutions per nucleotide position.

Red alga associated bacteria belonging to the genera *Sphingomonas*, *Bacillus*, *Paenibacillus* and *Labrenzia* were characterized by the phylogenetic diversity. A group of strains rh45, rh52, rh44, and strain rh62 showed 16S rRNA gene sequence identities less than 97-98 % to the members of the genera *Labrenzia* and *Streptomyces*, respectively, and therefore, they may represent novel species according to the criteria stated by Stackebrandt and Ebers [8].

The results obtained for the taxonomic compositions are mainly corroborated with the occurrence of many genera found in microbial communities associated with marine algae [9,10,11], hydrozoans [12,13], bryozoans [14], sponges [15,16,17], corals [18] collected from various sea habitats. It was summarized in the review of Goecke et al. [10] that bacteria isolated from *Polysiphonia* spp. and unidentified red algae belonged to the genera *Shewanella* (*Proteobacteria*), *Zobellia*, *Lacinutrix*, *Maribacter* (phylum *Bacteroidetes*), *Luteolibacter* (class *Verrucomicrobiae*); and *Alphaproteobacteria* and *Bacteroidetes* members in microbial associations with macroalgae have being observed as constant components. Interrelations between macroalgae and some species of associated bacteria were noted to be highly specific [10]. At the same time our knowledge regarding microbial communities of hydrozoans is very restricted. Stabili et al. [12] have reported luminous bacteria of the genus *Vibrio* in associations with hydrozoan and bryozoan species. Di Camillo et al. [31] found that epibiontic bacteria of *Ectopleura crocea* (*Cnidaria*, *Hydrozoa*) were dominated by members of *Comamonadaceae* and *Flavobacteriaceae*.

3.1. Antimicrobial Activity of *Hydrozoa* and Red Alga *Polysiphonia* sp. Bacterial Isolates

All isolates were tested by the agar diffusion method for their antimicrobial activity. Screening has resulted in 22 hydrozoan's strains and 24 those from red alga (40 % and 30 % of total, respectively) with inhibitory activity against from two to seven indicator strains, mainly, Gram-positive *Staphylococcus aureus*, *Enterococcus faecium*, *Bacillus subtilis*, *Candida albicans* and Gram-negative phytopathogenic

Xanthomonas sp. pv. *badrii*, and less against *Staphylococcus epidermidis* and *Escherichia coli* (Tables 1, 2). Most active hydrozoan's strains were phylogenetically close to *Phaeobacter* and *Streptomyces* members. Streptomycetes constituting about half of active strains contributed significantly to the overall antimicrobial activity. They grouped with recognized species of the genus *Streptomyces*, *S. fulvissimus* and *S. angustmyceticus* which have been reported to be producers of antibiotics, valinomycin and angustmycin, respectively. Strains having moderate or weak activity were found among representatives of the genera *Pseudoalteromonas*, *Shewanella*, *Sphingorhabdus*, *Cellulomonas*. Strains belonging to *Phaeobacter*, *Pseudoalteromonas*, *Shewanella* and *Streptomyces* are well-known antagonists, whereas *Sphingorhabdus* or *Cellulomonas* are rarely mentioned bacteria in this context. Hydrozoan's active strains belonging to the genus *Phaeobacter* clustered all together with *Phaeobacter inhibens* sharing gene sequence similarity of 99%. The genus *Phaeobacter* belongs to the *Roseobacter* lineage (class *Alphaproteobacteria*) which has been reported to be the essential component of microbial communities colonizing marine eukaryotes [19, 20]. The genus *Phaeobacter* was created by Martens et al. [21] as a result of reclassification of the *Roseobacter gallaeciensis* isolated from a rearing of the scallop *Pecten maximus* [22], and description of a new species *P. inhibens* from surface water of a tidal mud flat in the German Wadden Sea. During recent years *Phaeobacter* strains have attracted attention due to their ability to produce biologically active compounds [23, 24, 25], being of importance as probiotic agents in marine aquaculture [26] and as causative bleaching diseases in marine red algae [27].

Most active strains-associants of red alga *Polysiphonia* sp. were placed within the genera *Pseudovibrio*, *Paenibacillus*, *Bacillus* and *Streptomyces* sharing 99% gene sequence similarity to previous recognized species. *Streptomyces* strains occurred abundantly in both hydrobionts specimens and showed remarkable antibacterial properties. Unlike hydrozoan's bacterial consortium most *Streptomyces* strains from red alga were phylogenetically related to *S. albidoflavus* which also includes strains producing bioactive compounds [28]. Members of the genera *Bacillus*, *Paenibacillus*, *Streptomyces* from marine sources have been reported to be of importance as producers of bioactive compounds [3,29,30]. In previous studies we have reported

172 **Table 1. Antimicrobial activity of *Hydrozoa* associated strains.**

Strain	Nearest phylogenetic relative	Similarity (%)	Inhibitory zone (mm)						
			<i>B. subtilis</i>	<i>S. aureus</i>	<i>S. epidermidis</i>	<i>E. faecium</i>	<i>E. coli</i>	<i>Xanthomonas</i> sp. pt. <i>badrii</i>	<i>C. albicans</i>
H1	<i>Pseudoalteromonas hodoensis</i>	99	W*	-	-	W	-	-	12
H2	<i>Shewanella aestuarii</i>	98	-	-	W	-	-	-	12
H4	<i>Pseudoalteromonas tetraodonis</i>	99	W	-	-	W	-	-	12
H5	<i>Pseudoalteromonas tetraodonis</i>	99	-	-	-	W	-	-	12
H6	<i>Phaeobacter inhibens</i>	99	20	13	15	16	-	20	15
H7	<i>Phaeobacter inhibens</i>	99	17	15	17	16	-	21	15
H8	<i>Phaeobacter inhibens</i>	99	14	12	15	12	12	15	w
H9	<i>Streptomyces fulvissimus</i>	99	20	20	-	25	-	14	20
H10	<i>Streptomyces angustmyceticus</i>	98	15	22	-	12	-	15	20
H11	<i>Streptomyces fulvissimus</i>	98	18	22	-	18	-	18	21
H12	<i>Streptomyces fulvissimus</i>	98	25	16	21	15	-	15	17

H13	<i>Streptomyces fulvissimus</i>	100	20	22	-	-	-	15	20
H14	<i>Streptomyces fulvissimus</i>		15	16	12	W	15	12	21
H15	<i>Streptomyces fulvissimus</i>		14	15	20	15	12	12	20
H16	<i>Streptomyces fulvissimus</i>	100	15	16	15	W	12	20	21
H33	<i>Streptomyces fulvissimus</i>		W	12	13	25	12	12	20
H36	<i>Shewanella basaltis</i>	98	-	-	-	-	-	W	20
H38	<i>Pseudoalteromonas hodoensis</i>	99	-	-	W	-	-	12	18
H40	<i>Streptomyces fulvissimus</i>	99	W	-	12	-	12	12	22
H41	<i>Streptomyces fulvissimus</i>		W	-	12	-	12	12	20
H48	<i>Sphingorhabdus flavimaris</i>	99	W	W	12	-	-	-	W
H50	<i>Cellulomonas aerilata</i>	98	-	W	W	-	-	-	-

173

174 *W, weak activity, inhibition zones were observed as zones with reduced growth.

175

176

177 **Table 2. Antimicrobial activity of strains associated with red alga *Polysiphonia* sp.**

Strain	Nearest phylogenetic relative	Similarity (%)	Inhibitory zone (mm)						
			<i>B. subtilis</i>	<i>S. aureus</i>	<i>S. epidermidis</i>	<i>E. faecium</i>	<i>E. coli</i>	<i>Xanthomonas</i> sp. pt. <i>badrii</i>	<i>C. albicans</i>
rh6	<i>Streptomyces albidoflavus</i>	99	18	15	-	15	13	14	12
rh10	<i>Pseudovibrio ascidiaceicola</i>	99	-	-	-	W ⁺	12	-	-
rh11	<i>Pseudovibrio ascidiaceicola</i>	99	W	W	-	-	12	15	-
rh12	<i>Pseudovibrio ascidiaceicola</i>	99	15	16	-	15	W	28	14
rh16	<i>Pseudovibrio ascidiaceicola</i>	99	15	W	-	w	W	15	12
rh17	<i>Pseudovibrio ascidiaceicola</i>	99	15	W	-	15	W	15	12
rh20	<i>Streptomyces albidoflavus</i>	99	15	15	W	w	W	12	17
rh21	<i>Streptomyces albidoflavus</i>	99	16	15	12	15	12	15	18
rh22	<i>Streptomyces albidoflavus</i>	99	18	15	W	w	12	12	18

rh23	<i>Streptomyces albidoflavus</i>	99	-	-	W	15	-	-	12
rh30	<i>Bacillus murimartini</i>		-	12	12	14	-	-	-
rh31	<i>Streptomyces albidoflavus</i>	99	14	16	12	18	12	13	12
rh62	<i>Streptomyces intermedius</i>	93	14	15	W	18	12	14	15
rh63	<i>Streptomyces intermedius</i>		18	15	-	20	-	14	12
rh44	<i>Labrenzia marina</i>		-	12	-	15	-	15	-
rh45	<i>Labrenzia marina</i>	97	-	12	W	-	W	15	-
rh50	<i>Agrococcus jenensis</i>	99	-	14	-	15	11	12	12
rh56	<i>Paenibacillus xylanexedens</i>	99	15	20	-	20	-	12	-
rh57	<i>Paenibacillus terrae</i>	99	-	15	-	20	-	12	12
rh74	<i>Paenibacillus xylanexedens</i>		18	20	-	22	-	18	W
rh77	<i>Paenibacillus xylanexedens</i>		W	14	-	20	-	15	18
rh35	<i>Shewanella pacifica</i>	99	-	-	-	w	W	-	12
rh38	<i>Bacillus murimartini</i>	98	-	15	-	-	-	15	-

rh58	<i>Bacillus murimartini</i>	-	15	-	-	-	15	-
------	-----------------------------	---	----	---	---	---	----	---

178 *W, weak activity, inhibition zones were observed as zones with reduced growth.

bioactive substances produced by *Bacillus*, *Paenibacillus*, and *Saccharothrix* strains associated with ascidian *Halocynthia aurantium* and mollusk *Anadara broughtoni* inhabiting in the Sea of Japan seashore [5,31,32,33].

Bacteria of the genus *Pseudovibrio* (family *Rhodobacteriaceae*, class *Alpharroteobacteria*), to which red alga active isolates belonged, have been reported to be abundant and ubiquitous bacteria associated with diverse marine dwellers, sponges [15,16,17,34], ascidians [35], corals [18] or red alga [36]. In the present study *Pseudovibrio* strains with antimicrobial activity sharing 99% gene sequence similarity positioned all on the line of *Pseudovibrio ascidiaceicola*, strains of which were isolated from ascidians *Polycitor proliferus* and *Botryllidae* sp. collected from a Pacific Ocean beach, the Boso peninsula, Japan [35]. *Pseudovibrio* strains have been reported to produce antimicrobial compounds, tropodithietic acid, TDA, [36], the red pigment heptylprodigiosin [37], polypeptide pseudovibrocin [38].

Comparing microbial associations of hydrozoans and red alga specimens it can be concluded that each of hydrobionts harbored a certain bacterial composition. Despite the taxonomic differences they are united by the occurrence of representatives of *Phaeobacter* and *Pseudovibrio* which are known TDA-producing bacteria. TDA is a disulfide tropolone, which can exist in a tautomeric form with thiotropocin (Fig. 3) and exhibits antimicrobial activity against a broad spectrum of microorganisms [23,20,39]. TDA was firstly isolated from a soil bacterium *Pseudomonas* species [40], and subsequently from marine *Caulobacter* sp. [41] and from the *Roseobacter* clade bacteria including *Ruegeria* sp. and *Phaeobacter inhibens* [23,26,42,43].




Fig. 3. The chemical structure of TDA

Strains *Phaeobacter* sp. H8 and *Pseudovibrio* sp. rh17 were selected to be examined for the production of TDA, which could contribute their antimicrobial activity. Antimicrobial activities against Gram-positive

indicatory strains *B. subtilis*, *S. aureus* and *E. faecium* were revealed in ethyl acetate extracts of CFS of these strains (Fig. 4); very weak activity was observed towards Gram-negative *E. coli* and *Xanthomonas* sp. pv. *badrii*, and no activity against *S. epidermidis* and *C. albicans*.




Fig. 4. In vitro antimicrobial activity assay of ethyl acetate extracts obtained from strains *Phaeobacter* sp. H8 and *Pseudovibrio* sp. rh17 against *Bacillus subtilis* (a) and *Staphylococcus aureus* (b).

The UV spectra of active ethyl acetate extracts exhibited broad strong UV absorption bands at 303 nm, a shoulder at 356 nm and a weak broad absorption around 450 nm, that could be superimposed on the spectrum of thiotropocin or tropodithietic acid [24]. To further confirm the TDA production by strains H8 and rh17, active ethyl acetate extracts were redissolved in 10 mL 85% acetonitrile/15% MilliQ water and analyzed by liquid chromatography-mass spectrometry, including high resolution, (LC-HR-MS). Separation was performed on a Zorbax Eclipse XDB C18 column using a water-acetonitrile gradient solvent system, with both water (solvent A) and acetonitrile (solvent B) containing 0.1% formic acid. The peaks representing the TDA in the extracts of strains H8 and rh17 were t_R =17.8 and 17.2 min, respectively (Fig. 5, 6).





Fig. 5. LC-MS analysis of TDA produced by the strain *Phaeobacter* sp. H8: chromatographic profile (at the top) and corresponding MS spectra in positive ion mode (at the bottom).





Fig. 6. LC-MS analysis of TDA produced by the strain *Pseudovibrio* sp. rh17: chromatographic profile (at the top) and corresponding MS spectra in positive ion mode (at the bottom).

LC-HR-MS analysis confirmed the presence of TDA in the extracts of strains H8 and rh17 based on the unique accurate mass of the $[M+H]^+$ ion at m/z 212.9671 and 212.9678, respectively, (calcd. for $C_8H_4O_3S_2$

m/z 212.9675). To our knowledge, this is the first report concerning TDA production by bacteria associated with hydrozoans and red alga from the Peter the Great Bay, the Sea of Japan.

4. CONCLUSION

Overall our findings showed that hydrobionts studied harbored taxonomically diverse microbial communities with specific composition for each of them. A large proportion of associated bacteria was found to exhibit antimicrobial activities and can be considered as a potential source of bioactive compounds. This study provides evidence that TDA-producing strains belonging probably to *Phaeobacter inhibens* and *Pseudovibrio ascidiaceicola* occurred abundantly in microbial associations colonizing hydrozoans and red alga *Polysiphonia* sp., respectively, in the Sea of Japan seashore. TDA and other antimicrobials produced by bacteria-associants are conceivable to provide chemical protection of their host-organisms and serve for bacterial relationships in marine shallow environments.

REFERENCES

1. Li Z. Advances in marine microbial symbionts in the China Sea and related pharmaceutical metabolites. *Mar. Drugs*. 2009;7:113-129.
2. Hughes CC, Fenical W. Antibacterials from the sea. *Chem. Eur. J*. 2010;16:12512-12525.
3. Debbab A, Aly AH, Lin WH, Proksch P. Bioactive compounds from marine bacteria and fungi. *Microb. Biotech*. 2010;3:544-563.
4. Fuerst JA. Diversity and biotechnological potential of microorganisms associated with marine sponges. *Appl. Microbiol. Biotechnol*. 2014;98:7331-7347.
5. Romanenko LA, Uchino M, Kalinovskaya NI, Mikhailov VV. Isolation, phylogenetic analysis and screening of marine mollusc-associated bacteria for antimicrobial, hemolytic and surface activities. *Microbiol. Res*. 2008;163:633-644.
6. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol*. 2013;30:2725-2729.

- 271 7. Romanenko LA, Uchino M, Frolova GM, Tanaka N, Kalinovskaya NI, Latyshev N, *et al.* *Sphingomonas*
272 *molluscorum* sp. nov., a novel marine isolate with antimicrobial activity. *Int. J. Syst. Evol. Microbiol.*
273 2007;57:358-363.
- 274 8. Stackebrandt E, Ebers J. Taxonomic parameters revisited: tarnished gold standards. *Microb. Today.*
275 2006;45:153-15.
- 276 9. Egan S, Thomas T, Holmstrom C, Kjelleberg S. Phylogenetic relationship and antifouling activity of
277 bacterial epiphytes from the marine alga *Ulva lactuca*. *Environ. Microbiol.* 2000;2:343-347.
- 278 10. Goecke F, Labes A, Wiese J, Imhoff JF. Chemical interactions between marine macroalgae and
279 bacteria. *Mar. Ecol. Prog. Ser.* 2010;409:267-300.
- 280 11. Nylund GM, Persson F, Lindegarth M, Cervin G, Hermansson M, Pavia H. The red alga
281 *Bonnemaisonia asparagoides* regulates epiphytic bacterial abundance and community composition
282 by chemical defence. *FEMS Microbiol. Ecol.* 2012;71:84-93.
- 283 12. Stabili L, Gravili C, Tredici SM, Piraino S, Talá A, Boero F, *et al.* Epibiotic *Vibrio* luminous bacteria
284 isolated from some hydrozoa and bryozoa species. *Microb. Ecol.* 2008;56:625-636.
- 285 13. Di Camillo CG, Luna GM, Bo M, Giordano G, Corinaldesi C, Bavestrello G. Biodiversity of prokaryotic
286 communities associated with the ectoderm of *Ectopleura crocea* (Cnidaria, Hydrozoa). *PLoS ONE.*
287 2012;7(6): e39926. doi:10.1371/journal.pone.0039926
- 288 14. Heindl H, Wiese J, Thiel V, Imhoff JF. Phylogenetic diversity and antimicrobial activities of bryozoan-
289 associated bacteria isolated from Mediterranean and Baltic Sea habitats. *Syst. Appl. Microbiol.*
290 2010;33:94-104.
- 291 15. Hentschel U, Schmid M, Wagner M, Fieseler L, Gernert C, Hacker J. Isolation and phylogenetic
292 analysis of bacteria with antimicrobial activities from the Mediterranean sponges *Aplysina aerophoba*
293 and *Aplysina cavernicola*. *FEMS Microbiol. Ecol.* 2001;35:305-312.
- 294 16. Muscholl-Silberhorn A, Thiel V, Imhoff JF. Abundance and bioactivity of cultured sponge-associated
295 bacteria from the Mediterranean sea. *Microb. Ecol.* 2008;55:94-106.

- 296 17. Margassery LM, Kennedy J, O’Gara F, Dobson AD, Morrissey JP. Diversity and antibacterial activity
297 of bacteria isolated from the coastal marine sponges *Amphilectus fucorum* and *Eurypon major*. Lett.
298 App. Microbiol. 2012;55:2-8.
- 299 18. Chen Y-H, Kuo J, Sung P-J, Chang Y-C, Lu M-C, Wong T-Y, et al. Isolation of marine bacteria with
300 antimicrobial activities from cultured and field-collected soft corals. World J. Microbiol. Biotechnol.
301 2012;28:3269-3279.
- 302 19. Buchan A, Gonzalez JM, Moran MA. Overview of the marine *Roseobacter* lineage. Appl. Environ.
303 Microbiol. 2005;71:5665-5677.
- 304 20. Brinkhoff T, Giebel H-A, Simon M. Diversity, ecology, and genomics of the *Roseobacter* clade: a short
305 overview. Arch. Microbiol. 2008;189:531-539.
- 306 21. Martens T, Heidorn T, Pukall R, Simon M, Tindall BJ, Brinkhoff T. Reclassification of *Roseobacter*
307 *gallaeciensis* Ruiz-Ponte et al. 1998 as *Phaeobacter gallaeciensis* gen. nov., comb. nov., description
308 of *Phaeobacter inhibens* sp. nov., reclassification of *Ruegeria algicola* (Lafay et al. 1995) Uchino et
309 al. 1999 as *Marinovum algicola* gen. nov., comb. nov., and emended descriptions of the genera
310 *Roseobacter*, *Ruegeria* and *Leisingera*. Int. J. Syst. Evol. Microbiol. 2006;56:1293-1304.
- 311 22. Ruiz-Ponte C, Cilia V, Lambert C, Nicolas JL. *Roseobacter gallaeciensis* sp. nov., a new marine
312 bacterium isolated from rearings and collectors of the scallop *Pecten maximus*. Int. J. Syst. Bacteriol.
313 1998;48:537-542.
- 314 23. Brinkhoff T, Bach G, Heidorn T, Liang L, Schlingloff A, Simon M. Antibiotic production by a
315 *Roseobacter* clade affiliated species from the German Wadden Sea and its antagonistic effects on
316 indigenous isolates. Appl. Environ. Microbiol. 2004;70:2560-2565.
- 317 24. Bruhn JB, Nielsen KF, Hjelm M, Hansen M, Bresciani J, Schulz S, et al. Ecology, inhibitory activity,
318 and morphogenesis of a marine antagonistic bacterium belonging to the *Roseobacter* Clade. Appl.
319 Environ. Microbiol. 2005;71:7263-7270.

- 320 25. Prado S, Montes J, Romalde JL, Barja JL. Inhibitory activity of *Phaeobacter* strains against
321 aquaculture pathogenic bacteria. *Int. Microbiol.* 2009;12:107-114.
- 322 26. Porsby CH, Nielsen KF, Gram L. *Phaeobacter* and *Ruegeria* species of the *Roseobacter* clade
323 colonize separate niches in a Danish Turbot (*Scophthalmus maximus*)-rearing farm and antagonize
324 *Vibrio anguillarum* under different growth conditions. *Appl. Environ. Microbiol.* 2008;74:7356-7364.
- 325 27. Fernandes N, Case RJ, Longford SR, Seyedsayamdost MR, Steinberg PD, Kjelleberg S, *et al.*
326 Genomes and virulence factors of novel bacterial pathogens causing bleaching disease in the marine
327 red alga *Delisea pulchra*. *PLoS ONE.* 2011;6:e27387.
- 328 28. Roy RN, Laska S, Sen SK. Dibutyl phthalate, the bioactive compound produced by *Streptomyces*
329 *albidoflavus* 321.2. *Microbiol. Res.* 2006;161:121-126.
- 330 29. Jensen PR, Mincer TJ, Williams PG, Fenical W. Marine actinomycete diversity and natural product
331 discovery. *Antonie van Leeuwenhoek.* 2005;87:43-48.
- 332 30. Bull AT, Stach JEM. Marine actinobacteria: new opportunities for natural product search and
333 discovery. *Trends in Microbiology.* 2007;15:491-499.
- 334 31. Romanenko LA, Kalinovskaya NI, Mikhailov VV. Taxonomic composition and biological activity of the
335 microorganisms associated with marine ascidian *Halocynthia aurantium*. *Biologiya Morya. (Russ. J.*
336 *Mar. Biol.)* 2001;27:291-295.
- 337 32. Kalinovskaya NI, Kuznetsova TA, Ivanova EP, Romanenko LA, Voinov VG, Huth F, Laatsch H.
338 Characterization of surfactin-like cyclic depsipeptides synthesized by *Bacillus pumilus* from ascidian
339 *Halocynthia aurantium*. *Mar. Biotechnol.* 2002;4:179-188.
- 340 33. Kalinovskaya NI, Kalinovskiy AI, Romanenko LA, Dmitrenok PS, Kuznetsova TA. New angucyclines
341 and antimicrobial diketopiperazines from the marine mollusk-derived actinomycete *Saccharothrix*
342 *espanaensis* An 113. *Nat. Prod. Comm.* 2010;5:597-602.

343 34. O'Halloran JA, Barbosa TM, Morrissey JP, Kennedy J, O'Gara F, Dobson ADW. Diversity and
344 antimicrobial activity of *Pseudovibrio* spp. from Irish marine sponges. J. Appl. Microbiol.
345 2011;110:1495-1508.

346 35. Fukunaga Y, Kurahashi M, Tanaka K, Yanagi K, Yokota A, Harayama S. *Pseudovibrio ascidiaceicola*
347 sp. nov., isolated from ascidians (sea squirts). Int. J. Syst. Evol. Microbiol. 2006;56:343-347.

348 36. Penesyan A, Tebben J, Lee M, Thomas T, Kjelleberg S, Harder T, et al. Identification of the
349 antibacterial compound produced by the marine epiphytic bacterium *Pseudovibrio* sp. D323 and
350 related sponge-associated bacteria. Mar. Drugs. 2011;9:1391-1402.

351 37. Sertan-de Guzman AA, Predicala RZ, Bernardo EB, Neilan BA, Elardo SP, Mangalindan GC et al.
352 *Pseudovibrio denitrificans* strain z143-1, a heptylprodigiosin-producing bacterium isolated from a
353 Philippine tunicate. FEMS Microbiol. Lett. 2007;277:188-196.

354 38. Vizcaino M, Moeller P, Morris P. The Potential for the Marine Bacterium *Pseudovibrio denitrificans* to
355 Produce Novel Antibiotics. Available:[https://www.musc.edu/mbes-](https://www.musc.edu/mbes-lig/MBESOpenHouse/2010/abstracts/Vizcaino-SROH_July_16_2010.pdf)
356 [lig/MBESOpenHouse/2010/abstracts/Vizcaino-SROH_July_16_2010.pdf](https://www.musc.edu/mbes-lig/MBESOpenHouse/2010/abstracts/Vizcaino-SROH_July_16_2010.pdf).

357 39. Crowley SP, O'Gara F, O'Sullivan O, Cotter PD, Dobson ADW. Marine *Pseudovibrio* sp. as a Novel
358 Source of Antimicrobials. Mar. Drugs. 2014;12:5916-5929.

359 40. Kintaka K, Ono H, Tsubotani S, Harada S, Okazaki H. Thiotropocin, a new sulfur-containing 7-
360 membered-ring antibiotic produced by a *Pseudomonas* sp. J. Antibiot. 1984;37:1294-1300.

361 41. Kawano Y, Asada M, Inoue M, Nakagomi K, Oka S, Higashihara T. Biological activity of thiotropocin
362 produced by marine bacterium, *Caulobacter* sp. PK654. J. Mar. Biotechnol. 1998;6:49-52.

363 42. Geng H, Belas R. Expression of tropodithietic acid biosynthesis is controlled by a novel autoinducer. J.
364 Bacteriol. 2010;192:4377-4387.

365 43. Porsby CH, Webber MA, Nielsen KF, Piddock LJV, Gram L. Resistance and Tolerance to Tropodithietic
366 Acid, an Antimicrobial in Aquaculture, Is Hard To Select? Antimicrob. Agents Chemother. 2011;1332-
367 1337.