

1       **Anti-Coccidiosis Potential of Autoclaveable Antimicrobial Peptides**  
2       **from *Xenorhabdus budapestensis* Resistant to Proteolytic (Pepsin,**  
3       **Trypsin) Digestion based on *in vitro* studies**

4  
5  
6       **András Fodor<sup>1\*</sup>, László Makrai<sup>2</sup>, László Fodor<sup>2</sup>, István Venekai<sup>3</sup>, Ferenc**  
7       **Husvéth<sup>4</sup>, László Pál<sup>4</sup>, Andor Molnár<sup>4</sup>, Károly Dublecz<sup>4</sup>, Csaba Pintér<sup>4^</sup>**  
8       **Sándor Józsa<sup>4^</sup> and Michael G. Klein<sup>5</sup>**

9  
10       <sup>1</sup>*Department of Genetics, Faculty of Science and Informatics, University of Szeged,*  
11       *University of Szeged, Közép fasor 52, Szeged, H-6726, Hungary,*

12       <sup>2</sup>*Department of Microbiology and Infectious Diseases, University of Veterinary Medicine*  
13       *Budapest, Hungary*

14       <sup>3</sup>*Department of Biochemistry, Eötvös Loránd University, Budapest H-1117, Hungary*

15       <sup>4</sup>*Department of Animal Sciences and Animal Husbandry, Georgikon Faculty, University of*  
16       *Pannonia, Széchenyi Street, 11 Keszthely, H-8360 Hungary).*

17       <sup>4^</sup>*Adjunct Emeritus Professor, Georgikon Faculty, University of Pannonia, Széchenyi Street,*  
18       *11 Keszthely, Hungary (The place of the *in vivo* experiment)*

19       <sup>5</sup>*Adjunct Professor, Department of Entomology, The Ohio State University, 1680 Madison*  
20       *Ave., Wooster, OH-44691, USA*

21  
22  
23

---

24  
25

## ABSTRACT

**Aims:** To elucidate the anticoccidial potential of antimicrobial peptides from *Xenorhabdus budapestensis* on both causative pathogens (prokaryotic *Clostridium perfringens* and eukaryotic *Eimeria tenella*). **Objectives:** (1) To establish if the antimicrobial compounds of the cell-free culture media (CFCM) of the entomopathogenic symbiotic bacterium species, *X. budapestensis* DSM 16342 (EMA) and *X. szentirmaii* DSM 16338 (EMC) were active against 13 independent pathogenic isolates of *Clostridium perfringens in vitro*; (2) To create a sterile, autoclaved, bio-preparation called “XENOFood”, for future *in vivo* feeding studies, aimed at determining the efficacy, and side-effects, of EMA and EMC on *C. perfringens* in chickens.

**Study design:** *Clostridium perfringens* samples (LH-1-LH24) were collected from chickens and turkeys, and were deposited in the frozen stock collection of Department of Microbiology and Infectious Diseases, Faculty of Veterinary Science, Szent István University, Budapest, Hungary, where the *in vitro* assays were carried out on 13 of these isolates.

**Place and Duration of Study:** Department of Microbiology and Infectious Diseases, Faculty of Veterinary Science, Szent István University, Budapest, Hungary between September 2013 and February 2014.

**Methodology:** Adaptation of our previously published *in vitro* bioassays for aerobic tests for the anaerobic bacteria *Clostridium perfringens*. When preparing “XENOFood” we benefitted from our experimental data about the heat tolerance and endurance to proteolytic enzymatic digestion of the studied antimicrobial peptides.

**Results:** The studied antimicrobial peptides were heat-stable, trypsin and pepsin resistant. All but one of 13 *C. perfringens* isolates was sensitive to EMA-CFCM. XENOFood (made here) is not toxic for chicken, (unpublished).

**Conclusion:** Since these cell-free cultures killed *E. tenella* cells, but were toxic to permanent chicken liver (LMH) cells, we need to run *in vivo* feeding tests to determine the gastrointestinal (ileac), anti-*Clostridium* and anti-*Eimeria* biological effects of the these heat-, - and proteolysis tolerant antimicrobial peptides.

26  
27  
28  
29  
30

*Keywords: Clostridium perfringens, Xenorhabdus Antimicrobial Peptides; in-vitro Bioassay, Xenofood*

## 1. INTRODUCTION

31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

Multi-drug resistance (MDR) has gradually been increasing in both Gram-positive [1] and Gram-negative [2] pathogenic bacterium species. MDR has always been a phenotypic consequence of sequential accumulation of simultaneously appearing mutations, or the up-take of resistance plasmids harboring mobile genetic elements or genomic islands with resistance genes. These encode for either enzymes capable of destructing the antibiotics, or catalyzing biochemical reactions resulting in inhibition of either binding to, or permeation through, the cellular membrane (CM). The poultry gastro-intestinal (GI) flora is a seed-bed of MDR, as shown by the spectacular on-going evolution in *Enterococcus* [3] [4] [5], in *Clostridium* [6], as well as in *Salmonella* genera [7]. The explanation is that the poultry GI is an ideal “market place” for exchange and horizontally transferring resistance gene –carrying

43 plasmids, and mobile genetic elements, between coexisting bacteria. *Enterococcus*  
44 *cecorum*, for instance, once a simple commensal member of the intestinal microbiota, has  
45 become the causative pathogen of arthritis and osteomyelitis worldwide in chickens, such as  
46 in Hungary [8] and Poland [9]. Evidences of multidrug-resistant plasmid transfer from Gram  
47 positive [10] and Gram negative [11] [12] chicken pathogens via consumed chicken meat to  
48 human pathogens, has been accumulating. Apart from the veterinary aspects, this horizontal  
49 gene transfer is of critical clinical importance.

50 The anaerobic, Gram-positive, *C. perfringens* was first published as a globally threatening  
51 danger by Van Immerseel and his associates, [13] as the causative pathogen of necrotic  
52 enteritis. Since then it has become alarming from both veterinary and human clinical  
53 aspects. The incidence of *C. perfringens*-associated necrotic enteritis in poultry has also  
54 increased in countries that stopped using antibiotic growth promoters. Both the disease and  
55 its subclinical forms are caused by *C. perfringens* type A strains, which produce either the  
56 alpha toxin, (to a lesser extent type C), or both alpha and beta toxins [14]. A few *C.*  
57 *perfringens* type A isolates produce an enterotoxin at sporulation as well, causing disease in  
58 humans, [14].

59 As for the pathogenesis of necrotic enteritis in chickens [15], it is a result of a “joint venture”  
60 the eukaryotic *Eimeria* species and *C. perfringens*, [16] [17]. The lesions and damages in the  
61 gut wall tissues (mainly in the lamina muscularis mucosae and in the lamina mucosa)  
62 provide anaerobic conditions needed for propagation of the toxin-producing *Clostridium*,  
63 especially in the ileum. The *Eimeria* (most frequently) *tenella* infection is usually preceded by  
64 previous unfavorable changes in the GI biota. The latter might be an indirect consequence of  
65 non-appropriate diets which increases the viscosity of the intestinal contents and makes it  
66 predisposed to necrotic enteritis [15]. This important discovery provides an opportunity for  
67 nutrient scientists to help reduce *Clostridium* infections. In other words, the discovery that  
68 the gastrointestinal microbiota could significantly be restructured by nutritional factors,  
69 provides additional opportunities for nutrition scientists working on the problem of coccidiosis  
70 [18] [19] or similar problems such as *Campylobacter jejuni*, [20]

71 *Clostridium perfringens* type A cells release different toxins that causing diseases not only in  
72 chickens, but also in humans. One of them, the necrotic enteritis B-like toxin (NetB), is a  $\beta$ -  
73 barrel pore-forming one, which used to be a candidate vaccine [21]. Another one, called  
74 perfringolysin O (PFO, also called  $\theta$  toxin), is a pore-forming cholesterol-dependent cytolysin  
75 (CDC) [22]. PFO is secreted as a water-soluble monomer that recognizes and binds  
76 membranes via cholesterol. Membrane-bound monomer molecules undergo chemical  
77 structural changes that culminate in the formation of an oligomerized pre-pore complex on  
78 the membrane surface [22]. The pre-pore then undergoes conversion into the bilayer-  
79 spanning pore, playing an important role in so-called gas gangrene progression and necro-  
80 hemorrhagic enteritis in some mammals [22].

81 *Clostridium perfringens* strains which were isolated from epidemic outbreaks of necrotic  
82 enteritis, and were capable of secreting factors that inhibit growth of other (competitor) *C.*  
83 *perfringens* strains, including those isolated from the guts of healthy chickens [23]. This  
84 feature lends a selective virtue to respective NetB-toxin producing virulent strains, the  
85 causative factor of gut lesions. The factor providing this selective virtue to the virulent strains

86 is a novel, chromosomally encoded, heat-labile, trypsin - and proteinase-K sensitive protein  
87 with bacteriocin activity called perfrin [23]. The gene, which can only be found in *C.*  
88 *perfringens* NetB strains and nowhere else, (despite the fact that the NetB is a plasmid  
89 encoded toxin), could be transferred to and expressed in *E. coli*. Theoretically, it may also  
90 happen in the chicken GI at any time, and the recombinant gene product is antibacterial  
91 active at a large pH range [23].

92 Several data from the literature seem to support our opinion that although vaccination is an  
93 effective, but probably not an omnipotent, veterinary tool for controlling Gram-positive MDR  
94 pathogens such as *Clostridia*. The vaccination projects involving *Enterococcus* seem to be in  
95 a promising, but only very experimental stage [24]. (None of the seven respective  
96 publications have recently been available in PubMed include anything on poultry).

97 As for *Clostridia*, the vaccination of chickens against the fatal human pathogen type C  
98 (causing botulism), have fortunately been successful [25]. The vaccination against *C.*  
99 *perfringens* however, although seeming to be not too far from realization, but maybe not in  
100 the near future. The immunization with NetB genetic, or formaldehyde toxoids, seemed to be  
101 the most plausible approach [26], but only the double vaccination (on 3 and 12 days, with  
102 crude supernatant), were effective. Immunization with a single toxin molecule did not give  
103 satisfactory protection to chickens against necrotic enteritis lesions, which probably is not a  
104 realistic option for practical application [27].

105 This observation led Professor Dr. Van Immersee (Universiteit Gent, Belgium) and his  
106 associates to the conclusion that “*immunization with single proteins is not protective against*  
107 *severe challenge. Therefore combinations of different antigens are needed as alternative. In*  
108 *most published studies multiple dosage vaccination regimens were used. It is not a relevant*  
109 *way for practical use in the broiler industry*”, [28]. Some other less pessimistic reports, such  
110 as suggesting the use of *C. perfringens* recombinant proteins in combination with  
111 Montanide™ ISA 71 VG adjuvant as a vaccine [29] or anticoccidial live vaccine [30] have  
112 been noted. Nevertheless, we think that we'd better to accept the opinion of the Expert #1 in  
113 that research field: the vaccination against avian *C. perfringens* type A strains in broiler  
114 chicken is not yet available [28].

115 Consequently, there is a room for working on novel antimicrobials, especially on novel  
116 antimicrobial peptides which might be used to control *C. perfringens* A and also MDR  
117 pathogens in the GI system of broiler chickens. This approach needs a comprehensive  
118 strategy, based on Quantitative Structure – Activity Relation (QSAR) analysis and *in silico*  
119 modeling [31]. Chemical synthesis of modified analogs leading to new antimicrobial agents  
120 with novel modes of action should follow the molecular design to get new antimicrobial  
121 peptides, [31]. The structural design of AMP candidate molecules has aimed at improving  
122 endurance to proteolytic degradation, binding to, and the penetration through cellular  
123 membranes and other biological barriers [32]. This can be achieved by adding modules for  
124 passive or active transport [32]. Another approach is searching for efficient synergisms [33].

125 Another (ever-green) alternative research line is to search for new antimicrobials of  
126 completely novel modes of action in nature. Our research team has been searching for novel  
127 antimicrobials, which are not used in human medicine, are toxic only for chicken pathogens,  
128 but not toxic for the organisms to be protected. We expect to find the best candidates among

129 the natural antimicrobial peptides (AMPs), synthesized by the obligate bacterial symbionts  
130 (EPB) of entomopathogenic nematodes (EPN) [34]. These EPB-released AMPs are  
131 evolutionary products developed under severe selective pressure, and comprise a powerful  
132 chemical arsenal against a large scale of prokaryotic and eukaryotic organisms. They  
133 provide monoxenic conditions for a given respective EPN / EPB symbiotic complex in  
134 polyxenic (insect gut, soil) conditions. Many EPN-EPB complexes exist, and many AMP  
135 profiles could be determined. Considering that all but one [35] of the known AMPs can be  
136 produced by the bacterium *in vitro*, the EPN/EPB complexes provide a gold mine for  
137 researchers interested in new antimicrobials. The majority of EPB-produced AMPs were  
138 identified in the last 15 years [36] [37] [38] [39]. Each of these evolutionarily designed  
139 antibiotic arsenals has effectively overcome intruders representing a full scale of antibiotic  
140 resistance repertoire in their respective niche. Each EPB-AMP discovered so far is a non-  
141 ribosomal peptide (NRP), synthesized by multi-enzyme thiotemplate mechanisms, using  
142 non-ribosomal peptide synthetases (NRPS), fatty acid synthases (FAS), and / or related  
143 polyketide synthases (PKS), or a hybrid biosynthesis thereof [40]. The biosynthetic enzymes  
144 are encoded by gene clusters [41], determining the biosynthetic pathways.

145 Cabanillasin, from *X. cabanillasii*, exerts of a strong antifungal activity [42]. In our  
146 experiments, the cell-free culture media (CFCM) of *X. cabanillasii* was also extremely toxic  
147 to *Staphylococcus aureus*, *Escherichia coli* and *Klebsiella pneumoniae*, isolated from cows  
148 with mastitis syndromes [43]. In that experiment, the antibacterial activities of the CFCM of  
149 several *Xenorhabdus* species were compared.

150 We found that and those of *X. budapestensis* DSM 16342 (EMA), and *X. szentirmaii* DSM  
151 16338 (EMC) [44] proved far the best. The CFCM of EMA and EMC were also effective  
152 against *S. aureus* MRSA, (Fodor, McGwire and Kulkarni, unpublished). Furthermore, the  
153 CFCM from EMA and EMC also was effective against plant pathogens, including both  
154 prokaryotic *Erwinia amylovora*, *E. carotovora*, *Clavibacter michiganense* and several  
155 *Xanthomonas* species [45] [46] [47] and all tested eukaryotic Oomycetes (*Phytophthora*)  
156 species [42] (Muvevi et al., unpublished). Gualtieri confirmed our data, declaring that *X.*  
157 *szentirmaii* DSM16338 (EMC) was really a source of antimicrobial compounds of great  
158 potential, and he sequenced this strain [48]. One of the products (szentiamide) has been  
159 chemically synthesized [46].

160 We suppose that these antimicrobial peptides act in concert. The idea of a preparing a bio-  
161 product for oral administration to via chicken food, ("XENOFood"), is based on the intention  
162 to benefit from the joint action of cooperating AMP molecules produced by EMA and EMC  
163 cells, not only on a single molecule. We know that the strongest, predominant antibacterial  
164 peptide produced by both EMA and EMC species is fabclavine [51], but there are also others  
165 acting on eukaryotic pathogens as well, especially in EMC [48] [49]. (This is the explanation  
166 why we did not use only EMA CFCM alone, but a mixture of EMA and EMC CFCM instead in  
167 the experiments reported here).

168 Many of our experiments with EMA were repeated in the laboratory of Professor Helge B  
169 Bode (Goethe-Universität, Frankfurt – am – Main, Germany). They confirmed that EMA  
170 CFCM exhibited broad-spectrum bioactivity against *Bacillus subtilis*, *E. coli*, *Micrococcus*  
171 *luteus*, *Plasmodium falciparum*, *Saccharomyces cerevisiae*, *Trypanosoma brucei*, and *T.*

172 *cruzi* [51] as well. They subjected the CFCM from *X. budapestensis* to MALDI-MS analysis  
173 and found altogether 4 isomers of fabclavine, one of which was then purified, and its  
174 structure was determined. The details of biosynthesis were impressively reconstructed by  
175 the authors, but no data about the mode of action has been published so far [51].  
176 Fabclavines are considered a novel class of biosynthesized hybrid peptide–polyketide-  
177 polyamino natural compounds with extremely high antimicrobial potential in both prokaryotic  
178 and eukaryotic pathogen targets, but also with unwanted eukaryotic cell-toxicity. They are  
179 unambiguously the most effective antimicrobial *Xenorhabdus* peptide-products that have  
180 ever been discovered, and they are released by *X. budapestensis* and *X. szentirmaii* [44].  
181 (This is a spectacular example of present-day science, when on group of scientists are  
182 “sowing” while the other ones are “harvesting”).

183 We tested CFCM of EMA and EMC were in 2009 in the McGwire laboratory (Ohio State  
184 University, Columbus, OH, USA) against different targets, and found that, similarly to several  
185 other antimicrobial peptides [52] [53] they exerted apoptotic effects on eukaryotic cells of  
186 *Leishmania donovani*. They were also active against *Candida* sp., and *Phytophthora*  
187 *infestans* (A. Fodor et al., unpublished).

188 Considering that not only prokaryotic, but eukaryotic pathogens also exist, we decided to  
189 continue the “EMA-EMC” project. Coccidiosis is the best example of when a prokaryotic and  
190 a eukaryotic pathogen act together. Dr. Petra Ganas tested both CFCMs on a permanent  
191 chicken liver cell line at the Vet Med University of Vienna, Austria, and found them toxic to  
192 the tissue cultures (Ganas, personal communication, for details, see Discussion), even if the  
193 toxic cell concentration was 1 order of magnitude higher than the bactericide concentration.  
194 These data, and the identification of the most active component (fabclavine), might seem  
195 discouraging for the continuation of the project.

196 However, considering the presence of multidrug resistance, and even pan-resistance,  
197 problems in the GI system of broiler chicken, which may also threaten human health, and the  
198 limitation of vaccinations, we reconsidered it as a potential tool, on the prospects that orally  
199 applied compounds would not be absorbed into the meat of broiler chickens. Prior to *in vivo*  
200 feeding tests we carried out the *in vitro* bioassays presented here, and formulated a chicken  
201 food, Xenofood, to test in the *in vivo* tests. From this aspect, we believe that the results of  
202 this *in vivo* experiment are worthwhile, and our conclusions will be taken into consideration  
203 by coccidiosis specialists.  
204

## 205 **2. MATERIAL AND METHODS**

### 206 **2.1 Bacterium Strains**

207  
208

209 *Clostridium perfringens* NCAIM 1417 strain was obtained from the National Collection of  
210 Agricultural and Industrial Microorganisms –WIPO (of Hungary, Faculty of Food Sciences,  
211 Szent István University Somlói út 14-16 1118 Budapest, Hungary). *Clostridium perfringens*  
212 LH1-LH8; LH11-LH16; LH19, and LH20 are of chicken origin, and LH24 came from a pig;  
213 each has been deposited in the (frozen) stock collection of Department of Microbiology and  
214 Infectious Diseases, University of Veterinary Medicine Budapest, Hungary.



215 *Xenorhabdus* strains, *X. budapestensis* DSM 16342 (EMA), *X. szentirmaii* DSM 16338  
216 (EMC) [44] and *X. bovienii* NYH which had been isolated from the entomopathogenic  
217 nematodes *Steinernema bicornutum* [Tallósi] [54], *S. rarum* and *S. feltiae* HU1 [55], are  
218 originated from the Fodor laboratory, Eötvös University, Budapest, Hungary. EMA and EMC  
219 had also been deposited by us in the DSMZ, (Leibniz Institute Deutsche Sammlung von  
220 Mikroorganismen und Zellkulturen, Braunschweig, Germany) as DSM 16342 and DSM  
221 16338, respectively. *Xenorhabdus nematophila* ATTC 19061, was from Forst Laboratory at  
222 the University of Wisconsin – Milwaukee, USA) and *X. nematophila* DSM 3370 DSMZ,  
223 Braunschweig, Germany). *Steinernema cabanillasii* BP was isolated by us from infective  
224 dauer juveniles from the EPN *S. riobrave*.

## 225 **2.2 Overlay Bioassays for Comparing the Antibacterial Potential of Different** 226 ***Xenorhabdus* Strains**

227 Overlay bioassays for comparing the antibacterial potential of different *Xenorhabdus* strains  
228 (each representing a species), were carried out as previously described [43]. To make sure  
229 that we use the proper bacterium, an earlier experiment was repeated in which we compared  
230 the antibacterial activities of 5 different *Xenorhabdus* strains on *K. pneumoniae*.

231 To determine if the antimicrobial compounds from EMA were effective against *C.*  
232 *perfringens*, an overlay experiment was carried out [43]. To be sure that the intestinal  
233 proteolytic activities would not inactivate our compounds, samples of EMA CFCM were  
234 digested with pepsin, following the professional guidance of our coauthor Professor Ferenc  
235 Husvéth (University of Pannonia, Keszthely, Hungary), while another sample was digested  
236 with trypsin by István Venekei (Eötvös University, Budapest, Hungary)

## 237 **2.3 Agar-Diffusion Assay of EMA CFCM against *Clostridium perfringens* NCAIM 1417** 238 **Laboratory Strain**

239 Agar Diffusion Tests were similarly carried out, as described by [46], but we converted the  
240 method for the anaerobic specimen, *C. perfringens*. An agar diffusion test was conducted as  
241 follows: In a hole at the center of the agar plate, 100  $\mu$ l of EMA CFCM were pipetted and  
242 overlaid with 3 ml of a log phase *C. perfringens* suspension diluted to 1:250 with soft (0.6  
243 V/V%) agar. They were incubated for 24h under anaerobic conditions at 40 °C.

## 244 **2.4 Comparison of the Sensitivities (MID Values) of 13 *C. perfringens* Strains, Isolated** 245 **from Poultry, to Cell-Free Culture Media (CFCM) of *X. budapestensis* (EMA) in Liquid** 246 **Cultures**


### 247 **2.4.1 Determination of MID Values**

248 To quantify the sensitivity of the strains, the maximum inhibiting dilution (MID) values [43]  
249 [56] [46] [47] were determined as below. These studies were carried out in sterile 24-hole  
250 tissue culture plates, with 4 (A-D) rows and 6 (1-6) Columns, in 1 ml final volumes. Each  
251 *Clostridium* strain was used in a different tissue culture plate. Each hole contained 0.5 ml of  
252 2XRCM Reinforced Clostridium Media [57] liquid medium, and 0.5 ml of sterile, diluted EMA  
253 CFCM, with the following distribution: 100, 80, 60, 40, 20 and 0 volume / volume (V/V) % in  
254 column 1, 2, 3, 4, 5, and 6, respectively. There were 50, 40, 30, 20, 10 and 0% V/V final  
255 concentration of EMA CFCM in columns 1, 2, 3, 4, 5, and 6. Each culture in rows A, B and C  
256 were inoculated with loopful of the respective bacteria obtained from three separate colonies  
257 grown on sheep blood agar plates. The holes in row D were not inoculated, and served as  
258 sterile (negative) controls. Columns 6 served did not contain EMA CFCM and served as

259 positive controls. Each 1-ml culture was overlaid by 0.5 ml sterile (freshly autoclaved),  
260 paraffin oil to provide anaerobic conditions. Plates were then incubated at 37°C for 24h and  
261 then scored visually. After 24h culturing, the growing and inhibited cultures could  
262 unambiguously be identified. We considered the concentration as MID where none of the 3  
263 replicates contained visible growth.

#### 264 **2.4.2 Enumeration of *Clostridium perfringens* colony forming units (CFU)**

265 Samples were taken from the first hole in which bacterial proliferation was not visually  
266 detected. 0.5 ml of culture were sucked out cautiously from below the paraffin oil and serial  
267 dilutions were prepared up to  $10^{-5}$ , and 100  $\mu$ l volumes were simultaneously spread onto the  
268 surface of sheep blood agar (by D. László Makrai, see Fig. 1) and Tryptose-Sulfite-  
269 Cycloserine (TSC) agar [59] plates. The latter was designed as a highly selective solid  
270 medium for growing and enumerating *C. perfringens* colony forming units. The TSC allows  
271 virtually complete recovery *C. perfringens*, while it inhibits practically all facultative  
272 anaerobes tested, and is known as being more selective than SFP Agar. Three replicates  
273 were used for each dilution. In preliminary experiments, carried out by András Fodor and  
274 Andor Molnár, both then at the Department of Animal Sciences and Animal Husbandry  
275 (Georgikon Faculty, University of Pannonia, Keszthely, Hungary), TSC plates were  
276 incubated under anaerobic conditions at 40°C, and found the best readability between 48 –  
277 72h. The *C. perfringens* colonies were recognized by colony color and the black reduced  
278 sulfides granules around them, but the color of the agar also gave a kind of qualitative  
279 information (Fig. 1). The colonies used in these preliminary experiments were obtained from  
280 chicken ileal digests, and from the stock collection of Dr. L. Makrai, were reproducibly  
281 counted.



282 .

283 Figure 1 shows the *Clostridium* colonies to be counted on a blood agar plate (Photo: Dr.  
284 László Makrai, (Department of Microbiology and Infectious Diseases, University of  
285 Veterinary Science, Szent István University, Budapest, Hungary).

#### 286 **2.5 Study of the Endurance of the Antimicrobial Compounds in the Cell-Free Culture 287 Media (CFCM) of *X. budapestensis* and *X. szentirmaii* to Proteolytic Degradation**

288 **2.5.1. Trypsin-digested samples** were tested on Gram-positive (*Staph. aureus*) and Gram  
289 negative (*E.coli*) targets in agar diffusion assay, and compared with untreated CFCM  
290 samples. No differences were demonstrated.



291 **2.5.2. Pepsin resistance** was studied as follows: in the center of a Luria Broth plate, a  
292 Millipore filter of 0.22  $\mu\text{m}$  pore size was laid and infiltrated with HCl and pepsin. Then EMA  
293 CFCM was pipetted onto it. The pepsin preparations were prepared by Professor Ferenc  
294 Husvéth. After that the plate was overlaid with a *Pseudomonas aeruginosa* suspension  
295 diluted with soft agar as described [46] [47]. After 24 h incubation at 40 °C, the growth of the  
296 test bacterium lawn was checked.

## 297 **2.6 Preparation of XENOFood**

298  
299 XENOFood: XENOFood contained 5% soy-meal, which had been suspended with equal  
300 amount (w/w) of EMA, and another 5% suspended in equal amount (w/w) of EMC cells  
301 obtained from 5 days-old shaken (2000 rpm) liquid cultures, followed by high-speed  
302 (Sorwall; for 30 minute) centrifugation. The liquid cultures were in 2XLB (DIFCO),  
303 supplemented with meat extract equivalent to the yeast extract. Five days was optimal for  
304 antibiotic production at 25°C under these conditions [43] [45]. It had previously been  
305 discovered that both EMA and EMC grow and produce antibiotics in autoclaved soy-meal  
306 containing some water and yeast extract, or in autoclaved 0.5% w/w yeast (Fodor,  
307 unpublished). Therefore the original chicken food served as a semi-solid culture media for  
308 the *Xenorhabdus* cells. Both the separate EMA and EMC culturing semi-solid chicken food  
309 that we (Dr. László Pál) prepared daily were incubated under sterile conditions for another  
310 five days. Then the EMA and EMC culture media were combined, autoclaved (20 min, 121  
311 °C), and then dried by heat (70°C) overnight. The *Xenorhabdus* cells were killed in such a  
312 way, while the heat stable [43] antimicrobial compounds remained active.

313  
314

## 2.7. Statistical Analysis


315 ANOVA procedures were used following the procedures of the SAS 9.4 Software, mostly  
316 due to the unbalanced data set. The significant differences ( $\alpha = 0.05$ ) between treatment  
317 means were assessed using the Least Significant Difference (LSD).

## 318 **3. RESULTS**

319  
320  
321

### 3.1. Results of Experiments, Aimed at Helping to Choose the Best *Xenorhabdus* Strains for This Study

322 Results shown in Fig. 2, and a qualitative evaluation of the inactivation zones, indicated the  
323 appropriate bacteria to use. As expected, *X. budapestensis* (EMA) and *X. szentirmaii* were  
324 the best. Results of the overlay bioassay experiment with different *Xenorhabdus* strains on  
325 *K. pneumoniae* helped to make the right decision when choosing antimicrobial producing  
326 strains.



A: *X. nematophila* DSM 3370

B: *X. cabanillasii* BP

C: *X. nematophila* ATTC19061

D: *X. bovienii* NYH

E: *X. budapestensis* DSM16342<sup>T</sup>

F: *X. szentirmaii* DSM16338<sup>T</sup>

327

328 Figure 2 Comparison of the antimicrobial potential of different *Xenorhabdus* strains  
329 (representing species) in overlay bioassays [43]. (Photo: Andrea Máthé Fodor. The Ohio  
330 State University, Wooster, OH, USA)

331 **3.2. Endurance of the antimicrobial peptides of *X. budapestensis* to pepsin, - and**  
332 **trypsin digestion**

333

334 As demonstrated by Fig. 3, the overnight pepsin-digested EMA CFCM remained active  
335 against *Pseudomonas aeruginosa*. The trypsin-digested samples also preserved their anti-  
336 Gram-positive (on *S. aureus*) and anti-Gram-negative (*Es. coli*) activities, (not shown).





Fig. 3. Experimental evidence that the antimicrobial compounds of *X. budapestensis* cell-free media are resistant to the proteolytic activity of pepsin. After 24 h incubation at 37 °C a large inactivation zone could be seen, demonstrating a significant antimicrobial activity of the pepsin-treated EMA CFCM.

337 **3.3. Efficacy of EMA CFCM on *C. perfringens* Laboratory Strain NCAIM 1471**

338 The cell-free EMA CFCM exerted strong antimicrobial activity on *C. perfringens* laboratory  
339 strain NCAIM 1471 in an agar diffusion test. The large inactivation zone of 3.7 cm diameter  
340 shows the anti - *Clostridium* activity (Fig 3). The question arises as to whether the  
341 pathogenic poultry isolates were also sensitive.

342

343



344  
345  
346

347 Figure 4. Anti- *Clostridium* activity of cell-free culture medium of *Xenorhabdus budapestensis*  
348 on *Clostridium perfringens* NCAIM 1417 strain in agar diffusion test [46] [47]. (Photo: Dr.  
349 Csaba Pintér, University of Pannonia, Keszthely, Hungary)

350

351 **3.4 Results of the Comparison of the Sensitivities (MID values) of 13 *Clostridium***  
352 ***perfringens* strains isolated from Poultry to Cell-Free Culture Media (CFCM) of**  
353 ***Xenorhabdus budapestensis* (EMA) in Liquid Cultures**

354 Table 1 lists the MID values as a qualitative parameter of the sensitivity of each of the  
355 poultry isolates to the antibacterial compounds of *X. budapestensis*. A majority of the  
356 examined strains are sensitive but one of the 13 was resistant (LM24). No direct interrelation  
357 between the degree of EMA sensitivity and other behavior could be demonstrated. The  
358 results provide a good message: The majority of *C. perfringens* isolates are sensitive.  
359 However, they also provide a bad message: There are EMA-resistant resistant *C.*  
360 *perfringens* isolates, even if they are rare.

361 None of the samples taken from cultures with no visible proliferation contained any CFU,  
362 indicating that the toxicity was complete. Whether the differences in the sensitivities could  
363 relate to the cellular phenotype was not revealed by this experiment, although the *C.*  
364 *perfringens* isolates were rather different concerning colony morphology and hemolytic  
365 behavior (Fig 5).  
366



367  
368

369 Figure 5 *Clostridium perfringens* isolates LM1, LM2 and LM24 differing in colony  
370 morphology, sporulation, and hemolytic behavior. (Photo: Dr. László Makrai, (Department of  
371 Microbiology and Infectious Diseases, University of Veterinary Science, Hungary).

372

373 Table 1 MID values of *Clostridium perfringens* isolates from chicken differing in colony  
 374 morphology and hemolytic behavior

<i>C. perfringens</i> isolates from poultry (L. Makrai, unpublished)	Minimum Inhibiting Dilutions (MID) Values (V/V%) of the cell-free culture medium (CFCM) of <i>Xenorhabdus budapestensis</i> (EMA) Inhibiting Bacterial Proliferation	Conclusion
LM 1	< 10	Extremely sensitive
LM 2	< 30	Sensitive
LM 3	< 10	Extremely sensitive
LM 4	< 10	Extremely sensitive
LM 5	< 10	Extremely sensitive
LM 8	< 30	Sensitive
LM 11	< 10	Extremely sensitive
LM 14	< 10	Extremely sensitive
LM 15	< 10	Extremely sensitive
LM 16	< 10	Extremely sensitive
LM19	< 10	Extremely sensitive
LM20	< 30	Sensitive
LM 24	> 50	Resistant

375  
 376  
 377

#### 4. Discussion

378 The *in vitro* experiments demonstrated that antimicrobial peptides of *X. budapestensis*  
 379 (EMA) were highly toxic for all but one (LM 24) *C. perfringens* isolates. Dr. Klaus Teichmann  
 380 (Biomim, Tulln, Austria), as a courtesy, tested EMA and EMC CFCM preparations, obtained  
 381 from us. He declared that the CFCM of EMA exerted an extremely strong anticoccidial  
 382 activity on both *Clostridium* and *Eimeria* cells. He declared that he had not ever worked with  
 383 such an efficient anticoccidial preparation before as EMA CFCM. Dr. Teichmann found a  
 384 lower concentration range within which *E. tenella* cells died, while the cells of the chicken  
 385 tissue culture were not affected, (Klaus Teichmann, personal communication). These facts  
 386 are arguments for taking the potential use of EMA and EMC antimicrobial peptides, as  
 387 potential anticoccidial agents administered *per os*, into consideration.

388 But there are arguments against using XENOFOD as well, and they are those data which  
 389 showed *in vitro* cytotoxicity on the permanent chicken liver cell line LMH [60]. Dr. Ganas and  
 390 her associates (Aziza Amin, Irina Profjeva, and Micheal Hess) tested the cytopathogenic  
 391 effect of different dilutions of the same samples of sterile cell-free media (CFCM) of EMA  
 392 and EMC on permanent chicken liver LMH cells, as Dr. Teichmann. They demonstrated that  
 393 EMA CFCM at a dose of < 5% V/V concentration was harmless, but at >5%V/V  
 394 concentrations they seriously damaged the cell layer. Doses >10% V/V caused total  
 395 destruction of the cell layer, while that of 5 – 10% V/V resulted in about a 50% damage  
 396 within the first 24h, and this damage was not repaired in the next 72 hrs. As for EMC, only  
 397 the dose of 32% resulted in complete cell layer destruction, but the lower doses of 1-20%  
 398 V/V also resulted in ~ 50% permanent damage, calculated on the base of the score scale of  
 399 Amin et al. [60] (2012); (Petra Ganes et al., personal communication).

400 Fabclavines are the predominant antimicrobial compound produced by both EMA and EMC  
401 and were isolated and purified [51], and was not suggested as a future drug because of its  
402 extremely large target size and toxicity to eukaryotic targets. This kind of “certification” is  
403 usually quite enough to place a candidate drug molecule into the wastebasket, despite its  
404 super strong antimicrobial effects. However, an exception with fabclavine may be considered  
405 because of the following arguments:

406 First, there are not only prokaryotic, but eukaryotic pathogens also exist. Coccidiosis is the  
407 best example where a prokaryotic *C. perfringens* and a eukaryotic *E. tenella* cooperate in  
408 causing the disease, and both should be controlled.

409 Second, there is practically no vaccination technique against *C. perfringens* [28]. So the  
410 introduction of new antimicrobial compounds should be taken into consideration.

411 We are not the only team walking on this road. Recently, there have been several research  
412 directions attempting to solve the coccidiosis problem. A project includes a search for novel  
413 antibiotic-delivery systems, such as using ovotransferrin as a targeting molecule [61].  
414 Another approach is to improve the usefulness of commonly used anticoccidials and  
415 antibiotics, which have recently been tested on a subclinical necrotic enteritis model [62].  
416 Recently proline-rich antimicrobial peptides are considered as potential therapeutics against  
417 antibiotic-resistant bacteria [63]. The designer proline-rich antibacterial peptide A3-APO  
418 prevents the Gram-positive *Bacillus anthracis* mortality by deactivating bacterial toxins [64].  
419 Even more recently two (NZ2114 and MP1102) novel plectasin-derived peptides have been  
420 designed for targeting Gram-positive bacteria, and the tests on gas gangrene-associated *C.*  
421 *perfringens* provided encouraging results [65].

422 The hopes of applying probiotics have been also emerging [66] [67] [68]. The use of  
423 vegetative *Bacillus amyloliquefaciens* cells did not justify the hopes: they did not confer  
424 protection against necrotic enteritis in broilers, despite the high antibacterial activity of its  
425 supernatant against *C. perfringens* in vitro [69].

## 426 **5. Conclusions**

427 There are two alternative approaches to control coccidiosis in broiler chicken: the  
428 vaccination and the “chemotherapy”, (that is, a search for gastro-intestinally active,  
429 autoclaveable antimicrobial peptides active against both *C. perfringens* and *E. tenella*).

430 Considering that there are publications about antibiotic resistant and multiresistant pathogen  
431 *C. perfringens* [70] [71], and that the coccidiosis problem has not yet seem to be solved by  
432 using vaccination, the search for new efficient antimicrobials to control coccidiosis have  
433 probably been justified.

434 On the basis of *in vitro* studies, fabclavine alone (and / or as a component of interacting  
435 antimicrobial active peptide complexes present in the CFCM of EMA and EMC) fulfil the  
436 criteria of a promising chemotherapeutic agent *in vitro*, that is, acting as strong antibacterial  
437 on *C. perfringens* and as strong apoptotic cytotoxic compounds on the unicellular eukaryotic  
438 pathogen, *E. tenella*.

439



440 However, the cytotoxicity may pose a serious problem of practical use. Indeed, we found  
441 that the CFCM of both EMA and EMC were cytotoxic *in vitro* in permanent chicken liver cells.  
442 But the *in vitro* and the *in vivo* situation are completely different.

443

444 If it happened that the orally administered fabclavine (and/or the whole AMP complex), due  
445 to their proteolytic endurance), might act *in vivo* as strong anti-Clostridia and anti-Eimeria  
446 agents in the GI, without causing any harm of the organism to be protected, it would have a  
447 chance to be register and use Xenofood as an anticoccidial bio-preparation. This option  
448 cannot be ruled out if the adsorption from the gut, were similarly low as that of the orally  
449 administred vancomycin [72].

450

451 We believe that an *in vivo* XENOFood feeding experiment would be necessary to learn  
452 whether the orally administrated antimicrobial peptides produced by *X. budapestensis* (EMA)  
453 and *X. szentirmaii* (EMC), *in vitro* against both the prokaryotic (*C. perfringens*) and the  
454 eukaryotic (*E. tenella*) pathogens causing coccidiosis in chicken, could be used in broiler  
455 cockerels.

456 We are ready for *in vivo* bioassay and looking for cooperative partners.

457

## 458 **ACKNOWLEDGEMENTS**

459

460 A The authors express their sincere thanks and appreciations to our coauthor Professor  
461 **Károly Dublecz**, the previous Dean of the Georgikon Faculty, for generously providing full  
462 (moral and even manual) help needed for preparing and accomplishing the *in vivo*  
463 experiments and following *Clostridium* bioassays in his Unit.

464 We thank **Dr. Petra Ganas** for unselfishly accomplishing those experiments which clarified  
465 the cytotoxicity of the CFCM by EMA and EMC, and kindly provided all data to us with the  
466 generous permission of his boss, Professor Michael Hess (University of Veterinary, Vienna,  
467 Austria). We thank **Dr. Klaus Teichmann** (Biomim, Tulln, Austria) who tested the same  
468 samples (that Petra Ganas worked with) on *Eimeria* cells and shared his conclusions with us  
469 unselfishly.

470 We would like to express thanks and appreciations the professional technical help to the lab  
471 technicians in the Vet School for their invaluable technical help, especially **Miss Éva**  
472 **Kolozsvári** and **Miss Teréz Halasi**.

473 We are extremely grateful to **Mrs. Gwen Short** (Head of the Library in Ohio State University,  
474 Wooster, OH) for providing shelter and professional support to accomplish the MS properly.  
475 Many thanks for that.

476 This MS would have never been materialized in the absence of the strong and unselfish help  
477 and support of **Mrs. Andrea Máthé-Fodor**, (Molecular and Cellular Imaging Center, Ohio  
478 State University, Wooster, OH, USA) throughout in every stage of the work.

479 The *in vitro* *Clostridium* bioassays preceding Xenofood experiments were supported by  
480 **TÁMOP-4.2.2/B-10/1-2010-0025** project based on the financial support the State of Hungary  
481 and the EU.

482

483 **COMPETING INTERESTS**

484

485 The authors declare that the research was conducted in the absence of any commercial or  
486 financial relationships that could be construed as a potential conflict of interest.

487 Authors have declared that no competing interests exist.

488

489 **AUTHORS' CONTRIBUTIONS**

490

491 **Dr. Habil. András Fodor** was the project initiator and put the MS together. Furthermore, he  
492 provided the bacterium cultures and inoculated and fermented Xenofood with *X.*  
493 *budapestensis* and *X. szentirmaii*.

494 **Professors László Makrai** and **László Fodor** (University of Veterinary Sciences, Budapest,  
495 Hungary) were the intellectual leader and governing the *Clostridium* tests. We worked on the  
496 *Clostridium perfringens* isolates (LM1 - LM24), previously collected, identified and deposited  
497 by them in the Department's stock collection. All but a few experiments were carrying on in  
498 their Laboratory at Department of Microbiology and Infectious Diseases, Faculty of  
499 Veterinary Science, Szent István University, Budapest, Hungary.

500 **Associate Professor Dr. István Venekei** carried out the trypsin digestions and bio-assayed  
501 the antimicrobial activities of the digested preparations with A. Fodor at the Department of  
502 Biochemistry at Eötvös University in Budapest, Hungary.

503 **Professor Emeritus Ferenc Husvéth**, (University of Pannonia, Keszthely, Hungary  
504 biochemist, independently from the others, guided our experiments on testing the pepsin  
505 sensitivity of the antimicrobial active compounds EMA and EMC CFCM in Keszthely, many  
506 thanks for it. This work was essential and Dr. Husvéth should definitely be our coauthor.

507 **Associate Professor Dr. László Pál** (University of Pannonia, Keszthely, Hungary) prepared  
508 the media by mixing up the food components before A. Fodor inoculated it with *X.*  
509 *budapestensis* and *X. szentirmaii*. He weighted the animals daily and calculated the food  
510 conversion. His work was essential and Dr. Pál should definitely be one of the coauthors.

511 **Research Associate Dr. Andor Molnár** (University of Pannonia, Keszthely, Hungary) and  
512 A. Fodor has applied previously published overlay,- and agar diffusion technique to *C.*  
513 *perfringens* and he was the pioneer of the *Clostridium* experiments at the Department of  
514 Animal Sciences and Animal Husbandry, Georgikon Faculty, University of Pannonia,  
515 (Keszthely, H-8360 Hungary) working with chicken ileal digestions.

516 The statistical analyses were carried out or guided by **Adjunct Emeritus Professor Dr.**  
517 **Sándor Józsa** (University of Pannonia, Keszthely, Hungary).

518 **Adjunct Emeritus Professor Dr. Csaba Pintér** (University of Pannonia, Keszthely,  
519 Hungary) made all but one (made by Andrea **Máthé – Fodor**, Ohio State University,  
520 Wooster, OH, USA) photos published here

521 **Adjunct Professor Michael G. Klein** (The Ohio State University) made the final shaping  
522 and proofreading of the MS and was the spiritual rector of putting the data and ideas in the  
523 present form, which has been an invaluable contribution.

524 **REFERENCES**

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

1. Nawrocki KL, Crispell EK, McBride SM. Antimicrobial peptide resistance mechanisms of Gram-positive bacteria. *Antibiotics (Basel)*. 2014 Oct 13;3(4):461-92. doi: 10.3390/antibiotics3040461. 10.3390/antibiotics3040461. PMID: 25419466 Free PMC Article
2. Gruenheid S, Le Moual H. Resistance to antimicrobial peptides in gram-negative bacteria. *FEMS Microbiol Lett*. 2012 May;330(2):81-9. doi: 10.1111/j.1574-6968.2012.02528.x. Epub 2012 Mar 12. Review. PMID: 22339775
3. Borst LB, Suyemoto MM, Robbins KM, Lyman RL, Martin MP, Barnes HJ. Molecular epidemiology of *Enterococcus cecorum* isolates recovered from enterococcal spondylitis outbreaks in the southeastern United States. *Avian Pathol*. 2012 Oct;41(5):479-85. doi: 10.1080/03079457.2012.718070. Epub 2012 Sep 17.
4. Miller WR, Munita JM, Arias CA. Mechanisms of antibiotic resistance in enterococci. *Expert Rev Anti Infect Ther*. 2014 Oct;12(10):1221-36. doi: 10.1586/14787210.2014.956092. Review. PMID: 25199988 Free PMC Article
5. Palmer, K.L., van Schaik, W., Willems, R.J.L., and Gilmore, M.S. (2014) Enterococcal Genomics. In: Gilmore, M.S., Clewell, D.B., Ike, Y., Shankar, N., (Editors). *Enterococci: From Commensals to Leading Causes of Drug Resistant Infection* [Internet]. Boston, MA, USA: Massachusetts Eye and Ear Infirmary; PMID: 24649511 Free Books & Documents
6. Dahms C, Hübner NO, Wilke F, Kramer A. Mini-review: Epidemiology and zoonotic potential of multiresistant bacteria and *Clostridium difficile* in livestock and food. *GMS Hyg Infect Control*. 2014 Sep 30;9(3):Doc21. doi: 10.3205/dgkh000241. eCollection 2014.
7. Lu Y, Zhao H, Sun J, Liu Y1, Zhou X, Beier RC, Wu G, Hou X. Characterization of multidrug-resistant *Salmonella enterica* serovars Indiana and Enteritidis from chickens in Eastern China. *PLoS One*. 2014 May 2;9(5):e96050. doi: 10.1371/journal.pone.0096050. eCollection 2014.
8. Makrai L, Nemes C, Simon A, Ivanics E, Dudás Z, Fodor L, Glávits R. Association of *Enterococcus cecorum* with vertebral osteomyelitis and spondylolisthesis in broiler parent chicks. *Acta Vet Hung*. 2011 Mar;59(1):11-21. doi: 10.1556/AVet.59.2011.1.2. Crossref, PubMed, Google Scholar
9. Dolka B, Chrobak-Chmiel D, Makrai L, Szeleszczuk P. Phenotypic and genotypic characterization of *Enterococcus cecorum* strains associated with infections in poultry. *BMC Vet Res*. 2016 Jun 27;12(1):129. doi: 10.1186/s12917-016-0761-1. PMID: 27350248.
10. Lebreton F, van Schaik W, McGuire AM, Godfrey P, Griggs A, Mazumdar V, Corander J, Cheng L, Saif S, Young S, Zeng Q, Wortman J, Birren B, Willems RJ, Earl AM, Gilmore MS. Emergence of epidemic multidrug-resistant *Enterococcus faecium* from animal and commensal strains. *MBio*. 2013 Aug 20;4(4). pii: e00534-13. doi: 10.1128/mBio.00534-13.
11. Szmolka A, Nagy B. Multidrug resistant commensal *Escherichia coli* in animals and its impact for public health. *Front Microbiol*. 2013 Sep 3;4:258. doi: 10.3389/fmicb.2013.00258.

- 569 12. Hasman H, Hammerum AM, Hansen F, Hendriksen RS, Olesen B, Agersø Y, Zankari E,  
570 Leekitcharoenphon P, Stegger M, Kaas RS, Cavaco LM, Hansen DS, Aarestrup FM, Skov  
571 RL. Detection of mcr-1 encoding plasmid-mediated colistin-resistant *Escherichia coli*  
572 isolates from human bloodstream infection and imported chicken meat, Denmark 2015.  
573 *Euro Surveill.* 2015;20(49). doi: 10.2807/1560-7917.ES.2015.20.49.30085. Erratum in:  
574 *Euro Surveill.* 2015; 20(50). doi: 10.2807/1560-7917.ES.2015.20.50.30091.
- 575 13. Van Immerseel F1, De Buck J, Pasmans F, Huyghebaert G, Haesebrouck F, Ducatelle R.  
576 *Clostridium perfringens* in poultry: an emerging threat for animal and public health.  
577 *Avian Pathol.* 2004 Dec;33(6):537-49. PMID: 15763720 DOI:  
578 10.1080/03079450400013162 [Indexed for MEDLINE]
- 579 14. Timbermont L1, Lanckriet A, Gholamiandehkordi AR, Pasmans F, Martel A,  
580 Haesebrouck F, Ducatelle R, Van Immerseel F. Origin of *Clostridium perfringens* isolates  
581 determines the ability to induce necrotic enteritis in broilers. *Comp Immunol Microbiol*  
582 *Infect Dis.* 2009 Nov;32(6):503-12. doi: 10.1016/j.cimid.2008.07.001. Epub 2008 Sep 9.  
583 Epub 2008 Sep 9. PMID: 18783830 [Indexed for MEDLINE]
- 584 15. Van Immerseel F, Rood JI, Moore RJ, Titball RW. Rethinking our understanding of the  
585 pathogenesis of necrotic enteritis in chickens. *Trends Microbiol.* 2009 Jan;17(1):32-6.  
586 doi: 10.1016/j.tim.2008.09.005. Epub 2008 Oct 30. PMID: 18977143 DOI:  
587 10.1016/j.tim.2008.09.005 [Indexed for MEDLINE]
- 588 16. Stanley D, Wu SB, Rodgers N, Swick RA, Moore RJ. Differential responses of cecal  
589 microbiota to fishmeal, *Eimeria* and *Clostridium perfringens* in a necrotic enteritis  
590 challenge model in chickens. *PLoS One.* 2014 Aug 28;9(8):e104739. doi:  
591 10.1371/journal.pone.0104739. eCollection 2014. PMID: 25167074 Free PMC Article
- 592 17. Kitessa SM, Nattrass GS, Forder RE, McGrice HA, Wu SB, Hughes RJ. Mucin gene mRNA  
593 levels in broilers challenged with *eimeria* and/or *Clostridium perfringens*. *Avian Dis.*  
594 2014 Sep;58(3):408-14. PMID: 25518436 DOI: 10.1637/10757-122313-Reg.1 [Indexed  
595 for MEDLINE].
- 596 18. Dahiya, J.P., Hoehler, D., Van Kessel, A.G. and Drew, M.D. (2007) Dietary encapsulated  
597 glycine influences *Clostridium perfringens* and *Lactobacilli* growth in the gastro-  
598 intestinal tract of broiler chickens. *J Nutr.* 2007 Jun;137(6):1408-14. PMID: 17513399  
599 Free Article
- 600 19. Teirlynck E, Bjerrum L, Eeckhaut V, Huyghebaert G, Pasmans F, Haesebrouck F, Dewulf J,  
601 Ducatelle R, Van Immerseel F. The cereal type in feed influences gut wall morphology  
602 and intestinal immune cell infiltration in broiler chickens. *Br J Nutr.* 2009 Nov;  
603 102(10):1453-61. doi: 10.1017/S0007114509990407. Epub 2009 Aug 7. PMID:  
604 19664304 [Indexed for MEDLINE]
- 605 20. Molnár A1, Hess C, Pál L, Wágner L, Awad WA, Husvéth F, Hess M, Dublec K.  
606 Composition of diet modifies colonization dynamics of *Campylobacter jejuni* in broiler  
607 chickens. *J Appl Microbiol.* 2015 Jan;118(1):245-54. doi: 10.1111/jam.12679. Epub  
608 2014 Nov 25. PMID: 25358748 [Indexed for MEDLINE]
- 609 21. Keyburn AL, Bannam TL, Moore RJ, Rood JI. NetB, a pore-forming toxin from necrotic  
610 enteritis strains of *Clostridium perfringens*. *Toxins (Basel).* 2010 Jul;2(7):1913-27. doi:  
611 10.3390/toxins2071913. Epub 2010 Jul 23. Review. PMID: 22069665 Free PMC Article
- 612 22. Verherstraeten S, Goossens E, Valgaeren B, Pardon B, Timbermont L, Haesebrouck F,  
613 Ducatelle R, Deprez P, Wade KR, Tweten R, Van Immerseel F. *Perfringolysin O*: The

- 614 Underrated Clostridium perfringens Toxin? Toxins (Basel). 2015 May 14;7(5):1702-21.  
615 doi: 10.3390/toxins7051702. Review. PMID: 26008232 Free PMC Article
- 616 23. Timbermont L, De Smet L, Van Nieuwerburgh F, Parreira VR, Van Driessche G,  
617 Haesebrouck F, Ducatelle R, Prescott J, Deforce D, Devreese B, Van Immerseel F. Perfrin,  
618 a novel bacteriocin associated with netB positive *Clostridium perfringens* strains from  
619 broilers with necrotic enteritis. Vet Res. 2014 Apr 5;45:40. doi: 10.1186/1297-9716-  
620 45-40. PMID: 24708344 PMCID: PMC3992141 DOI: 10.1186/1297-9716-45-40  
621 [Indexed for MEDLINE] Free PMC Article
- 622 24. Romero-Saavedra F, Laverde D, Wobser D, Michaux C, Budin-Verneuil A, Bernay B,  
623 Benachour A, Hartke A, Huebner J. Identification of peptidoglycan-associated proteins  
624 as vaccine candidates for enterococcal infections. PLoS One. 2014 Nov 4;  
625 9(11):e111880. doi: 10.1371/journal.pone.0111880. eCollection 2014. PMID:25369230  
626 Free PMC Article [Indexed for MEDLINE] Free PMC Article
- 627 25. Dohms JE, Allen PH, Cloud SS. The immunization of broiler chickens against type C  
628 botulism. Avian Dis. 1982 Apr-Jun;26(2):340-5. PMID:7049149 [Indexed for MEDLINE]
- 629 26. Fernandes da Costa SP, Mot D, Bokori-Brown M, Savva CG, Basak AK, Van Immerseel F,  
630 Titball RW. Protection against avian necrotic enteritis after immunisation with NetB  
631 genetic or formaldehyde toxoids. Vaccine. 2013 Aug 20;31(37):4003-8. doi:  
632 10.1016/j.vaccine.2013.05.063. Epub 2013 May 29 PMID:23727000 Free PMC Article
- 633 27. Mot, D., Timbermont, L., Delezie, E., Haesebrouck, F., Ducatelle, R., and Van Immerseel,  
634 F. Day-of-hatch vaccination is not protective against necrotic enteritis in broiler  
635 chickens. Avian Pathol. 2013 Apr;42(2):179-84. doi: 10.1080/03079457.2013.778955.  
636 PMID: 23581446 [Indexed for MEDLINE]
- 637 28. Mot D, Timbermont L, Haesebrouck F, Ducatelle R, Van Immerseel F. Progress and  
638 problems in vaccination against necrotic enteritis in broiler chickens. Avian Pathol.  
639 2014; 43(4):290-300. doi: 10.1080/03079457.2014.939942.PMID: 24980518 DOI:  
640 10.1080/03079457.2014.939942 [Indexed for MEDLINE]
- 641 29. Jang SI, Lillehoj HS, Lee SH, Lee KW, Lillehoj EP, Hong YH, An DJ, Jeong W, Chun JE,  
642 Bertrand F, Dupuis L, Deville S, Arous JB. Vaccination with Clostridium perfringens  
643 recombinant proteins in combination with Montanide™ ISA 71 VG adjuvant increases  
644 protection against experimental necrotic enteritis in commercial broiler chickens.  
645 Vaccine. 2012 Aug 3;30(36):5401-6. doi: 10.1016/j.vaccine.2012.06.007. Epub 2012  
646 Jun 17. PMID: 22713719.
- 647 30. Bangoura B, Alnassan AA, Lendner M, Shehata AA, Krüger M, Dausgschies A. Efficacy of  
648 an anticoccidial live vaccine in prevention of necrotic enteritis in chickens. Exp  
649 Parasitol. 2014 Oct; 145:125-34. doi: 10.1016/j.exppara.2014.08.004. Epub 2014 Aug  
650 14. PMID: 25131774. [Indexed for MEDLINE]
- 651 31. Mojsoska B, Jenssen H. Peptides and Peptidomimetics for Antimicrobial Drug Design.  
652 Pharmaceuticals (Basel). 2015 Jul 13;8(3):366-415. doi: 10.3390/ph8030366. Review.  
653 PMID: 26184232 Free PMC Article
- 654 32. Ötvös L Jr, Wade JD. Current challenges in peptide-based drug discovery. A Specialty  
655 Grand Challenge Article. Front Chem. 2014 Aug 8;2:62. doi: 10.3389/fchem.2014.00062.  
656 eCollection 2014. PMID: 25152873 PMCID: PMC4126357. Free PMC Article
- 657 33. Lin L, Nonejuie P, Munguia J, Hollands A, Olson J, Dam Q, Kumaraswamy M, Rivera H Jr,  
658 Corriden R, Rohde M, Hensler ME1, Burkart MD, Pogliano J, Sakoulas G, Nizet V.  
659 Azithromycin Synergizes with Cationic Antimicrobial Peptides to Exert Bactericidal and



- 660 Therapeutic Activity Against Highly Multidrug-Resistant Gram-Negative Bacterial  
661 Pathogens *EBioMedicine*. 2015 Jun 10; 2(7):690-8. doi: 10.1016/j.ebiom.2015.05.021.  
662 eCollection 2015 Jul. PMID: 26288841 PMCID: PMC4534682 DOI:  
663 10.1016/j.ebiom.2015.05.021 [Indexed for MEDLINE] Free PMC Article. Comment in  
664 Rethinking the Antibiotic Discovery Paradigm. [*EBioMedicine*. 2015]
- 665 34. Forst S, Nealson K. Molecular biology of the symbiotic-pathogenic bacteria  
666 *Xenorhabdus* spp. and *Photorhabdus* spp. *Microbiol Rev*. 1996 Mar;60(1):21-43.  
667 Review. PMID: 8852894 PMCID: PMC239416 [Indexed for MEDLINE] Free PMC Article
- 668 35. Nollmann FI, Dauth C, Mulley G, Kegler C, Kaiser M, Waterfield NR, Bode HB. Insect-  
669 specific production of new GameXPeptides in *Photorhabdus luminescens* TTO1,  
670 widespread natural products in entomopathogenic bacteria. *Chembiochem*. 2015 Jan  
671 19;16(2):205-8. doi: 10.1002/cbic.201402603. Epub 2014 Nov 25. PMID: 25425189  
672 DOI: 10.1002/cbic.201402603 [Indexed for MEDLINE]
- 673 36. Vivas EI, Goodrich-Blair H. *Xenorhabdus nematophilus* as a model for host-bacterium  
674 interactions: rpoS is necessary for mutualism with nematodes. *J Bacteriol*. 2001  
675 Aug;183(16):4687-93. PMID: 11466270 PMCID: PMC99521 DOI:  
676 10.1128/JB.183.16.4687-4693.2001 [Indexed for MEDLINE] Free PMC Article
- 677 37. Park D, Ciezki K, van der Hoeven R, Singh S, Reimer D, Bode HB, Forst S. Genetic  
678 analysis of xenocoumacin antibiotic production in the mutualistic bacterium  
679 *Xenorhabdus nematophila*. *Mol Microbiol*. 2009 Sep;73(5):938-49. doi: 10.1111/j.1365-  
680 2958.2009.06817.x. Epub 2009 Aug 4. PMID: 19682255 [Indexed for MEDLINE]
- 681 38. Gualtieri M, Aumelas A, Thaler JO. Identification of a new antimicrobial lysine-rich  
682 cyclolipopeptide family from *Xenorhabdus nematophila*. *J Antibiot (Tokyo)*. 2009 Jun;  
683 62(6):295-302. doi: 10.1038/ja.2009.31. Epub 2009 Apr 17. PMID: 19373275 DOI:  
684 10.1038/ja.2009.31 [Indexed for MEDLINE]
- 685 39. Bode HB, Brachmann AO, Jadhav KB, Seyfarth L, Dauth C, Fuchs SW, Kaiser M,  
686 Waterfield NR, Sack H, Heinemann SH, Arndt HD. Structure Elucidation and Activity of  
687 Kolossin A, the D-/L-Pentadecapeptide Product of a Giant Nonribosomal Peptide  
688 Synthetase. *Angew Chem Int Ed Engl*. 2015 Aug 24; 54(35):10352-5. doi:  
689 10.1002/anie.201502835. Epub 2015 Jun 26. PMID: 26118790 DOI:  
690 10.1002/anie.201502835 [Indexed for MEDLINE]
- 691 40. Reimer D, Bode HB. A natural prodrug activation mechanism in the biosynthesis of non-  
692 ribosomal peptides. *Nat. Prod. Rep*. 2013, 31(2): 154-159. doi: 0.1039/c3np70081j.  
693 Review. PMID: 24356302
- 694 41. Medema MH, Kottmann R, Yilmaz P, 1, Cummings M, Biggins JB, Blin K, Claesen J, et al.,,  
695 Minimum Information about a Biosynthetic Gene cluster. *Nat Chem Biol*. 2015 Sep;  
696 11(9):625-31. doi: 10.1038/nchembio.1890. doi:10.1038/nchembio.1890. PMID:  
697 26284661 PMCID: PMC5714517 DOI: 10.1038/nchembio.1890 [Indexed for MEDLINE]  
698 Free PMC Article
- 699 42. Houard J, Aumelas A, Noël T, Pages S, Givaudan A, Fitton-Ouhabi V, Villain-Guillot P,  
700 Gualtieri M. Cabanillasin, a new antifungal metabolite, produced by entomopathogenic  
701 *Xenorhabdus cabanillasii* JM26. *J Antibiot (Tokyo)*. 2013 Oct;66(10):617-20. doi:  
702 10.1038/ja.2013.58. Epub 2013 Jun 12. PMID: 23756685 DOI: 10.1038/ja.2013.58  
703 [Indexed for MEDLINE]
- 704 43. Furgani G, Böszörményi E, Fodor A, Máthé-Fodor A, Forst S, Hogan JS, Katona Z, Klein  
705 MG, Stackebrandt E, Szentirmai A, Sztaricskai F, Wolf SL. *Xenorhabdus* antibiotics: a

- 706 comparative analysis and potential utility for controlling mastitis caused by bacteria. *J*  
707 *Appl Microbiol.* 2008 Mar; 104(3):745-58. Epub 2007 Nov 1. PMID: 17976177 DOI:  
708 10.1111/j.1365-2672.2007.03613.x [Indexed for MEDLINE]
- 709 44. Lengyel K1, Lang E, Fodor A, Szállás E, Schumann P, Stackebrandt E. Description of four  
710 novel species of *Xenorhabdus*, family Enterobacteriaceae: *Xenorhabdus budapestensis*  
711 sp. nov., *Xenorhabdus ehlersii* sp. nov., *Xenorhabdus innexi* sp. nov., and *Xenorhabdus*  
712 *szentirmaii* sp. nov. *Syst Appl Microbiol.* 2005 Mar; 28(2):115-22. Erratum in: *Syst Appl*  
713 *Microbiol.* 30 (1): 83. Also in March/April 2014 Volume 2 Issue 2 e00190-14 Genome  
714 Announcements genomea.asm.org ehlersii sp. nov., *Xenorhabdus innexi* sp. nov., and  
715 *Xenorhabdus szentirmaii* sp. nov. *Syst. Appl. Microbiol.* **28**: 115. [http://dx.doi.org/10.1](http://dx.doi.org/10.1016/j.syapm.2004.10.004.227)  
716 [016/j.syapm.2004.10.004.227](http://dx.doi.org/10.1016/j.syapm.2004.10.004.227) PMID: 15830803 DOI: 10.1016/j.syapm.2004.10.004.  
717 [Indexed for MEDLINE]
- 718 45. Böszörményi E, Érsek T, Fodor A, Fodor AM, Földes LS, Hevesi M, Hogan JS, Katona Z,  
719 Klein MG, Kormány A, Pekár S, Szentirmai A, Sztaricskai F, Taylor RA. Isolation and  
720 activity of *Xenorhabdus* antimicrobial compounds against the plant pathogens *Erwinia*  
721 *amylovora* and *Phytophthora nicotianae*. *J Appl Microbiol.* 2009 Sep;107(3):746-59.  
722 doi: 10.1111/j.1365-2672.2009.04249.x. Epub 2009 Mar 23. PMID: 19320949 DOI:  
723 10.1111/j.1365-2672.2009.04249.x [Indexed for MEDLINE]
- 724 46. Vozik D, Bélafi-Bakó K, Hevesi M, Böszörményi E, Fodor A. Effectiveness of a peptide-  
725 rich fraction from *Xenorhabdus budapestensis* culture against fire blight disease on  
726 apple blossoms. *Not. Bot. Horti. Agrobo.* 2015 43(2): 547-553. DOI:  
727 10.15835/nbha4329997, Available online: [www.notulaebotanicae.ro](http://www.notulaebotanicae.ro)
- 728 47. Vozik D. [Agricultural Utilization of Antimicrobial Compounds produced by  
729 Entomopathogenic Bacteria, PhD Thesis in Hungarian] Entomopatogén Baktériumok  
730 Antimikrobiális Hatású Anyagcseretermékeinek Felhasználhatósága a  
731 Növényvédelemben Ph. D. Thesis, University of Pannonia, Veszprém, Hungary, 2017.
- 732 48. Gualtieri M, Ogier JC, Pagès S, Givaudan A, Gaudriault S. Draft Genome Sequence and  
733 Annotation of the Entomopathogenic Bacterium *Xenorhabdus szentirmaii* Strain  
734 DSM16338. *Genome Announc.* 2014 Mar 13; 2(2). pii: e00190-14. doi:  
735 10.1128/genomeA.00190-14. PMID: 24625876 PMCID: PMC3953197 DOI:  
736 10.1128/genomeA.00190-14 Free PMC Article
- 737 49. Nollmann FI, Dowling A, Kaiser M, Deckmann K, Grösch S, Ffrench-Constant R, Bode HB.  
738 Synthesis of szentiamide, a depsipeptide from entomopathogenic *Xenorhabdus*  
739 *szentirmaii* with activity against *Plasmodium falciparum*. *Beilstein J Org Chem.*  
740 2012;8:528-33. doi: 10.3762/bjoc.8.60. Epub 2012 Apr 11. PMID: 22563351 PMCID:  
741 PMC3343279 Free PMC Article
- 742 50. Fuchs SW, Sachs CC, Kegler C, Nollmann FI, Karas M, Bode HB. Neutral loss  
743 fragmentation pattern based screening for arginine-rich natural products in and  
744 *Photorhabdus*. *Anal Chem.* 2012 Aug 21; 84(16):6948-55. doi: 10.1021/ac300372p.  
745 Epub 2012 Aug 6. PMID: 22873683 [Indexed for MEDLINE]
- 746 51. Fuchs SW, Grundmann F, Kurz M, Kaiser M, Bode HB Fabclavines: bioactive peptide-  
747 polyketide-polyamino hybrids from *Xenorhabdus*. *Chembiochem.* 2014 Mar  
748 3;15(4):512-6. doi: 10.1002/cbic.201300802. Epub 2014 Feb 13 PMID: 24532262 DOI:  
749 10.1002/cbic.201300802 [Indexed for MEDLINE]
- 750 52. Kulkarni MM, McMaster WR, Kamysz W, McGwire BS. Antimicrobial peptide-induced  
751 apoptotic death of leishmania results from calcium-dependent, caspase-independent

- 752 mitochondrial toxicity. *J Biol Chem.* 2009 Jun 5; 284(23):15496-504. doi:  
753 10.1074/jbc.M809079200. Epub 2009 Apr 8. PMID: 19357081 PMCID: PMC2708846  
754 DOI: 10.1074/jbc.M809079200 [Indexed for MEDLINE] Free PMC Article
- 755 53. Marr AK, McGwire BS, McMaster WR. Modes of action of Leishmanicidal antimicrobial  
756 peptides. *Future Microbiol.* 2012 Sep;7 (9):1047-59. doi: 10.2217/fmb.12.85.  
757 Review.PMID: 22953706. [Indexed for MEDLINE]
- 758 54. Kaya H Aguilera MM, Alumai A, Choo HY, de la Torre M, Fodor A. et al., Status of  
759 entomopathogenic nematodes and their symbiotic bacteria from selected countries or  
760 regions of the world. *Biol Cont.* **38** (2006): 134–155 PMID: 15782660 DOI:  
761 10.1556/AVet.53.2005.1.7 [Indexed for MEDLINE]
- 762 55. Tóth EM, Márialigeti K, Fodor A, Lucskai A, Farkas R. Evaluation of efficacy of  
763 entomopathogenic nematodes against larvae of *Lucilia sericata* (Meigen, 1826)  
764 (Diptera: Calliphoridae). *Acta Vet Hung.* 2005;53(1):65-71. PMID: 15782660 DOI:  
765 10.1556/AVet.53.2005.1.7 [Indexed for MEDLINE]
- 766 56. Fodor A Fodor AM Forst S Hogan JS Klein MG Lehoczky É. Comparative analysis of  
767 antibacterial activities of *Xenorhabdus* species on related and non-related bacteria in  
768 vivo. *J Microbiol Antimicrobials* 2010; 2(3): 30-35.
- 769 57. Romond, C., Beerens, H., Criquelion, J. and Lepage, C. De'nombrement en milieu liquide  
770 de *C. perfringens* dans les viandes analysées cinq fois, après dilution décimale de la  
771 viande. *Annales des Falsifications de l'Expertise Chimique et de Toxicologie*  
772 1981; 74: 181–184.
- 773 58. Current Trends in Consumption of Animal Products. Chapter 1 In: Designing Foods:  
774 Animal Product Options in the Marketplace. National Research Council (US) Committee  
775 on Technological Options to Improve the Nutritional Attributes of Animal Products.  
776 Washington (DC): National Academies Press (US); 1988.
- 777 59. Harmon SM, Kautter DA, Peeler JT. Improved medium for enumeration of *Clostridium*  
778 *perfringens*. *Appl Microbiol.* 1971; 22(4): 688–692. PMCID: PMC376387
- 779 60. Amin A1, Bilic I, Berger E, Hess M. *Trichomonas gallinae*, in comparison to  
780 *Tetratrichomonas gallinarum*, induces distinctive cytopathogenic effects in tissue  
781 cultures. *Vet Parasitol.* 2012 May 25; 186(3-4):196-206. doi:  
782 10.1016/j.vetpar.2011.11.037. Epub 2011 Nov 20. PMID: 22172581 DOI:  
783 10.1016/j.vetpar.2011.11.037 [Indexed for MEDLINE]
- 784 61. Ibrahim HR, Tatsumoto S, Ono H, Van Immerseel F, Raspoet R, Miyata T. A novel  
785 antibiotic-delivery system by using ovotransferrin as targeting molecule *Eur J Pharm*  
786 *Sci.* 2015 Jan 23;66:59-69. doi: 10.1016/j.ejps.2014.10.005. Epub 2014 Oct 12.  
787 PMID:25315410 [Indexed for MEDLINE]
- 788 62. Lanckriet A, Timbermont L, De Gussem M, Marien M, Vancraeynest D, Haesebrouck F,  
789 Ducatelle R, Van Immerseel F. The effect of commonly used anticoccidials and  
790 antibiotics in a subclinical necrotic enteritis model. *Avian Pathol.* 2010 Feb; 39(1):63-8.  
791 doi: 10.1080/03079450903505771. PMID: 20390538. [Indexed for MEDLINE]
- 792 63. Li W, Tailhades J, O'Brien-Simpson NM, Separovic F, Ötvös L. Jr, Hossain MA, Wade JD.  
793 Proline-rich antimicrobial peptides: potential therapeutics against antibiotic-resistant  
794 bacteria. *Amino Acids.* 2014 Oct; 46(10):2287-94. doi: 10.1007/s00726-014-1820-1.  
795 Epub 2014 Aug 21. Review. PMID: 25141976
- 796 64. Ötvös L. Jr, Flick-Smith H, Fox M, Ostorházi E, Dawson RM, Wade JD The designer  
797 proline-rich antibacterial peptide A3-APO prevents *Bacillus anthracis* mortality by  
798 deactivating bacterial toxins. *Protein Pept Lett.* 2014 Apr; 21(4):374-81. PMID:  
799 24164262

- 800 65. Zheng X, Wang X, Teng, Mao R, Hao Y, Yang N, Zong L, Wang J. Mode of action of  
801 plectasin-derived peptides against gas gangrene-associated *Clostridium perfringens*  
802 type A. PLoS One. 2017 Sep 21; 12(9):e0185215. doi: 10.1371/journal.pone.0185215.  
803 eCollection 2017. PMID: 28934314 Free PMC Article
- 804 66. Ducatelle R, Eeckhaut V, Haesebrouck F, Van Immerseel F A review on prebiotics and  
805 probiotics for the control of dysbiosis: present status and future perspectives. Animal.  
806 2015 Jan;9(1):43-8. doi: 10.1017/S1751731114002584. Epub 2014 Oct 22. Review.  
807 PMID:25336177 [Indexed for MEDLINE]
- 808 67. Skrivanova E Van Immerseel F Hovorkova P Kokoska L. In Vitro Selective Growth-  
809 Inhibitory Effect of 8-Hydroxyquinoline on *Clostridium perfringens* versus  
810 Bifidobacteria in a Medium Containing Chicken Ileal Digesta. PLoS One. 2016 Dec  
811 9;11(12):e0167638. doi: 10.1371/journal.pone.0167638. eCollection 2016. PMID:  
812 27936245 Free PMC Article
- 813 68. Eeckhaut V, Wang J, Van Parys A, Haesebrouck F, Joossens M, Falony G, Raes J, Ducatelle  
814 R, Van Immerseel F. The Probiotic *Butyrivibrio pullicaecorum* Reduces Feed  
815 Conversion and Protects from Potentially Harmful Intestinal Microorganisms and  
816 Necrotic Enteritis in Broilers. Front Microbiol. 2016 Sep 21; 7:1416. eCollection 2016.  
817 PMID: 27708624 Free PMC Article
- 818 69. Geeraerts S, Delezie E, Ducatelle R, Haesebrouck F, Devreese B, Van Immerseel F.  
819 Vegetative *Bacillus amyloliquefaciens* cells do not confer protection against necrotic  
820 enteritis in broilers despite high antibacterial activity of its supernatant against  
821 *Clostridium perfringens* in vitro. Br Poult Sci. 2016 Jun; 57(3):324-9. doi:  
822 10.1080/00071668.2016.1169246. PMID:27122203.
- 823 70. Gholamiandehkordi A, Eeckhaut V, Lanckriet A, Timbermont L, Bjerrum L, Ducatelle R,  
824 Haesebrouck F, Van Immerseel F. Antimicrobial resistance in *Clostridium perfringens*  
825 isolates from broilers in Belgium. Vet Res Commun. 2009 Dec;33(8):1031-7. doi:  
826 10.1007/s11259-009-9306-4. PMID:19597952
- 827 71. Ngamwongsatit B, Tanomsridachchai W, Suthienkul O, Urairong S, Navasakuljinda W,  
828 Janvilisri T. Multidrug resistance in *Clostridium perfringens* isolated from diarrheal  
829 neonatal piglets in Thailand. Anaerobe. 2016 Apr;38:88-93. doi
- 830 72. Aradhyula S, Manian FA, Hafidh SA, Bhutto SS, Alpert MA. Significant absorption of oral  
831 vancomycin in a patient with *Clostridium difficile* colitis and normal renal function.  
832 South Med J. 2006 May; 99(5):518-20. PMID: 16711316 DOI:  
833 10.1097/01.smj.0000216477.06918.a3 [Indexed for MEDLINE]  
834

835 **ABBREVIATIONS**

836 **List of Abbreviations**

837 **AMP** = Antimicrobial Peptides

838 **CFCM** = cell-free culture media

839 **EMA** = *Xenorhabdus budapestensis*, (obligate bacterium symbiont of the nematode  
840 *Steinernema bicornutum* but can easily be grown in vitro, even in supplemented chicken  
841 food)

842 **EMC** = *Xenorhabdus szentirmaii*, (obligate bacterium symbiont of the nematode  
843 *Steinernema rarum* but can easily be grown in vitro, even in supplemented chicken food)

844 **EPB** = entomopathogenic (nematode-symbiotic) bacterium

845 **EPN** = entomopathogenic nematode

846 **GI** = gastro-intestinal system

847 **MDR** = multi drug resistance (multiple antibiotic resistance)