

COMPARATIVE STUDY OF THE ALKYLSULPHATASE ACTIVITIES OF BACTERIA FOUND IN SOIL CONTAMINATED WITH DETERGENT IN ONDO STATE, NIGERIA

ABSTRACT

Aims: To isolate, characterize and identify detergent degrading bacteria from detergent contaminated soil in Ondo State, Nigeria and also to compare and quantify enzyme production and biodegrading potentials of each of the bacterial isolates.

Place and Duration of Study: Ondo state, Nigeria, between February and July, 2017.

Methodology: Detergent degrading bacteria were isolated from detergent contaminated soil samples by supplementing minimal salt media with test surfactant. The bacteria isolated were subjected to enzyme analysis to study the alkylsulphatase enzyme production/activity in relation to growth pattern.

Results: Some bacterial isolates showed remarkable potential for alkylsulphatase production. In the enzyme study, *Bacillus subtilis* (1.53 mM/min), *Pseudomonas putida* (1.36 mM/min) and *Pseudomonas fluorescens* (1.33 mM/min) showed better enzymatic action. *Bacillus subtilis* showed the highest enzymatic activity of 1.53 mM/min.

Conclusion: It can be concluded that *Bacillus subtilis*, *Pseudomonas putida* and *Pseudomonas fluorescens* can be found in soil environment polluted with detergent. They are capable of surviving the toxic effect of the pollutant and efficiently producing alkylsulphatase; thus can be employed in enzyme production. They are capable of degrading detergent as a pollutant; thus can be utilized in the bioremediation of soil environments contaminated with surfactants

Keywords: Bioremediation, soil, alkylsulphatase, detergent, enzyme.

INTRODUCTION

Soil is a mixture of minerals, organic matter, gases, liquids and countless organisms that support life on earth. Soil continually undergoes development by way of numerous physical, chemical and biological processes, which include weathering with associated erosion. The soil functions as a medium for plant growth [1], it purifies, stores and supply water [2], and it influences distribution of plant species and provides a habitat for a wide range of organisms [3]. Soil is fundamental to human life on earth. Most plant requires a soil substrate to provide water and nutrients, and whether we cultivate the plants directly or consume animals that feed on the plants; we don't eat without soil [3]. Soil pollution is typically caused by industrial activity, chemicals used in agriculture and improper disposal of waste. Contaminants in the soil have major consequences on human health [3]. Long term exposure to polluted soil affects the genetic makeup of the body and may cause congenital illness and chronic health diseases.

Detergent is one of the major pollutants found in the soil after being used mostly in laundry processes [3]. Surfactants are routinely deposited in numerous ways on land and into water systems, whether as part of an intended process or as industrial and house hold waste causing pollution [4]. They are known to be toxic to animals, ecosystems and humans, and can increase the diffusion of other environmental contaminants [4]. Biodegradation of surfactants is being performed by soil or aquatic microorganisms leading to the generation of water, biomass, salts and carbon (iv) oxide gas [5]. The alkylsulphatase enzyme produced by some microorganism is involved in the biodegradation of detergents, which hydrolyses inorganic sulphate from its ester linkage with alcohols, the later being readily assimilated through normal metabolic pathways [6].

This research therefore, assess the biodegrading capabilities of bacteria isolated from soil contaminated with detergents on surfactants, in Ondo State, Nigeria by comparing the alkylsulphatase activities of each bacterial isolate.

METHODOLOGY

Collection of Samples

Soil samples were collected from carwash parks, in selected towns in Ondo State. The samples were collected in sterile containers, labelled and transported to the laboratory for Analysis.

Isolation of Detergent Degrading Bacteria

Isolation of detergent degrading bacteria from the soil samples was done by collecting the soil samples in sterile containers from the carwash parks; where the waste water effluent is being deposited. Serial dilutions were carried out on the soil samples. The serial diluted samples were inoculated onto minimal salt composition media supplemented with test surfactant. The inoculated plates were incubated aerobically at 28°C for 48hrs. At the end of the period of incubation, the plates were checked for growth [7]. The cultural characteristics of pure culture were noted for bacterial characterization [8]. The bacterial isolates were subjected to Gram's reaction and biochemical tests to identify the isolates.

Determination of Alkylsulphatase Production

Preparation of Enzyme Extract

Minimal salt composition media was prepared in broth form and supplemented with SDS at 0.01%, and it was inoculated with the bacterial isolates. The culture broth was incubated in an orbital shaker at 150 rpm. Fifty millilitres of the broth culture was collected at the end of twelve hours and it was centrifuged for 15 minutes at 4°C. The supernatant was decanted off. The cell pellets at the base of the centrifugation tube was collected using one millilitre (1ml) of tris buffer. The pellets were homogenized for 15 minutes. The homogenized pellets were then centrifuged for 15 minutes at 4°C. The supernatant was decanted and kept for the enzyme assay [9].

Alkylsulphatase Enzyme Assay

Four hundred and fifty micro litres (450 µl) of fifty millimolar (50 mM) Tris-hydrochloric acid (pH 7.5) and five hundred micro litres (500 µl) of one hundred millimolar (100 mM) SDS was pipette into a container of fifty micro litres (50 µl) of the enzyme. It was then incubated for a period of time (15 minutes). One hundred micro litres (100 µl) of the mixture, 9.9 ml of distilled water, two and a half millilitres (2.5 ml) of methylene blue solution and one millilitre (1ml) of chloroform was pipette into a separating funnel and shaken vigorously for 40 seconds. A chloroform layer was formed. The chloroform layer formed was carefully collected and the absorbance which indicates the quantity of enzyme produced was read at 652nm [9].

Analysis of Data

Data obtained were subjected to descriptive one way analysis of variance, using SPSS version 16 Microsoft windows 7 and treatment means were separated with Duncan's Multiple Range Test.

RESULTS AND DISCUSSION

The detergent degrading bacteria isolated from the contaminated soil were *Xanthomonas campestris*, *Bacillus subtilis*, *Pseudomonas putida*, *Bacillus panthoteticus*, *Bacillus funiculus*, *Echerichia coli*, *Pseudomonas haloplanktis*, *Bacillus cereus*, *Pseudomonas fluorescens* and *Bacillus anthracis*. Some of which were isolated in other related research [10] and [4]. Figure 1 depicts the enzyme activity of *Xanthomonas campestris* having its highest enzyme activity as 0.70 mM/min, while its optical density was 0.9 at this point. Figure 2 illustrates the enzyme activity of *pseudomonas putida* having its highest enzyme activity as 1.36 mM/min, its optical density was 0.15 at this point. Figure 3 shows the enzyme activity of *Escherichia coli*, the highest enzyme activity of *Escherichia coli* was 0.70 mM/min and its optical density was 0.99 at this point. Figure 4 depicts the enzyme activity of *Bacillus subtilis* having its highest activity as 1.53 mM/min at an optical density of 1.56. Figure 5 depicts the enzyme activity of *Klebsiella oxytoca*, it was able to produce an highest enzyme activity of 0.95 mM/min at an optical density of 0.83.




Fig. 1. Alkylsulphatase activity (AST) of *Xanthomonas campestris*




Fig. 2. Alkylsulphatase activity (AST) of *Pseudomonas putida*




Fig. 3. Alkylsulphatase activity (AST) of *Escherichia coli*




Fig. 4. Alkylsulphatase activity (AST) of *Bacillus subtilis*






Fig. 5. Alkylsulphatase activity of *Klebsiella oxytoca*

From Figure 6, *Proteus mirabilis* was able to produce an enzyme activity of 0.78 mM/min, which was the highest. The optical density at this point was 0.87. Figure 7 depicts the enzyme activity of *Bacillus cereus*, it was able to produce an enzyme activity of 1.05 mM/min, which was the highest. Its optical density at this point was 0.90. From Figure 8, *Pseudomonas fluorescens* produced an enzyme activity of 1.33 mM/min, which was its highest. The optical density was 1.68 at this point. From figure 9, *Bacillus anthracis* was able to produce an enzyme activity of 0.92 mM/min, which was its highest, while its optical density at this point was 0.60.




124 Fig. 6. Alkylsulphatase activity (AST) of *Proteus mirabilis*




125

126 Fig. 7. Alkylsulphatase activity (AST) of *Bacillus cereus*



127

128 Fig. 8. Alkylsulphatase activity (AST) of *Pseudomonas fluorescens*



129

130 Fig. 9. Alkylsulphatase activity (AST) of *Bacillus anthracis*

The bacterial isolates were able to produce the alkylsulphatase enzyme; possessing the mechanisms to carry out biodegradation of surfactants. There were variations in the quantity of alkylsulphatase enzymes produced by the bacterial isolates. Some of the bacteria showed better biodegrading potentials, and this could be as a result of the genetic makeup of the microorganisms [9]. Increase in optical density was an index of microbial growth. Growth pattern increased with increase in enzyme production. The results suggest that bioremediation by the bacterial isolates are promising for the biodegradation of surfactants as pollutants in the soil environment.

CONCLUSION

The study was able to illustrate the pattern of enzyme production and activity of the various isolates with respect to time and microbial growth. The study indicates an array of bacteria that could be selected for the remediation of soil environment contaminated with detergent. The study indicates that enzyme activity increases with time and microbial growth. It can be concluded that *Bacillus subtilis*, *Pseudomonas putida* and *Pseudomonas fluorescens* can be found in soil environment polluted with detergent. They are capable of producing alkylsulphatase; thus can be employed in enzyme production. They are capable of surviving the toxic effect of the pollutant, being able to break down the surfactant molecule and utilize it for their own growth; thus they can be applied in the bioremediation of environments contaminated with detergent.

REFERENCES

1. Dominati, E., Patterson, M. and Mackay, A. A framework for classifying and quantifying the natural capital and ecosystem services of soils. *Ecological Economics*. 2010; 69(9):1858-68.
2. House, C.H., Bergmann, B.A., Stomp, A.M. and Frederick, D.J. Combining constructed wetlands and aquatic and soil filters for reclamation and reuse of water. *Ecological Engineering*. 1999;12(1-2):2738.
3. Mishra, R.K., Mohammad, N. and Roychoudhury, N. Soil pollution: Causes, effects and control. *Tropical Forest Research Institute*. 2015; 3(1): 20-30.
4. Emmanuel, E., Hanna, K., Bazin, C., Keck, G., Clement, B. and Perrodin, Y . Fate of glutaraldehyde in hospital wastewater and combined effects of glutaraldehyde and surfactants on aquatic organisms. *Environment International*. 2005; 31 (3): 399–406.
5. Schleheck, D. ,Dong,W. and Denger, K . An Alphaproteobacterium Converts Linear Alkylbenzenesulfonate sulfophenylcarboxylates and linear alkyldiphenyletherdisulfonate surfactants into sulfodiphenylethercarboxylates. *Applied and Environmental Microbiology*.2000; 66, 1911-1916.
6. Toesch, M., Schober, M. and Faber, K. Microbial alkyl- and aryl- sulphatase: mechanism, occurrence, screening and stereoselectivities. *Applied Microbiology and Biotechnology*. 2014; 98:1487.
7. Sushma, P., Anvita, A., Nismitha, S. and Melwyn, S. Degradation of Anionic Surfactants by *Bacillus subtilis* and *Bacillus cereus*. *Journal of pharmacy and biological science*. 2012; 3(1):42-45.
8. Okpokwasili, G.O. and Nwabuzor, C.N. Primary biodegradation of anionic surfactants in laundry detergents. *Chemosphere*. 1998.17: 2175 – 2182.
9. Arotupin, D.J and Yusuf, A. Bacteria Associated with Selected Rivers in Akure, Nigeria and their Alkylsulphatase Activity/Production. *British Microbiology Research Journal*. 2016;13(6):1-7.
10. Ojo, O.A. and Oso, B.A. Isolation and characterization of synthetic detergent-degraders from wastewater. *African Journal of Biotechnology*. 2008; 7: 3753 – 3760.