

1 **Role of Optical Coherence Tomography In Macular Disorders**

2 **Abstract**

3 Optical coherence tomography (OCT) is an emerging technology for performing high-
4 resolution cross-sectional imaging. OCT is analogous to ultrasound imaging, except that
5 it uses light instead of sound. OCT can provide cross-sectional images of tissue structure
6 on the micron scale *in situ* and in real time.

7 High resolution cross-sectional imaging of the retina is useful for identifying, monitoring and
8 quantitatively assessing macular diseases. Optical coherence tomography (OCT) uses low
9 coherence or white light interferometer to perform high resolution measurements and imaging.

10 Optical coherence tomography (OCT) has a great potential as a diagnostic and research tool.

11 However it has its own limitations in the presence of media opacities like vitreous hemorrhage,
12 corneal edema and dense cataract. Hence, an attempt has been made to study the role of
13 Optical coherence tomography in diagnosis of macular disorders.

14 15 **Introduction**

16 The macula in retina is responsible for sharp, clear central vision and the ability to perceive
17 color. The tightly packed photoreceptor cells in the macula control all of the eye's central vision
18 and are responsible for the ability to read, drive a car, watch television, see faces, and
19 distinguish detail. There are two types of photoreceptor cells in the retina, rods and cones. The
20 rods provide vision at low light levels, while the cones provide sharp vision and discrimination.

High resolution cross-sectional imaging of the retina is useful for identifying, monitoring and quantitatively assessing macular diseases. Image resolutions of 1 to 15 μm can be achieved one to two orders of magnitude higher than conventional ultrasound. Cross-sectional images of the retina are obtained at the resolution of 1-10 micron. Optical coherence tomography (OCT) uses low coherence or white light interferometer to perform high resolution measurements and imaging. The high resolving power (10 μm - Time Domain, 5 μm – Spectral Domain) provides excellent detail for evaluating the vitreo-retinal interface, neurosensory retinal morphology, and the RPE-choroid complex. The infra-red light beam has a wave length of 820nm. OCT has made it possible to understand better the vitreoretinal relationship at the macula. Optical coherence tomography (OCT) has a great potential as a diagnostic and research tool. However it has its own limitations in the presence of media opacities like vitreous hemorrhage, corneal edema and dense cataract, which attenuate the incident and reflected light. Also, it cannot assess function and can only provide details about structural alterations. Hence, an attempt has been made to study the role of Optical coherence tomography in diagnosis of macular disorders. [1,2]

Literature

Anatomy of the macula [1,2]

The macula lutea is an oval, yellowish area of the center of the posterior part of the retina. It measures about 5.5 mm in diameter and lies about 3mm to the lateral side of the optic disc.

The yellowish coloration of the macula is caused by a yellow carotenoid pigment, xanthophylls, which is present in the retinal layers from the outer nuclear layer inward.

Parts of the macula [1,3,4]

- ❖ Umbo
- ❖ Foveola
- ❖ Foveal avascular zone
- ❖ Fovea
- ❖ Parafoveal area
- ❖ Peri foveal area

Umbo

It is a tiny depression in the very centre of the Foveola .which corresponds to the ophthalmoscopically visible foveolar reflex.

Foveola (350microns)

It is the small central region in which the thickness of the retina is reduced so as to contain only photoreceptors, glial cells and Mullers cells.

Foveal avascular zone (800 microns)

It is located inside the fovea but outside the foveola.

Fovea (1500microns)

It is a small depression where the retina is reduced to about half its normal thickness moving towards the centre of the retina. The inner nuclear layer is reduced to a double row of cells at the edge of the fovea.

The foveal sides of the depression are called the clivus ,the floor of the depression is the foveola. There are no blood vessels overlying the fovea and no rod cells in the floor of the fovea. It is here that there is the highest concentration of cones

Parafoveal area (500 microns)

It is characterized by the densest accumulation of the nerve cells in the entire retina especially ganglion cells and the inner nuclear layer. The outer boundary is the point where the ganglion cell layer has four rows of the nuclei.[3,4]

Peri foveal area (1500microns)

It ends where the ganglion cells are reduced to a single layer.[4,5]

Histology of the macula

The centre of the macula is the fovea containing the following layers

- Internal limiting membrane
- Outer plexiform layer
- Outer nuclear layer
- Layer of cones
- Retinal pigment epithelium.

Among the 301 patients, 428 images were studied, reviewed and diabetic macular edema, CSR, ARMD and occlusive diseases were diagnosed using SD-OCT, concluded that SD-OCT was useful in complementing the clinician's assessment of macular diseases. [6,7]

Among the 4 eyes in 4 patients of macular degeneration by optical coherence tomography (OCT). The series comprised one eye of Alport syndrome and 3 eyes of cone or cone-rod dystrophy [1]. In Alport syndrome, OCT showed thinning of sensory retina and high-reflective layer corresponding to retinal pigment epithelium and choriocapillaris. In 3 eyes of cone or cone-rod dystrophy, OCT showed absence of low-reflective layer corresponding to the photoreceptors. The foveal thickness in the 4 eyes ranged from 25 to 73 micron, average 56.0 $\pm$ 21.4 microns. It averaged 136.3 $\pm$ 12.3 microns in normal eyes. The difference was significant ($p < 0.05$). The diameter of macular depression averaged 1,452.8 $\pm$ 149.3 microns in the 4 eyes and 1,258.5 $\pm$ 47.1 microns in normal eyes. The difference was significant ($p < 0.05$). [1]

OCT scan protocols in macular disease [8,9]

The protocols that are helpful in macular diseases are the following:

1. Line scan

The line scan gives an option of acquiring multiple line scan without retaining to main window. Both the length and angle can be changed according to resolution.

2. Radial lines

3. Macular thickness map

98 **4. Fast macular thickness map**

99 **5. Raster line**

100 **6.Repeat**

101 **Principle [9,10]**

102 Optical coherence tomography (OCT) is an emerging optical imaging method, which allows the
103 acquisition of cross-sections in a non-destructive and contactless manner. Nowadays, OCT
104 systems provide axial resolutions ranging from 1 – 10 μm and allow real time in situ and in vivo
105 imaging.

106 Optical coherence tomography is based on the phenomenon of white light interferometry and
107 very sensitive to small changes in the refractive index of the sample. Therefore it delivers
108 complementary information to other imaging techniques like X-ray computed tomography (CT)
109 or magnetic resonance imaging (MRI). Near infrared light waves are partially reflected at
110 different depths within the sample, and their arrival times at a detector are interferometrically
111 compared with a reference wave. The detected signal contains information on the position of
112 the scatterers within the sample, on their reflectivity, velocity and polarisation properties.
113 Cross-sectional images can be reconstructed by collecting depth scans at different adjacent
114 positions.[10]

115

116

117 **Various types of OCT-**

118 Time domain OCT[9,10,11]

119

120 Frequency domain OCT (FD-OCT)(12,13)

121

122 Spatially encoded frequency domain OCT (spectral domain or Fourier domain OCT)

123

124 Time encoded frequency domain OCT (also swept source OCT)(17)

125

126 Optical coherence tomography

127 Optical coherence tomography (OCT) allows characterization of the profile and the intra-
128 retinal structural changes of many macular disorders .[13]

129 Diabetic macular edema [5,14]

130 This has five distinct patterns on OCT

131 ▪ Sponge-like retinal thickness

132 ▪ Cystoid macular edema

133 ▪ Subfoveal serous detachment

134 ▪ Vitreomacular traction

135 ▪ Taut posterior hyaloid

136 OCT is very useful in defining treatment options and for monitoring response to intervention

137 [14]

138 **Macular hole:** OCT is extremely use for assessing the following:

139 ▪ Early macular holes

- 140 ▪ Presence of PVD
- 141 ▪ Size of macular holes
- 142 ▪ Closure pattern following Vitrectomy [6]
- 143 **ARMD:** OCT has become very popular in ARMD because of early disease detection, monitoring
- 144 of therapy and early evidence of reactivation [9]

145 **ERM:** These are clearly visible on the vitreoretinal interface on OCT [11,12,13]

146 **CSR:** OCT allows detection of early CSR that is clinically observe .[12,13]

147

148 **Macular disorders**

149 **Age-related macular degeneration [16,17]**

150 Age-related macular degeneration (ARMD), also called senile macular degeneration, is a

151 bilateral disease. It is a leading cause of blindness in developed countries, in population above

152 the age of 65 years.

153 It is of two types

- 154 • Non-neovascular(Dry)
- 155 • Neovascular(Wet or exudative)

156 **Non-neovascular ARMD**

157 **Drusen**

158

159 Drusen can be hard drusen or soft drusen.

160 **Hard drusen in OCT**

161 On OCT they are seen as focal elevation of the RPE with mildly thinned overlying retina. The
162 elevated RPE does not shadow the reflection.

163 **Soft drusen in OCT** images show small modulation in the contour of the retinal pigment
164 epithelium with lack of shadowing below the contour changes, similar to serous RPE
165 detachment. However, unlike a PED, soft drusen have shallow margins and mild back scatter
166 into the choroid.

167 **Geographic atrophy**

168 OCT shows a thinned out overlying retina and decreased reflectivity of RPE with enhanced
169 choroidal layer.

170 **Neovascular ARMD (exudative)**

171 OCT shows retinal thickening with mild back scattering from outer retinal layers, with
172 disruption in the RPE in an area of moderate reflectivity with fusiform thickening of the
173 reflective band corresponding to the CNVM. The lesions merge with RPE choriocapillaries
174 layer.[\[15,16\]](#)

175 **Occult CNVM**

176 OCT shows an enhanced optical scatter signal at the choroidal level extending into sub-RPE
177 space which is thicker than the adjacent reflection from the pigmented epithelium and

178 choriocapillaries, consistent with a neovascular membrane. The retina overlying the lesion
179 appears thickened and elevated, with reduced reflectivity, suggesting fluid accumulation.

180 **Retinal pigment epithelium detachment(PED) [10,17]**

181 PED can be differentiated into the following type

182 Serous PED

183 Fibro vascular PED

184 Hemorrhagic PED

185 Drusenoid PED

186 **Serous PED**

187 Serous PED-clinically seen as an area of RPE detachment.

188 **Fibro vascular PED**

189 OCT shows elevated RPE, along with mild to moderate back scatter persisting beneath
190 the RPE, consistent with the Fibro vascular PED. It may have associated overlying neurosensory
191 detachment or overlying cystoid macular edema.

192 **Hemorrhagic PED**

193 OCT through the hemorrhagic PED shows elevated RPE with moderate back scatter,
194 which attenuates quickly with depth below the RPE. This back scatter, appearing immediately
195 below the raised RPE is consistent with the hemorrhagic. The RPE band may be

196 indistinguishable from hemorrhagic. There may be associated neurosensory elevation and
197 cystoid changes in the overlying retina.

198 **Drusenoid PED**

199 OCT defines a RPE elevation with optical back scatter below the RPE, with small
200 modulations in the RPE contour with moderate optical reflectivity below the lesion.

201 **Feeder vessels**

202 OCT may reveal areas of increased back scatter corresponding to the scarring
203 membrane, with associated thickening of RPE choriocapillaries layer.

204 **Loculated fluid**

205 OCT shows a well defined region of hyper-reflectivity at the level of the RPE-
206 choriocapillaries, consistent with the CNVM. The neurosensory retina above the membrane
207 shows thickening as well as focal areas of reduced optical, corresponding with the fluid.

208 **RPE rip**

209 OCT through the lesion shows the elevated neurosensory retina and RPE which shadows
210 the reflections from the choroid below. In the area of RPE rip two highly reflective layers are
211 seen below the sensory retina corresponding with a folded double layer of pigment epithelium,
212 with attenuation of reflections below the layer. In the area corresponding with absent RPE,
213 enhanced reflectivity and optical penetration of the choroid is noted.

214 **Disciform scar**

OCT reveals a focal area of enhanced back scatter corresponding to the chorioretinal scar. The overlying neurosensory retina shows thinning throughout. (Natarajan et al 2005)

Central Serous ChorioRetinopathy (CSCR)[5,11]

OCT shows an elevated neurosensory retina with an optically clear space underneath and a reflective layer corresponding to the RPE and choriocapillaries underneath it, corresponding to the CSR. With the advent of OCT the existence of PED with CSR has been shown to be much more than thought previously. In cases of associated PED; below the elevated neurosensory retina lies an optically clear space below which a highly reflective dome shaped elevation with attenuated shadows below it suggest the presence of PED. There may be hyper reflective spots in the optically clear space below the neurosensory retina corresponding with the sub retinal exudation and fibrin seen in resolving or chronic CSCR. CSCR may also have associated choroidal neovascular membrane (CNVM). In such situations, there will be a delineated hyper fluorescence on angiogram corresponding with the CNVM. OCT will show high reflectivity at the level of RPE choriocapillaries suggestive of a CNVM.

Epi-retinal membrane (ERM)[8,12,13]

OCT is a very useful tool in grading the ERM, and providing information about the underlying retinal traction and secondary changes in the retina. OCT reveals a hyper-reflective membrane at the vitreoretinal interface, adherent to the underlying retina, with distortion of the inner retinal layer. The traction on the underlying retina is seen as folds in the inner retinal

layer. On OCT, ERMs are seen as clearly separable where clear space is visible between the epiretinal membrane and inner retinal surface and global ERM where no area of separation is seen between the ERM and inner retinal surface. An increased thickness of retina with optically empty spaces in retinal layers along with reduced back scattering from outer retinal layers will indicate Cystoid macular edema; a lamellar macular hole may be associated with ERM. A macular pseudo hole formed by the ERM needs to be differentiated from a true macular hole. A macular pseudo hole will show an abnormally steep foveal contour resembling a partial thickness macular hole, but the outer retinal layer is seen intact throughout the image ruling out a full thickness macular hole. It is important to distinguish an ERM from a detached posterior hyaloid as both appear as a reflective band anterior to the retina on OCT. The posterior hyaloid has a weak, patchy reflection that is inner than an ERM.

Macular hole [15,16,17]

In OCT the following are seen:

1. stage IA of the macular hole is seen as foveal pseudo cyst in the retinal layers.
2. stage IB is seen as partial disruption of the outer retinal layers. The posterior vitreous seen attached to its edges. It may develop to a lamellar macular hole.
3. stage II reveals as a full thickness macular dehiscence with attached posterior vitreous. The surrounding retina may show thickening and cystic changes.
4. Stage III .reveals a large macular hole with attached posterior vitreous.

5. Stage IV reveals a macular hole larger than 400μ in diameter, with a detached posterior hyaloid. There may be an associated optically empty space below the neurosensory retina at the edges of the dehiscence suggestive of sub retinal fluid, along with thickening and cystic changes in the retinal layer.[8]

Heredomacular degeneration (HMD)

Best's disease:

OCT shows a characteristic dome shaped lifting of retina including the RPE layer, with a high reflective deposit in the sub RPE space, corresponding to the accumulated lipo fuschin. This is in contrast to the presentation of a serous PED.

Foveal schisis

OCT is diagnostic as it reveals an elevation of the foveal pit with peri foveal splitting of the neurosensory retinal layers.

Stargardt's disease

OCT reveals thinned out neurosensory retina along with increased reflectivity of the RPE choriocappileries layer.

Retinitis pigmentosa

OCT reveals a thinned out neurosensory retina; associated CME may be seen, (Natarajan 2005).(8)

CONCLUSION

OCT can certainly play a vital role along with the clinical examination in the diagnosis and evaluation of macular disorders. OCT can be included as a part of evaluation of any disorder affecting the macula which can be performed and interpreted by an ophthalmologist. It is useful for both screening and monitoring macular diseases and for deciding intervention. It is useful for assessing response to intervention such as laser photo coagulation, intravitreal injections and surgery.

References

1. Chiako K, Optical coherence tomographic findings in atrophic macular disorders. Japanese journal of clinical ophthalmology;(2000) 54 (8): 1429-1433.
2. Gupta V, Amod Gupta. Atlas of Optical Optical Coherence Tomography of macular diseases. Jaypee, Medical publisher, India first edition, (2004).
3. Haouchine B, Massin P, Tadsyoni R, Erginay A and Gaudric A. Diagnosis of macular pseudoholes and lamellar macular holes by optical coherence tomography. American Journal of Ophthalmol(2004) 138:5
4. Hee MR, Puliafito CA, Duker Js et al. Topography of diabetic macular edema with optical coherence tomography, Ophthalmology (1998)105:360-70.
5. Hee MR, Puliafito CA, Wong C et al. Optical coherence tomography of central serous chorioretinopathy. AM J Ophthalmol (1995)120:65-73

6. Johnson MW. Improvements in understanding and treatment of macular holes, Curr Opin Ophthalmol (2002) 13: 152 -160.

7. Massin P, Duguid G, Erginay A, Haouchine B, Gaudric A, Optical Coherence Tomography for evaluating diabetic macular edema before and after Vitrectomy. American Journal of Ophthalmol (2003) 135:169-77.

8. Natarajan S, Sharma S, Hitendra B Mehta HB. Diagnostics in macular disorders. Jaypee, Medical publisher, India , First edition (2005)

9. Rogers AH, Martidis A, Greenberg PA, Puliafito CA. optical coherence tomography findings following photo dynamic therapy of choroidal neovascularisation, Am J Ophthalmol (2002) 134:566-576

10. Singh M. Spectral domain optical coherence tomography imaging of retinal diseases in Singapore; Ophthalmic surgery, Lasers and imaging, (2008);39:S120-S125.

11. Wilkins JR, Puliafito CA, Hee MR et al. Characterization epiretinal membranes using optical coherence tomography. Ophthalmology. (1996)103:2142- 2151.

12. A. F. Fercher and E. Roth, "Ophthalmic laser interferometry. Proc. SPIE vol. 658, pp. 48-51. 1986.

13. Fercher, AF; Mengedocht, K; Werner, W (1988). "Eye-length measurement by interferometry with partially coherent light." **13** (3): 186–8.


14 .A. F. Fercher, "Ophthalmic interferometry," Proceedings of the International

Conference on Optics in Life Sciences, Garmisch-Partenkirchen, Germany, 12–16 August 1990.


15. Huang, D; Swanson, EA; Lin, CP; Schuman, JS; Stinson, WG; Chang, W; Hee, MR; Flotte, T; et al. (1991). "Optical coherence tomography". *Science* **254** (5035): 1178–81.

16. Zysk, AM; Nguyen, FT; Oldenburg, AL; Marks, DL; Boppart, SA (2007). "Optical coherence tomography: a review of clinical development from bench to bedside.". *Journal of biomedical optics* **12** (5): 051403.


17. A. F. Fercher, C. K. Hitzenberger, W. Drexler, G. Kamp, and H. Sattmann, " In Vivo Optical Coherence Tomography," *Am. J. Ophthalmol*, vol. 116, no. 1, pp. 113-114. 1993.




CME before Treatment




CME after Treatment




CNVM before Treatment




CNVM after Treatment Fig (1)




CSR before Treatment



CSR after Treatment




Macular Edema before Treatment




Macular Edema after Treatment


Fig (2)




ERM before Treatment



ERM after Treatment



Macular Hole before Treatment



Macular Hole after Treatment

Fig (3)

Source of pictures: taken from the databse of patients treating for the respective diseases.

376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550