

EFFECT OF COLOUR VISION ON OCULAR DEVIATION

Abstract

Purpose: The aim of the present study is to correlate effect of colour vision in the presence of ocular deviation.

Methods: A pilot, cross sectional, observational study was performed at tertiary eye care centers. Subjects with Ocular deviation between 10 to 40 prism diopters, Corrected distance Visual Acuity should be greater than 6/18 and Age should be between 10 to 40 years of age were included in the study. Colour vision was assessed with Fransworth Munsell D 15 colour vision test.

Results: 30 subjects were included in the study. Out of that, 16 subjects were in the age group of 11-20 years, 12 subjects were in the age group of 21-30 years and 2 subjects were in the age group of 31-40 years. 60% subjects were Female and 40% subjects were Male. Mean scores of vision parameters were taken. In 50 % of cases, colour vision was found normal, 33 % were having mild tritanopia and 17% were having moderate tritanopia.

Conclusions: In cases of ocular deviation, mild to moderate tritanopia can occur.

Keywords: Ocular Deviation, colour vision

INTRODUCTION:

Colour vision plays an important role in Binocular Single Vision. For colour vision, cone cells are precisely responsible. In cases of normal ocular deviation, images of an object fall on the foveal region. Due to anatomical deformity,

number of cone cells varied from region to region in macular area. That's why, in the ocular deviation, there are chances of deterioration of colour vision in the presence of ocular deviation. Image of an object will fall on the para foveal region in cases of ocular deviation, so there are chances that colour vision will deteriorate.

METHODOLOGY:



A pilot, cross sectional, observational study was performed at tertiary eye care centers. Subjects with Ocular deviation between 10 to 40 prism diopters, Corrected distance Visual Acuity should be greater than 6/18 and Age should be between 10 to 40 years of age were included in the study. Individuals with any other systemic disease(specially which can affect study), Individuals with any other Ocular Pathology, with any active ocular infection, any ocular anomalies like Corneal Scar etc, ocular deviation if less than 10 degree and Significant amount of amblyopic patient were excluded from the study. Full refractive correction along with detailed fundus evaluation was performed in each and every patient. Colour vision was assessed with Fransworth Munsell D 15 colour vision test.



44

RESULTS:


30 subjects were included in the study. Graph 1 shows distribution of subjects in various age groups. 16 subjects were in the age group of 11-20 years, 12 subjects were in the age group of 21-30 years and 2 subjects were in the age group of 31-40 years. Graph 2 shows gender wise distribution of the subjects. 60% subjects were Female and 40% subjects were Male. Mean scores of visual parameters were taken using SPSS Software version 20. Graph 3 shows 50 % of cases, colour vision was found normal, 33 % were having mild tritanopia and 17% were having moderate tritanopia.



54


55

56 GRAPH 1: SHOWS AGE WISE DISTRIBUTION OF THE SUBJECT



57

58 GRAPH 2 shows gender wise distribution of the subjects



59

60


61

62

63

64

65 Graph 3: shows effect of colour vision in the presence of ocular deviation.



66

67

68 Discussion:

69 Colour vision was tested with Fransworth Munsell D-15 colour vision test.

70 Here 30 subjects were evaluated. Out of that 15 were having normal colour
71 vision, 10 were having mild tritanopia and 5 were having moderate tritanopia.

72 Severe tritanopia was not found in any patient. According to statistical analysis,

73 colour vision will deteriorate significantly with increase in ocular deviation. It

74 mainly occurs due to number of cone cell variation in the macular region. The

75 number of cone cells are highest in foveal region compare to parafoveal region

76 Cone cell is completely responsible for Colour vision, so any anomaly in cone

77 cells will lead to deterioration of colour vision.

78 Conclusion :

79 In cases of ocular deviation, mild to moderate tritanopia can occur.

80 CONSENT :

81 Oral/ written consent was obtained from patient as well as from tertiary eye care
82 centres.

83 ETHICAL APPROVAL:

84 It is not applicable.

85 References:

- 86 1. HANDBOOK OF PEDIATRIC STRABISMUS AND AMBLYOPIA,
87 Kenneth W. Wright, Peter H. Spiegel, Lisa Thompson, First ed, 2006
- 88 2. Hui Zhu et al., “Association between Childhood Strabismus and
89 Refractive Error in Chinese Preschool Children “Journal of Plos One,
90 March 2015
- 91 3. Zhale Rajavi et al, “Prevalence of Colour Vision Deficiency and its
92 Correlation with Amblyopia and Refractive Errors among Primary School
93 Children”, Journal of Ophthalmic and Vision Research, 2015; vol. 10,
94 issue 2, pg 130-138...
- 95 4. Anika K. Tandon et al. “Binocular Inhibition in Strabismic Patients is
96 Associated with Diminished Quality of Life”, Journal of American
97 Association for Pediatric Ophthalmology and Strabismus, October 2014,
98 volume 18, issue 5, pg – 423-426
- 99 5. Ye et al, “Strabismus genetics across a spectrum of eye misalignment
100 disorders”, Journal of clinical genetics, 2014, vole 86, pg 103-111
- 101 6. A.G. KOCAK-ALTINTAS et al “Visual Acuity and Colour Vision
102 deficiency in Amblyopia” European Journal of Ophthalmology, 2000,
103 vol. 10, no.1, pg 77-81
- 104 7. ALAN W. FREEMAN et al. “Components of Visual Acuity Loss in
105 Strabismus” published in the journal of vision research, 1996, Vol. 36,
106 No. 5, Pg. 765-774

107