

AMETROPIAIN CHILDREN: A REVIEW OF 432 CASES

MAKITA Chantal ¹, NGANGA NGABOU Géraud Charles Freddy ¹, KOULIMAYA
Reinette Christelle ¹, DIATEWA Bénédicte ¹, MASSAMBA Alphonse ²

¹Ophthalmology Department, University and Hospital Center of Brazzaville,
Congo

²Biostatistics and Biosciences Department, Brazzaville Institute of Health and
Sports, Brazzaville –Congo

Authors' contributions

This work was carried out in collaboration between all authors. Authors MC, NNGCF, KRC
and DB were the doctors consulting patients. The author MA performed the statistical
analysis and writing of the study. The author MC was responsible supervising and writing the
study. All authors have read and agree with the final version.

Author corresponding:

Dr Chantal Makita

BP: 14626, Brazzaville-Congo

E-mail: chantalmakita@gmail.com

Original Research Article

Abstract

Aims: To determine the epidemiological and clinical aspects of ametropia and their impact on low
vision. **Study Design:** Retrospective: **Place and Duration of Study:** Department of Ophthalmology,
University and Hospital Center of Brazzaville Congo between January 2014 and December 2016.
Methodology: The records of 1,126 children aged 5–15 years who attended the ophthalmic
department were assessed. A refractive error was considered visual acuity less than 0.7(decimal
notation of visual acuity) on the measurement of visual acuity by far in monocular to 5
meters using optotypes adapted for age. **Results:** Of the 1,126 records, 432 children had a
refractive disorder, with a relative frequency of 38.3%. Of these, 242 (56%) were males, and 190
(44%) were females. Concerning the reasons for consultation, a headache ranked first 218 (50.5%),
followed by problems with visual acuity 106 (24.5%). Astigmatism was the most common type of
ametropia 348 (80.6%), followed by myopia 52 (12%) and farsightedness 32 (7.4%). Astigmatism was

present in 52.4% of boys and 28.2% of girls ($P<0.05$). Astigmatism was associated with myopia in 15.7% of cases and hyperopia in 7% of cases. The mean age of the patients was 10.4 years, with children aged 8–10 years accounting for the majority of cases ($n = 231, 53.4\%$). Visual acuity was ≤ 0.7 in 432 patients with refractive errors. Ametropia did not improve in 9 (2%) cases. There were 2 cases (0.5%) of anisometropia. In the majority of 296 (68.5%) cases, ametropia was not severe. In 225 (52.9%) ametropic cases, the astigmatism was the simple form. **Conclusion:** The prevalence of ametropia observed in this study points to the need for action to enable early detection and treatment.

Keywords: Child, visual acuity, ametropia, astigmatism

1. Introduction

The ametropia or a refractive error is most often accompanied by a decrease in visual acuity, with the severity depending on the underlying pathology [1]. In children, ametropia can have adverse effects on learning [1]. It can also impede social aspects of life. Worldwide, ametropia is the most common cause of declines in visual acuity and ophthalmic consultations [2]. Ametropia requires early detection and appropriate management [3]. In developing countries, loss of visual acuity is not life threatening. However, in under-developed countries, such as Tunisia, it poses a major threat. According to a study conducted in Tunisia in 2000, 57.2% of schoolchildren aged 6–15 years were ametropic [4]. The aim of the present study was to determine epidemiological and clinical aspects of ametropia and the impact of these on low vision in Congo Brazzaville.

2. Material and methods

2.1 Study population

We conducted a retrospective study of the records of patients who attended the Department of Ophthalmology, University and Hospital Center of Brazzaville Congo from 1 January 2014 to 31 December 2016. A data collection sheet was compiled based on information in patient records, including the reasons for consultations, medical histories and results of ophthalmic examinations. Children aged 5–15 years were included in the study. A subjective measure of visual acuity was used based on a logarithmic scale ≤ 0.7 decimal in at least one eye and spherical or cylindrical ametropia of $\geq 0.5D$. The exclusion criteria were: aged less than 5 and greater than 15 years, incomplete files, and existence of organic amblyopia. The study was approved by the ethics committee of the Medical

60 Society of Congo-Brazzaville. The variables analysed were age, sex, reason for consultation, visual
61 acuity and refractive power of each eye.

62 **2.2. Ophthalmic Examinations**

63 All the patients underwent an ophthalmological examination, including measurements of uncorrected
64 monocular visual acuity at a viewing distance of 5 m using age-appropriate optotypes; automatic
65 refractometry using a TOPCON AR-8900 refractometer before and after the induction of cycloplegia
66 (achieved using cyclopentolate eyedrops); anterior segment biomicroscopy; automatic tonometry; and
67 a fundus assessment.

68 **2.3 Definitions**

69 Ametropia was divided into three categories: myopia, hypermetropia and astigmatism. The simple
70 form was defined as positive or negative cylindrical refractive values, and the compound form was
71 defined as positive and spherical refractive values and negative or negative values. Three degrees of
72 severity were recognized: low spherical ametropia was considered refraction less than 3 D (dioptres),
73 moderate spherical ametropia was considered refraction between 3 and 6 D, and strong spherical
74 ametropia was considered refraction greater than 6 D, all at algebraic values [5]. Astigmatism was
75 considered low when it was less than 2 D, moderate when it was between 2 and 4 D and strong when
76 it exceeded beyond 4 D. Anisometropia was defined by a difference in refraction between two eyes
77 greater than 2 D [6].


78 **2.4 Statistical Analysis**

79 The analysis of the data was carried out using Epi Info, version 2012 software. To compare the
80 percentages of the various parameters the S test of Sokal was conducted. *P* values were considered
81 significant when the threshold of significance was ≤ 0.05 .

82 **3. Results**

83 Of a total of 1,126 patient records of children aged 5–15 years who presented during the study period,
84 432 children had a refractive disorder, a relative frequency of 38.3%. Of these, 242 (56%) patients
85 were males, and 190 (44%) were females, with no significant difference ($P > 0.05$). The average age of
86 the patients was 10.4 years (range: 5–15 years). As shown in Figure 1, a headache was the most
87 common reason for consultation 218 (50.5%) of cases, asthenopia was observed in 106 (24.5%) of

88 cases, a decrease in visual acuity was noted in 72(16.7%) of cases and 36(8.3%) wore glasses of
 89 correction for prior optical equipment
 90



92 **Fig.1 Frequency distribution of functional signs**

93 The frequency by type of ametropia and sex is presented in Table 1. Astigmatism was the most
 94 common ametropia (n=348, 80.6%), followed by myopia (n=52, 12%) and farsightedness (n=32,
 95 7.4%).

97 **Table 1. Distribution of ametropia by sex**

Sex	Male		Female		Total	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Hypermetropia	12	37.5	20	63.5	32	7.4
Myopia	30	57.7	22	42.3	52	12.0
Astigmatism	200	57.5	148	42.5	348	80.6
Total	242	56	190	44	432	100

98 *n = number of patients*

99 Table 2 shows the distribution of astigmatisms. Astigmatism was present in 52.4% of boys and 28.2%
 100 of girls ($P < 0.05$). Compound astigmatism 22.7% of cases, associated with near-sightedness in 15.7%
 101 and hypermetropia (7%); for a total of 27.7% cases of myopia and 14.4%, cases of hypermetropia.

102

Table 2. Distribution of astigmatism by sex

Sex	Male	Female	Total
	%	%	%
Simple	34.7	23.2	57.9
astigmatism			
Compound	12.2	3.5	15.7
myopic			
astigmatism			
Compound	5.5	1.5	7
hypermetropic			
astigmatism			
Total	52.6	28.2	80.6

105 The different types of ametropia, analysed by age group, are shown in Table 3. Ametropia was most
 106 common among those aged 8–10 years ($n = 231$, 53.4%) ($P < 0.05$). Compound astigmatism was
 107 common in those aged 10–13 years ($n = 128$, 36.7%).

Table 3. Distribution of type of ametropia by age group

Age range	5–7years		8–10years		11–13years		14–15years		Total	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Hypermetropia	1	3.1	20	6.5*	6	18.7	5	15.7	32	7.4
Myopia	2	3.8	25	48.1	20	38.5	5	9.6	52	12.0
Astigmatism	38	10.9	186	53.4*	74	21.3	50	14.4	348	80.6
Total	41	9.4	231	53.4	100	23.1	60	13.8	432	100

110 * $P < 0.05$

111 Visual acuity was ≤ 0.7 in 432 ametropia cases. It was not improved in 9 (2%) cases. There were 2
 112 (0.5%) cases of anisometropia. In terms of the severity of ametropia, it was classified as low in the
 113 majority 296 (68.5%) of cases. The astigmatism was the simple form in 225 (52.9%) cases.

4. Discussion

Ametropia is one of the most common pathologies encountered in schoolchildren, with a reported prevalence of 28–60%, depending on the series [3]. The frequency (38.3%) found in the present study was close to that (43.1%) reported in a previous study conducted in Cameroon [7]. Studies conducted in China and Chile reported a higher prevalence of 50.8% and 56.3%, respectively, [8, 9]. A study in Tunisia found a prevalence of 57.2% [4]. The high recorded prevalence of ametropia is likely attributed to the use of automatic refractometry measurements after the induction cycloplegia [4]. Other studies conducted in China (18.8%) [10], Benin (10.6%) [2] and Nepal (2.2%) [11] found low frequencies of ametropia.

In the present study, sex was not associated with the occurrence of ametropia in children. This finding is in accordance with that reported in studies conducted in Tunisia, Cameroon and Japan [4, 7, 12].

In this study, astigmatism was the most common type of refractive error (80.6%), especially the simple form (57.9%). Our observations are comparable with those of a previous study, which reported a prevalence of 91.9% for total astigmatism and 46.4% for simple astigmatism in a paediatric population in Benin [2]. On the other hand, studies conducted in Iran and Tunisia reported a predominance of hypermetropia (56.6% and 67%, respectively) [13, 14]. Disparities in the prevalence of ametropia are likely attributed to ethnic rather than genetic factors [15], as can be seen in the present study. In our study, the proportions of patients with myopia (27.7%) and hyperopia (14.4%) were similar to those reported in studies conducted in Benin (33.9% and 19.7%, respectively) [2] and the Democratic Republic of Congo (33% and 22%, respectively) [16]. In the present study, two boys for one girl have astigmatism, but this difference did not appear to be a risk factor for ametropia.

The frequency of ametropia in our study increased until the age of 10 years and then gradually decreased. The same trend was observed for each type of ametropia. A previous study found similar results, with no statistically significant relationship between age and ametropia ($P>0.05$) [2].

All children with visual acuity ≤ 0.7 are considered to have a refractive anomaly. However, subjective visual acuity alone is insufficient to make a diagnosis of ametropia. According to Roche [17], a large number of children with visual acuity ≤ 0.7 were emmetropic. The refraction objective by cycloplegia undoubtedly minimizes the risk of refractive error.

In the present study, nine (2%) cases of visual acuity could not be improved. These consisted of five cases of strong myopia, three cases of compound myopic astigmatism and one case of amblyogenic

all-round hypermetropia. Both cases of anisometropia were due to unilateral strong myopia. These findings high light the importance of early detection of ametropia and its best correction of refraction anomalies to prevent low vision.

In the majority of cases (68.5%), the severity of ametropia was classified as low,with 225 (52.9%) cases of ametropia due to simple astigmatism. Sounouvou and al. and Odoulami-Yehouessi and al. [2,5] presented similar findings, reporting, respectively, 89.3%, 45.5% and 71.8%, 17.1%.

5. Conclusion

Refraction anomalies are common in daily practice. In this study, they were present in 38.3% of cases, with a strong predominance of astigmatism (80.6%) and low ametropia (68.5%). Amblyopia (2%) was also detected. The prevalence of ametropia observed in this study points to the need for action to enable early detection and treatment.

Declaration of interest

The authors declare that they have no links of interest.

References

- 1- Bois C, BinotTM, Jonqua F, Guillemot G, Brémont-Gignac D. Screening visual disorders between 3 and 5 years: experience of the departmental service of maternal and child protection in the Hauts-de-Seine. J FrOphthalmol, 2007; 30: 570-6.
- 2- Sounouvou I, Chiabi S, Doutetien C, Sonon F, Yehouessi-Odoulami L, BassabiSK. Amétropies in primary school in Cotonou (Benin). J FrOphthalmol, 2008; 31: 771-5.
- 3- Jeddy-Blouza A, Loukil I, Mhenni A, Khayati L, Mallouche N, Zouari B. Management of the farsightedness of the child. J FrOphthalmol, 2007; 30: 255-9.
- 4- Ayed T, Sokkah M, Charfi O, and El Matri L. Epidemiology of refractive errors in schooled, socio-economically disadvantaged children in Tunisia. J Fr Ophthalmol, 2002; 25: 712-7
- 5- Odoulami-Yehouessi L, Chiabi S, Sounouvou I, Deguenon J, Doutetien C, Bassabi SK. The refraction of the child attending the CNHU Cotonou. Mal Med, 2005; 20: 24-6.
- 6- Shin YF, Hsiao CH, Wen SH, LL, CJ Chen, HungPT. Prevalence of anisometropia in taiwanese school children. J Formos Med Assoc, 2005; 104: 412-7.

176 7- OmgbwaEballé A, Assupta Bella L, Owona D, Mbome S, EbanaMvogo. The pathology of
177 the child from 6 to 15 years: hospital study of Yaoundé. Cah Santé, 2009; 19: 61-6.

178 8- He M, Zeng J, Liu Y, Xu J, Pokharel GP, Ellwein LB. Refractive error and visual
179 impairment in urban children in Southern China. Invest Ophthalmol Vis Sci, 2004; 45; 793-
180 9.

181 9- Maul E, Barroso ST, Munoz SR, Sperduto RD, Ellwein LB. Refractive error study in
182 children: Result from florida, chile. Am J Ophthalmol 2000; 129: 445-54.

183 10- Zhao J, Pan X, Sui R, Munoz SR, Sperduto RD, Ellwein LB. Refractive error study in
184 children: results from shunyi china district. Am J Ophthalmol 2000; 129: 427-35.

185 11- Pokharel GP, Negrel AD, Munoz SR, Ellwein LB. Refractive error study in children:
186 Result from Mechi area Nepal. Am J Ophthalmol 2000; 129: 436-44.

187 12- Matsumura H, Hirai H. Prevalence of myopia and refractive changes in students from 3 to
188 17 years of age. On Ophthalmol 1999; 44: 109-15.

189 13- Jeddi A, BelHadji A, Louane W, Hamoud M, Azaied S, Zouari B. Contribution of the
190 total optical correction after cycloplegia in headaches. J FrOphthalmol 2002; 25: 270-3.

191 14- Negrel AD, Maul E, Pokharel GP, Zhao J, Ellwein LB. Refereed error study in children:
192 Sampling and measurement methods for contrysurvey. Am J Ophthalmol 2000; 129: 525-7.

193 KaimbowaKaimbo D. Missotten L. Ocular refraction in Zaire. Bull Soc BelgianOphthalmol
194 1996; 261: 101-5.

195 16- Roche JJ. Assessments of visual impairments of five-year-old children in mayotte, Med
196 Trop 2005; 65: 465-72.

197