

Case study**Evaluation of Heavy metals occurrences in Sediment of Lagos Lagoon using Environmental Indices****Abstract**

Occurrence of five heavy metals (Mn, Pb, Cr, As and Zn) and their seasonal distribution between 2012 to 2013 in the sediment of Lagos Lagoon using index analysis tools: *Contamination Factor (C_f)*, *Degree of Contamination (C_d)* and *Pollution load Index (PLI)* was studied. Six sampling stations were established based on the activities within the sampling stations which include industrial effluents, domestic waste and sewage discharges. Stations 4 and 5 have high concentration of Pb above the permissible level during the wet season while the Pb concentrations in the remaining four stations were within the permissible limit of 40mg/kg of USEPA and DPR. Mn level in all the sediments sampling station analyzed during the two season exceeded USEPA and DPR of 40 mg/kg. Cr and Zn are within the permissible limit in all the stations during the wet and dry season hence, reflecting their natural background levels in the sediment. The concentration of As detected in all the sediment analyzed exceeded USEPA of 9.8 mg/kg while station 5 had the highest concentration of 72.86 mg/kg in the dry and 89.65 mg/kg in wet seasons respectively. Comparison between the results obtained from the sediment analyzed during dry and wet seasons for metal concentration, indicated that all the contaminants are significant at ($p < 0.05$) level during the dry and wet seasons except for Pb and Cr that are significant at ($P < 0.01$) level while As and Mn are not significantly correlated between the two seasons. The metals concentrations are arranged in increasing order in this area during the dry and wet seasons were: $Mn > As > Pb > Zn > Cr$. The mean concentration of metals in sediment for the two seasons is significantly correlated at (α level=0.05 and 0.01) which implied that the metals increase within the season in the matrix of sediment. Assessments of heavy metal contamination in sediment require knowledge of pre anthropogenic metal concentrations to act as a reference against which measured values can be compared. Sediment contamination by heavy metal based on the above index showed that the study area have very low degree of metal contamination.

Keywords: Heavy metal, sediment, USEPA, DPR, Contamination Factor , Degree of Contamination, Pollution load Index.

Introduction

The Lagos Lagoon is vastly impacted by numerous wastes which posed a stern threat to the community at large depends on it for the source of income especially for the masses living along the water's edge. Wastes of anthropogenic origin often contaminate the Lagoon and creek in Lagos [1]. Majority of the debris are largely plastics, nylon bags, empty cans of food and drinks, glass bottles, used needles and syringes and used car tyres *etc.* Marine sediment is frequently used as sensitive foundation of indicator for both spatial and temporal trends in monitoring of the aquatic ecosystem. The contamination of sediment is one of the main environmental problems that are often mandatory for frequent evaluation in order to maintain the aquatic biota and 70% of Lagos Lagoon is a reservoir for industrial, surface runoffs, and domestic discharge as well as input from several rivers [2]. Heavy metals such as cadmium, mercury, lead, copper, Arsenic, Copper, Iron and zinc are consider the most important pollutants of aquatic ecosystems owing to their persistence, toxicity and ability to be integrated into food chains [3]. Globally, metals generally persist and bioaccumulate in the living organisms in the natural ecosystem when they exceed their permissible level [4]. Quantity of wastes discharge to the aquatic environment often lead to bioaccumulation of heavy metal in the aquatic ecosystem, consequently leading to decline of the physical, chemical and biological change of the aquatic environment [5].

The amount of heavy metals in water are majorly in minute quantities whereas excessive concentration can be identified in the sediment sample thereby, affecting aquatic biota and causing risk to consumers of aquatic seafood [6]. The assessment of sediments developed recently into a foremost issue of significance that can revealed the existing quality of the environment as well as provide vital information on the anthropogenic activities within the aquatic biota. Sediment can be defined as loose


47 sand, clay, silt and other soil particles that settle at the bottom of body of water [7]. Corrosion of bedrock and soils leads to
 48 accumulation of sediment of current natural and anthropogenic activities as well as their component [8]. The major contributing
 49 factor to metal contamination in the sediment are through numerous pathway such as industrial disposal of liquid effluents,
 50 surface runoff, domestic discharge and leachates carrying chemicals originating from numerous urban and agricultural activities
 51 as well as atmospheric deposition [9]. Consequently, data from sediment can provide series information on the impact of
 52 anthropogenic on the broader environment since sediment is the final sink for metals accumulation. Constitute of sediment
 53 progression present the most efficient natural archives of current environmental changes within the aquatic biota. Universally,
 54 sediments are significant medium of assessment of anthropogenic inputs in the aquatic ecosystem. Therefore, metal
 55 concentration and distribution in sediment can best provide current information about spatial extent as well as magnitude of
 56 anthropogenic induced chemical change of the environment thereby leading to useful indicators of contaminant in aquatic
 57 ecosystems

58 The objective of this study was to evaluate the occurrence of heavy metal in the sediment of Lagos Lagoon environment and the
 59 ecological deduction that can be made from the results using contamination factor, degree of contamination and pollution load
 60 index tools.

61

62 Materials and Methods

63 **Study Area:** Lagos lagoon is one of the largest Lagoon systems of the Gulf of Guinea. The Lagos Lagoon receives a number
 64 of important large rivers including: Oshun, Yewa, Ogun and Ona rivers. This Lagoon empties directly into the Atlantic Ocean
 65 at Lagos Harbor; it is with between longitude 30 10' and 30 4' SE and latitude 60 5' and 60 36' N[Fig 1].



66

67 Fig 1: Map of the study area showing the sampling locations

68

69 Table 1: Description of sampling station

Station	Longitude	Latitude	Sampling area description	Location	Waste Type
1	3° 23' 5.602"	6° 28' 02.7"	Based on the visitation of itinerant tankers to the jetty to discharge raw faeces.	Iddo	Sewage
2	3° 22' 37.8"	6° 27' 54.7"	Wastes are majorly from sea food, cement bag washing, as well as influx of sewage discharged from Iddo all find their way into Ijora.	Ijora	Industrial
3	3° 23' 48.6"	6° 26' 17.9"	Receives wastes from Ikoyi, Lagos Island and Victoria Island.	Five cowries	Domestics and industrial.
4	3° 22' 42.4"	6° 26' 48.5"	Seaport activities with various effluents from the sea well as ship discharge from the passing vessel.	Apapa Port	Industrial
5	3° 21' 37.44"	6° 25' 46.8"	It houses most of the oil port and tank farms in the Lagos Lagoon as well as effluents from the Port novo creek, Badagry Creek and the commodore channel.	Tincan Creek	Industrial and domestic
6	3° 24' 26.0"	6° 25' 14.1"	Characterized by different types of wastes from ship discharged as well influx of others domestics' wastes from Victoria Island.	Commodore Channel	Industrial and domestic

Sampling Collection: Sediments were collected with the aid of Van Veen grab at each station and stored immediately in polythene bag. Sediments collected were stored at 4°C in an ice-box and transported to the laboratory. The entire samples were separately air dried in a laboratory. After drying the sediment, it was homogenized and sieved to remove big particulates. Homogenized sediment samples were then digested as follows: 1g of the powdered sediment samples were weighed into a 100 ml beaker. 25ml of freshly prepared mixture of aqua regia (HNO₃ /HCl) ratio 1:3 were added to each sample and covered with a wash glass. It was allowed to stand for 30 minutes during which initial reaction subsided [10]. Digestion was carried out on hot plate whose temperature was allowed to rise gradually until it reached a maximum temperature of 160°C in a fume cupboard. Heating was continued for about 2 hours, reducing the volume in the beaker to about 2 - 5ml. The beaker and its contents were allowed to cool and the content was transferred with whatman filtration into a 50ml volumetric flask and made up to mark with distilled water. The digested samples were then analyzed for Pb, Zn, Mn and Cr using a Flame Atomic Absorption Spectrophotometer model Varian SpectAA 400. While As was detected with the aid of Varian AA240 hydride generation atomic absorption spectrophotometer (AAS).

Statistical Analysis: The Metals were evaluated using the one-way analysis of variance (ANOVA) and means were compared using Duncan multiple range test. P values < 0.05 were considered significant.

1. Contamination factor: The extent of contamination of sediment by a metal is often expressed mathematically in terms of a contamination factor calculated by $CF = \frac{\text{Metal content in the sediment}}{\text{Background level of metal}}$ Equation [1]

Where CF < 1 refers to low contamination, $1 \geq CF \geq 3$ means moderate contamination, $3 \geq CF \geq 6$ indicates considerable contamination, and CF > 6 indicates very high contamination.

2. Degree of contamination (Cd) : This is used to evaluate the extreme values of monitored elements in soil sediment samples. It is expressed mathematically, it is expressed as given in Equation 3. C_iF is the contamination factor for the i-th element.

$C_d = \sum C_iF$ Equation[2]

Where (**Cd**) <6 refer to Low degree of contamination, $6 \geq Cd \geq 12$ mean moderate degree of contamination, $12 \geq Cd \geq 24$ refer to high degree of contamination and $Cd > 24$ refer to very high degree of contamination

3. **Pollution Load Index (PLI)**: this is used to evaluate specific sampling site, according to the method proposed by Tomilson *et al.* [11] and it is expressed as:

$$PLI = [CF1 \times CF2 \times \dots \times CFn]^{1/n} \dots \dots \text{Equation [3]}$$

Where, n is the number of metals in the present study and CF is the contamination factor.

Results

1. Evaluation of Heavy Metal Concentrations according to Mean and one way-ANOVA using Duncan multiple ranges test.

The mean concentration of each metal examined in the sediment from each sampling station is shown in Table 2 and fig 2-6. Mean metal concentrations ranged within the following intervals during the dry season: Mn: 124.42 -60.3mg/kg , As: 70.46-11.3mg/kg; Pb: 28.6-9.73mg/kg; Cr: 14.19-2.43mg/kg and Zn:29.8-6.4mg/kg, and their mean concentrations during the wet season were: Mn: 219.13-139.07mg/kg; As:90.26-22.5mg/kg; Pb: 41.89-21.17mg/kg; Cr: 19.7-4.92mg/kg and Zn: 35.21-6.5mg/kg. The metals are arranged in terms of their occurrence in the sampling station during the dry season as follows: Mn>As>Zn>Pb>Cr. While the mean metal concentration arrangements for wet season as follows: Mn>As>Pb>Zn>Cr. The high levels of heavy metals obtained in the sediments of the Lagos Lagoon could probably be associated with the following activities such as sewage, cement bags washing domestics and industrials waste that took place within and around the Lagoon. The highest mean metal concentration values for Mn and Zn were registered at Station 6, while Station 4 and 5, respectively, recorded the highest mean metal concentrations for Pb, As and Cr in the dry season; these was expected due to large number of fishing vessel and fuel dispensing points concentrated at these sample stations. In the wet season the highest mean metal concentration value for Mn and As were detected in the station 5 while the highest mean metal concentration recorded for Pb, Zn and Cr in Station 4 and 6. The results signify that mean concentrations of all heavy metals in the sediments sampled analyzed during dry season was higher than concentrations of all heavy metals during wet season were probably due to dilution by rainwater which influences concentration and heavy metal mobility. ANOVA was used to establish whether cluster of variables have the same means on data that are continuous distributed and with homogeneous variance using Duncan multiple range. Furthermore, it was used to evaluate the relationship between heavy metal concentrations and their elemental interaction between sections of the stream. There was significant difference ($p < 0.05$) between the level of heavy metals during different seasons except in the Manganese Mn (Table 3).

Table 2: Mean Concentration and Standard deviation of Heavy Metals (mg/kg) from Each Sampling Station and their standard geochemical background value.

Station	Mean and standard deviation of Metal concentration in mg/kg in Dry season					Mean and standard deviation of Metal concentration in mg/kg in Wet				
	Mn	As	Pb	Cr	Zn	Mn	As	Pb	Cr	Zn
Ijora	82.19±0.92	61.52±0.28	13.5±1.56	2.43±0.08	13±1.47	176.73±1.37	55.46±1.37	21.17±1.46	5.18±1.56	10.47±1.88
Idoo	104.05±1.24	11.39±1.98	17.09±1.19	2.93±1.63	14.19±1.23	139.07±2.58	62.48±1.49	24.18±1.97	4.92±0.90	18.81±1.83
Five cowries creek	109.66±0.93	25.78±1.67	9.73±1.22	2.48±1.05	3.57±1.73	159.35±0.91	25.99±1.60	25.9±2.36	6.1±0.91	13.78±1.74
Apapa port	60.3±1.73	15.49±1.63	28.61±1.78	13.96±0.69	17.39±1.49	108.26±3.42	22.5±1.59	40.64±2.01	19.71±0.71	23.04±1.80
Tincan creek	114.01±1.77	15.49±1.84	22.96±1.67	14.74±1.07	6.4±1.82	219.13±2.21	90.26±1.54	31.7±2.34	17.04±1.24	6.5±1.02
Commodore channel	124.42±3.52	15.49±1.65	25.34±1.58	11.8±1.05	29.8±1.29	179.37±3.76	29.74±1.46	41.89±1.47	17.6±1.32	35.21±1.63
World surface rock	720	13	20	97	129	720	13	20	97	129

value									
-------	--	--	--	--	--	--	--	--	--

128

129 *Table 3: shows the mean of heavy metal and Duncan's multiple range tests using One way-ANOVA*

Station	Mean Metal concentration in mg/kg in Dry season					Mean Metal concentration in mg/kg in Wet				
	Mn	As	Pb	Cr	Zn	Mn	As	Pb	Cr	Zn
Ijora	82.19 ¹	61.52 ^{2*}	13.50 ^{1 2*}	2.43 ^{1*}	13.00 ^{2&3*}	176.73 ^{2 & 3}	55.46 ^{1&2*}	21.17 ^{1*}	5.18 ^{1*}	10.47 ^{1*}
Idoo	104.05 ¹	11.39 ^{1*}	17.09 ^{1, 2 & 3*}	2.93 ^{1*}	14.19 ^{3*}	139.07 ^{1&2}	62.48 ^{1&2*}	24.18 ^{1&2*}	4.92 ^{1*}	18.81 ^{2&3v}
Five cowries creek	109.66 ¹	25.78 ^{1*}	9.73 ^{1*}	2.48 ^{1*}	3.57 ^{1*}	159.35 ^{1, 2 & 3}	25.99 ^{1*}	25.90 ^{1&2*}	6.10 ^{1*}	13.78 ^{1&2*}
Apapa port	60.30 ¹	15.49 ^{2*}	28.61 ^{4*}	13.96 ^{2*}	17.39 ^{3*}	108.26 ¹	22.50 ^{1*}	40.64 ^{2*}	19.71 ^{3*}	23.04 ^{3*}
Tincan creek	114.01 ¹	70.46 ^{2*}	22.96 ^{2, 3 & 4*}	14.74 ^{1 & 2*}	6.40 ^{1&2*}	219.13 ³	90.26 ^{2*}	31.70 ^{3*}	17.04 ^{2*}	6.50 ^{1*}
Commodore channel	124.42 ¹	16.34 ¹	25.34 ^{3 & 4*}	11.80 ^{2*}	29.80 ^{4*}	179.37 ^{2&3}	29.74 ^{1*}	41.89 ^{3*}	17.60 ^{2*}	35.21 ^{4v}

Values are the means of metals per station. Means with the same letter are not significantly different from each other. ¹⁻⁴Means in a row without a common superscript letter differ ($P < 0.05$), as analyzed by one-way ANOVA.

130
131
132
133




Fig 2: Manganese concentration in the sediment




Fig 3: Arsenic concentration in the sediment

134
135
136




Fig 4: Lead concentration in the sediment




Fig 5: Chromium concentration in the sediment




Fig 6: Zinc concentration in the sediment

2. Assessment According to the Correlation between the Elements

Pearson's correlation coefficient matrix among the selected heavy metals is presented in Tables 4. Significant correlations between the contaminants during the dry season was between Pb and Cr ($r=0.865$), Pb and Zn ($r=0.692$), Cr and Pb ($r=0.865$), which probably indicate the related source input during the dry season while for wet season, the significant correlation between the metal concentration is between Mn and Pb ($r=0.731$), Pb and Mn ($r=0.731$), Pb and Cr ($r=0.809$), Pb and Zn ($r=0.802$), Cr and Pb ($r=0.809$), Zn and Pb ($r=0.802$) may possibly signify the similar source contribution. Lagos Lagoon is surrounded by very old buildings roofed by means of zinc sheets with lead paints. Hence, the high correlation between Zn and Pb in sediment could be attributable to roofing sheets and paints from the homes located along the shore of the Lagos Lagoon. The availability of As at a location has been accredited to the unsuitable disposal of electronic waste, industrial effluents, fertilizers in agricultural surface runoff and atmospheric depositions [12]. By comparing the result obtain from metal concentrations during the Dry and Wet season respectively, it could be shown that all the metals are significant at $p > 0.05$ level during the dry and

154 wet seasons except for Pb and Cr that are significant at 0.01 level while As is not significantly correlated between the two
155 season.

156
157
158
159

Table 4: Pearson correlation matrix for heavy metal concentration in surface sediments from The Lagos Lagoon during the Dry and Wet Season

	Station	Mn.Dry	As.Dry	Pb.Dry	Cr.Dry	Zn.Dry	Mn.Wet	As.Wet	Pb. .Wet	Cr.Wet	Zn. .Wet
Station	1										
Mn.Dry	0.245	1									
As.Dry	-0.113	0.291	1								
Pb.Dry	.565*	0.316	0.027	1							
Cr.Dry	.747**	0.199	0.133	.841**	1						
Zn.Dry	0.399	0.233	-0.321	.631**	0.388	1					
Mn.Wet	0.225	.699**	.673**	0.12	0.226	-0.094	1				
As.Wet	-0.079	0.234	.622**	-0.059	0.071	-0.25	0.408	1			
Pb.Wet	.786**	0.215	-0.219	.854**	.805**	.665**	-0.043	-0.307	1		
Cr.Wet	.843**	-0.024	-0.044	.764**	.891**	0.431	0.031	-0.143	.845**	1	
Zn.Wet	.469*	0.062	-.605**	0.44	0.216	.829**	-0.379	-0.44	.683**	0.367	1

*. Correlation is significant at the 0.05 level (2-tailed).

**. Correlation is significant at the 0.01 level (2-tailed).

160
161
162
163
164
165
166
167
168

3. Assessment according to USEPA sediment quality guideline

Evaluations of the chemical contaminations in the sediment were estimated by comparison with the sediment quality guideline proposed by United States Environmental Protection Agency [13] and Department of Petroleum Resource. These criteria are shown in Table 5. The Mean concentration of metals in the present study shows that all the sites according to the USEPA scale indicated a heavy pollution level for Mn and As while for Pb in station 4 is moderately polluted during the wet season. For Cr and Zn are within the permissible limit for the two seasons in all the stations, hence, not polluted.

Table 5: Comparison with USEPA (2003) Guidelines for Sediment

Metal	USEPA Pollution Level (mg/kg)			Present study(mg/kg)	
	Not polluted	Moderately polluted	Heavily polluted	Dry	Wet
Mn	<46	>46	>60	55.17 – 137.56	105.25 - 194.76
Pb	<40	40-60	>60	9.62 - 27.59	15.65 - 89.65
Cr	<25	25-75	>75	2.09 - 12.07	3.89 - 19.65
As	<9.8	>9.8	>10.94	10.94 - 72.86	44.18 - 18.98
Zn	<90	90-200	>200	3.88 - 29.06	6.05 - 34.56

169
170
171
172
173
174
175
176
177
178
179
180
181
182

4. Assessment According to the Contamination Factor (C_F) and Degree of Contamination (C_d) in the sediment of Lagos Lagoon:

The values of the metals studied from all five sampling stations are shown in Table 6. In this present study, the highest C_F value was obtained for As, which registered a very high contamination factor with low contamination factors being registered for Cr and Zn. Based on the mean C_F values, the sediments sampled analyzed in during the two season may probably be considered to be enriched by the metals examined in the following order: As > Mn > Pb > Cr > Zn. C_F values for metals analyzed in the sediment shows higher (>1) values due to the influence of anthropogenic inputs except in Zn during the dry season. The cumulative effect of each individual mean metals presents in the sediment sample at each of the stations signify that station 5 has the highest degree of contamination, while Station 2 record the lowest contamination during the dry season while station 5 also registered the highest degree of contamination and station 4 has the lowest degree of contamination during the wet season.

Table 6: Contamination Factor (C_F) and degree of contamination (C_d) Values for Sediments of Lagos Lagoon

station	Contamination Factor (C_F) during Dry season					Degree of Contamination (C_d)	Contamination Factor (C_F) during Wet season					Degree of Contamination (C_d)
	Mn	As	Pb	Cr	Zn		Mn	As	Pb	Cr	Zn	
1	1.79	6.3	0.3	0.1	0.14	8.65(2 nd)	3.84	5.66	0.53	0.21	0.12	10.36(3 rd)
2	2.26	1.2	0.4	0.12	0.16	4.13(6 th)	3.02	6.38	0.6	0.2	0.21	10.41(2 nd)
3	2.38	2.6	0.2	0.1	0.04	5.39(4 th)	3.46	2.65	0.65	0.24	0.15	7.15(5 th)
4	1.31	1.6	0.7	0.56	0.19	4.36(5 th)	2.35	2.3	1.02	0.79	0.26	6.72(6 th)
5	2.48	7.2	0.6	0.59	0.07	10.9(1 st)	4.76	9.21	0.79	0.68	0.07	15.51(1 st)

	6	2.7	1.7	0.6	0.47	0.33	5.8(3 rd)	3.9	3.03	1.05	0.7	0.39	9.07(4 th)
Total per metal C _F		12.92	20.51	2.93	1.94	0.93	39.23	21.33	29.23	4.64	2.82	1.2	59.22

5. **Assessment According to the Pollution Load Index in the sediment of Lagos Lagoon:** The values of Pollution Load Index (fig. 7) were found to be generally higher (<1) in all the studied stations except in Zn which is 0.93 during the dry season. The variation in index results obtained was probably due to the disparity in sensitivity of these indexes towards the sediment contamination [14]. These confirmed that Lagos Lagoon is currently undergoing possible environmental contamination especially with hazardous metals Pb and As.




Fig 7: Pollution load index of heavy metals in the sediment of Lagos Lagoon

Discussion

Heavy metals contamination in sediment is of major concern particularly in many modern countries due to their toxicity, persistence and bioaccumulative nature [15]. It has been reported that the high level of metal concentration observed in all the stations during the wet season may be due to influx of waste which influences their concentration and metals mobility which depends directly on the environment and some metals properties such as pH, organic matter concentration, silt and clay concentration, salinity, dissolved oxygen concentration [16]. However, it has been reported that mobility of heavy metals depends not only on the total concentration in the soil and sediment but also on the sediment properties, metal properties and environmental factors. Occurrence and Spatial distribution patterns of Mn, As, Cr, Zn and Pb in Lagos Lagoon sediments were illustrated in Fig. 2- 6. The mean concentrations of Pb in the sediment are within 40mg/kg in USEPA Sediment Quality Guidelines except in station 4 and 5 that exceed the maximum permissible level. The results indicate that urbanization has contributed to the rise of Pb content in the sediment. Domestic and industrial effluents, municipal runoffs and atmospheric deposition may be the major sources. The contribution of Pb from the use of leaded petrol in ship and speed boat operator is possibly significant. Sadip, *et al.*, [17] reported that low concentrations of Pb still might pose a threat to life in a marine environment in comparison with other heavy metals. As mean concentration exceeds the permissible level of 9.8 mg/kg in all the sampling stations during the dry and wet season according to USEPA and DPR sediment quality guideline and have potential damage to the ecological communities according to the environmental protection agency of USA .The relatively low concentration detected in chromium and Zinc in the sediment reflects their natural background levels in the sediment. Hall *et al.*,[18] reported that Zn is a micronutrient for aquatic life in all natural water and sediments and even at low concentration , Zn can become toxic to aquatic organisms at higher concentration level than the threshold required contents.

Mn in sediment during the wet season was higher than that of the dry season and this may be due to rainfall and run off. This agrees with the findings of Lokeshwary and Chandrappa [19]. Mn has been found to occur at high concentrations in Lagos Lagoon [20]. Previous studies also have indicated elevated levels of heavy metals in aquatic systems receiving effluents from urban areas and the disposal of untreated sewage has been reported to contribute significantly to the level of heavy metals in the aquatic environment [21]. The finding in this study shows that metals concentration is higher in the wet season than in the dry season in all the stations which are properly due to influx and other anthropogenic activities within the sampling station in the Lagoon. The Chromium and Zinc detected within all the sampling station are within the permissible limit of USEPA for sediment quality guideline of 25 mg/kg and 40 mg/kg respectively while for Mn and As exceeded their permissible level in all the sediment sampled analyzed expect for Lead that exceed the permissible level in station 4. Generally, the pollution load index value significant increased as anthropogenic activity increased as well as dispersion of metal content with increasing distance from point of discharged. The relative low value of pollution load index in Zn indicates no significant input from anthropogenic sources. Table 6 indicates the contamination factor of each element and the degree of contamination in each of the sampling station. According to Hakanson classification [22], the contamination factor for all the sampling station fall within the high contamination factor and the values of contamination degree indicates high degree of contamination. The degree of contamination for the individual metal recline within the range of 4.13-10.9 during the dry season while degree of contamination during the wet season are within the range of 6.72-15.51 which are more than 1.5 indicating very high C_d . The degree of contamination data signifies anthropogenic impact in all the sampling station. Pollution load index were found to increase in sediments upstream along Lagos Lagoon (Fig. 7). These is attributed to industrial emissions, domestic and industrial effluents such as sewage discharge, cement bags washing and influx from the nearby river loaded with various waste are subsequently deposited into the Lagos Lagoon sediments.

234
235

236 CONCLUSION

The results obtained in this studying indicate the deleterious effect of anthropogenic activities on the Lagos Lagoon which are apparent in the high level of Mn, As and Pb at the sampling station studied. Contaminations in the sediments were further evaluated by applying contamination factor, degree of contamination and pollution load index. The results of contamination factor, degree of contamination illustrated that the sampling station falls within high to very high contamination, Calculation of Pollution load Index the study area is regarded as polluted. Based on the studied areas, the framework for mandatory action should be initiated for regular assessment of Lagos Lagoon and indexes are very useful tools for identification of sediment contamination.

244
245
246

245 References

- 247 1. Ajao, E. A., Okoye, B. C. O., and Adekanmbi, E. O., (1996). Environmental Pollution in the Nigerian Coastal Waters:
248 A Case Study of the Lagos lagoon. *Water Quality Monitoring and Environmental Status in Nigeria. FEPA Monograph*, 6:101-
249 112.
- 250 2. Olatunji, A. S., Abimbola A. F., (2010). Geochemical Evaluation of the Lagos Lagoon Sediments and Water, *World*
251 *Appl. Sci. J.*, , 9(2): 178-193.

252
253

- 254 3. Turner, A., (1996). Trace metals portioning in estuaries importance of salinity and particulate concentration. *Marine*
255 *Chemistry*, 54, 27 – 39.
- 256 4. Chakraborty, R., Zaman, S., Mukhopadhyay, N., Banerjee, K., and Mitra, A., (2009). Seasonal variation of Zn, Cu and
257 Pb in the estuarine stretch of west Bengal. *Indian Journal of Marine Science*, 38 (1), 104 –109.
- 258
- 259 5. Ridgway, J., and Shimmield, G., (2002). Estuaries as repositories of historical contamination and their impact on shelf
260 areas. *Estuarine, Coastal and Shelf Science*, 55, 903–928.
- 261
- 262 6. Mitra, A., Trivedi, S., Gupta, A., (1996). Distribution of trace metals in the sediments from Hooghly estuary, India.
263 *Pollution Research*, 15, 137 – 141.
- 264
- 265 7. Forstner, U. and Wittmann, G.T.W., (1983). *Metal Pollution in the Aquatic Environment*. Spring Verlag Berlin
266 Heidelberg, New York. 1983, 486Pp.
- 267
- 268 8. Davies, O. A. and Abowei, J. F. N. (2009). Sediment quality of lower reaches of OkpokaCreek, Niger Delta, Nigeria.
269 *European Journal of Scientific Research*. ,26 (3): 437 – 442.
- 270
- 271
- 272 9. Adefemi, S. O., S. S. Asaolu and O. Olaofa, (2008). Determination of heavy metals in *Tilapia mossambicus* fish,
273 associated water and sediment from Ureje dam in South-Western Nigeria. *Res. J. Environ. Sci.*, 2: 151-155
- 274
- 275 10. APHA. (2005). *Standard Methods for the examination of water and wastewater*, American Public Health Association,
276 Washington, DC. 21st Edition. 1368 Pp.
- 277
- 278 11. Tomilson, D.C.; Wilson, J.G.; Harris C.R. (1980). Jeffrey, D.W. Problems in assessment of heavy metals in estuaries
279 and the formation of pollution index. *Helgol Meeresunters.*, 33,566-575.
- 280
- 281
- 282 12. Talbot, V. and Chegwiddden, A., (1983). Heavy metals in the sediment of Cockbun Sound, Western Australia and its
283 surrounding area. *Environ. Pollution.*, 5: 187-193.
- 284
- 285 13. USEPA (2003). The incidence and severity of sediment contamination in surface waters of the United States: Volume
286 1: National Sediment Quality Survey.. Washington, District of Columbia. United States Environmental Protection Agency,
287 EPA/823/R-97/006
- 288
- 289 14. Department of Petroleum Resources, (2002), Environmental guidelines and Standards for the petroleum industries in
290 Nigeria Department of Petroleum Resources, Ministry of Petroleum and Mineral Resources, Abuja, Nigeria.
- 291
- 292 15. Mohammed, N. K., (2000). The Optimization of Energy Dispersive X-ray Fluorescent (EDXRF) Spectrometer for the
293 study of Heavy Metal Pollution. M.Sc. Thesis, University of Dar Es Salaam, Tanzania.
- 294
- 295 16. Yaron B, Calvet R, Prost R.(1996) Soil pollution: processes and dynamics [Internet]. Berlin: Springer; 315
- 296
- 297 17. Martin, J. M, Maybeck M. (1979). Elemental mass of material balance carried by major world rivers. *Marine*
298 *Chem.*7:173-206.
- 299
- 300
- 301 18. Sadip, R., Husain, T., Bose, N., Veitch, B., (2003).Distribution of Heavy Metals in Sediment Pore Water due to
302 offshore discharges: an ecological risk assessment. *Environmental Modeling and Software.*216 18, 451–461.

303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318

19. Hall, L.W.J., Scott, M.C., Killen, W.D.,(1997). A screening level probabilistic ecological risk assessment of copper and cadmium in the Chesapeake Bay Watershed. Chesapeake Bay Program.US EPA, Annapoils, MD.
20. Lokeshwary, H. and Chandrappa, G. T.,(2006) Impact of Heavy Metal Contamination of Bellandur Lake on Soil and Cultivated Vegetation. *Current science.*, 91(5): 622-627.
21. Nabulo, G.; Oryem-origa, H.; Nasinyama, G.; Cole, D.,(2008) Assessment of Zn, Cu, Pb and Ni contamination in wetland soils and plants in the Lake Victoria basin. *Int. J. Environ. Sci. Tech.*,. 5 (1), 65-74 .
22. Nyangababo, J. T.; Henry, E.; Omutange, E., (2005).Lead, cadmium, copper, manganese and zinc in wetland waters of Victoria Lake Basin, East Africa. *Bull. Environ. Contam. Toxicol.*,. 74 (5), 1003-1010.
23. Hakanson, L., (1980).Ecological Risk Index for Aquatic Pollution Control. A Sedimentological Approach. *Water Research.* 14: 975-1001.