

1 **Fermi problem and superluminal signals in quantum electrodynamics**

2 **Abstract**

3 Using as an example the Fermi problem dealing with nonstationary excitation trans-
4 formation from one atom to another the reason of superluminal signals appearance in quan-
5 tum electrodynamics is clearing. It is shown that the calculation using the conventional me-
6 thods in Heisenberg and Schrödinger representations in nonstationary problems lead to dif-
7 ferent results. The Schrödinger representation predicts the existents of specified quantum su-
8 perluminal signals. In Heisenberg representation the superluminal signals are absent. The rea-
9 son of nonidentityof representations is close connected with using of the adiabatic hypothesis.

10 PACS numbers: 42.50.-p

11 **1. Introduction**

12 In 1932 year E. Fermi [1] by developing the Dirac theory [2] of quantum transpositions had
13 considered the problem dealing with nonstationary transformation radiation between excited
14 atom and another atom being in its ground state. He calculated the probability of suchprocess
15 as a square of corresponded matrix element. It was shown that the radiation transformation
16 has the retarded character and is described by character construction $t = R / c$. Here t is the
17 time of excitation transformation, R is the distance between atoms, c is the light velocity in
18 vacuum. The result was repeated in many following theoretic papers [3,4] .

19 The detailed analysis of Fermi calculations performed in the paper [5] had shown that the
20 retarded character of signal defined by formula $t = R / c$ is only the approximation connected
21 with using the pole approximation. More punctual calculations show the appearance of a
22 small superluminal forerunner placed before the classical electromagnetic wave front at a dis-
23 tance of the order of one wave length. The author supposes that such a fact does not have the
24 physical sense. In his paper [5] he tried to proof this fact in general form using the Heisen-
25 berg representation.

26 In paper [6] the Fermi result was analyzed again. The appearance of the superluminal forerun-
27 ner forces the authors to clean the reason of its appearance and revise the Dirac theory [2] of
28 quantum transpositions. In the paper [6] one postulates the incorrectness of representation of
29 quantum transpositions probability as a square of consequence matric elements. One proposes
30 to evaluate the observed values as quantum average values of consequence quantum operators.

31 These average values have to be calculated in Heisenberg representation. Such way leads to the
 32 exact realization of the expression $t = R/c$ and like the method proposed in [5]. The
 33 Schrödinger representation was not investigated.

34 In a paper [6] as in a paper [5] using the Heisenberg representation the authors came
 35 to the conclusion of impossibility of the appearance in quantum electrodynamics the super-
 36 luminal signal.

37 Last years the interest for the optical superluminal signals has risen supplementary.
 38 Such signals were discovered in many experimental works [7-14]. The necessity of their theo-
 39 retical description has appeared. All attempts of theory constructions in present days (the fluc-
 40 tuations excluded) deal extremely with using the classical representation of the internal struc-
 41 ture of electromagnetic field [15-19]. The exception represents the paper [20]. In this work
 42 using the interaction representation an evident nonequality $\langle \hat{E}^2 \rangle \geq \langle \hat{E} \rangle^2$, \hat{E} being the strength
 43 operator of electromagnetic field, one shows the appearance in electrodynamics of excited
 44 media the superluminal signals. Such signals do not have the classical analogs. For the appear-
 45 ance of such signals the inversion population of atom states in media is not necessary. Such
 46 superluminal signals were experimentally observed and evidently are in a good coincidence
 47 with experimental data [13]. The reason and their appearance conditions in connection with
 48 experiments mentioned above are very interesting. In present work such questions are solved
 49 using the Fermi-problem as example. In such a way one shows that the quantum radiation trans-
 50 fer in quantum electrodynamics at the finite times in Heisenberg and Schrödinger representa-
 51 tions are described in different way. Other words these representations are non-identical. Such
 52 result possesses not only the methodic character. The fact is that the superluminal signals ap-
 53 pear only in Schrödinger representation. In Heisenberg representation they are absent. This fact
 54 permits to understand the result differences in the calculations using the different methods.
 55 Namely this fact opens the possibility for prediction the analogous results in other situations.

56 We doubt not in the results of calculations in papers [5] and [6] but we doubt in the fi-
 57 nite conclusions in these works. In these works the conclusions about the absent of superlu-
 58 minal signals in quantum electrodynamics follows from Heisenberg representation. But in
 59 these works the analysis using Schrödinger representation is absent. We revise in the follow-
 60 ing the solutions of Fermi-problem by using both representations. We shall show that non-

identity of Schrödinger and Heisenberg representations in nonstationary problems is naturally and connected closely with using in quantum electrodynamics the adiabatic hypothesis.

2. The state of the problem

Let us suppose that the test atom (1) being in its ground state is placed at the point \mathbf{R}_1 and is attacked by the radiation of excited atom (2) placed in the point \mathbf{R}_2 . The excited atom begins interact with electromagnetic field at a moment of time t_0 . Each atom possesses only one electron. We neglect the spin variables. One supposes the atoms are placed in wave zone at a large distance between them that permits to neglect in the exchange effect and in the longitudinal electromagnetic field. Suppose that each atom possesses only two energetic levels. But these levels may have energetic sublevels. The excited and ground states of primary excited atom (2) are describes consequently by indexes j_{ex} and j_g . The energetic states of non-excited atom (1) are described by indexes i_{ex} and i_g . The Hamiltonian of the problem in Schrödinger representation and quasi-resonant approximation is written in the following form

$$\hat{H} = \hat{H}^0 + \hat{H}', \quad \hat{H}^0 = \int \hat{\psi}_1^+(\mathbf{r}_1) \hat{H}_1 \hat{\psi}_1(\mathbf{r}_1) d\mathbf{r}_1 + \int \hat{\psi}_2^+(\mathbf{r}_2) \hat{H}_2 \hat{\psi}_2(\mathbf{r}_2) d\mathbf{r}_2 + \hat{H}_{ph}$$

$$\hat{H}' = -\frac{e}{mc} \int \hat{\psi}_1^+(\mathbf{r}_1) \hat{p}_{r_1}^{v_1} \hat{A}^{v_1}(\mathbf{r}_1) \hat{\psi}_1(\mathbf{r}_1) d\mathbf{r}_1 - \frac{e}{mc} \int \hat{\psi}_2^+(\mathbf{r}_2) \hat{p}_{r_2}^{v_2} \hat{A}^{v_2}(\mathbf{r}_2) \hat{\psi}_2(\mathbf{r}_2) d\mathbf{r}_2 \theta(t - t_0), \quad (1)$$

$\theta(t - t_0)$ being the Heaviside step function that fixed the moment of time appearance of the interaction of radiated atom with electromagnetic field. Over the repeated indexes one supposes the summation,

$$\hat{\psi}_1(\mathbf{r}_1) = \sum_i \psi_i(\mathbf{r}_1 - \mathbf{R}_1) \hat{b}_i, \quad \hat{\psi}_2(\mathbf{r}_2) = \sum_j \psi_j(\mathbf{r}_2 - \mathbf{R}_2) \hat{b}_j, \quad \hat{H}_{ph} = \sum_{\mathbf{k}\lambda} \hbar c k \left(\hat{\alpha}_{\mathbf{k}\lambda}^+ \hat{\alpha}_{\mathbf{k}\lambda} + \frac{1}{2} \right),$$

$$\hat{A}^v(\mathbf{r}) = \sum_{\mathbf{k}\lambda} \sqrt{\frac{\hbar c}{2kV}} e_{\mathbf{k}\lambda}^v \left(\hat{\alpha}_{\mathbf{k}\lambda} e^{i\mathbf{k}\mathbf{r}} + \hat{\alpha}_{\mathbf{k}\lambda}^+ e^{-i\mathbf{k}\mathbf{r}} \right).$$

The wave functions ψ_i and ψ_j denote the behavior of electrons in atoms (1) and (2), \hat{b}_i^+ and \hat{b}_j^+ denote the electron creations operators at the same states. By $\hat{\alpha}_{\mathbf{k}\lambda}$ and $\hat{\alpha}_{\mathbf{k}\lambda}^+$ the annihilation and creation photon operators in states (\mathbf{k}, λ) are denoted. Here \mathbf{k} is the photon wave vector, λ is the index of its polarization. The photons have only the transversal polarization λ

85 =(1,2). The rationalized Gauss unite system is used. For the fulfil numbersequal to unity the
 86 form of operator commutation relations does not change the finite results. That is why for the
 87 sake of simplicity one supposes all the operators being the Bose-Einstein field operators.

88 Instead of Schrödinger representation it will be convenient to use the equivalent inte-
 89 raction representation. If $\Psi(t)$ is the system wave function in Schrödinger representation
 90 than in interaction representation the wave function $\tilde{\Psi}(t)$ has the following view

$$91 \quad \tilde{\Psi}(t) = \exp\left(i\frac{\hat{H}^0}{\hbar}t\right)\Psi(t).$$

92 For the initial state in which the atom (1) is in its ground state and atom (2) is in excited state
 93 and photons are absent the view of wave function is the following

$$94 \quad \tilde{\Psi}^0 = \hat{b}_{i_g}^+ \hat{b}_{j_{ex}}^+ |0\rangle,$$

95 were $|0\rangle$ being the wave function of vacuum state. If the photon field differs from the va-
 96 cuumstate and any conglomerate of free photons with fulfil numbers $\mathbf{N}(\mathbf{k}) = \dots, N_{\mathbf{k}\lambda}, \dots$ is-
 97 placed in it than the wave function of such state will be denoted as $|\mathbf{N}(\mathbf{k})\rangle$. After the appear-
 98 ance in space of excited atom (2) the wave function $\tilde{\Psi}(t)$ of total system at any moment of
 99 time $t > t_0$ may be expressed as a set over the self-functions of \hat{H}^0 operator

$$100 \quad \tilde{\Psi}(t) = \sum_{ij} c_{ij}^{(1)}(t) \hat{b}_i^+ \hat{b}_j^+ |0\rangle + \sum_{ij\mathbf{N}(\mathbf{k})} c_{ij}^{(2)}(t, \mathbf{N}(\mathbf{k})) \hat{b}_i^+ \hat{b}_j^+ |\mathbf{N}(\mathbf{k})\rangle. \quad (2)$$

101 The summation over $\mathbf{N}(\mathbf{k})$ means the summation over all possible photon field conglome-
 102 rates. We are interested in the probability of exciting (1) atom at amoment of time $t > t_0$. Ac-
 103 cording to Dirac theory [2] the condition probability of such event by the transition at the
 104 same time of atom (2) at its ground state atthe absence of free photons in space is $\left|c_{i_{ex}j_g}^{(1)}(t)\right|^2$.

105 The condition probability of exciting (1) atom at a presence in space photons in state $|\mathbf{N}(\mathbf{k})\rangle$
 106 is $\left|c_{i_{ex}j_g}^{(2)}(t, \mathbf{N}(\mathbf{k}))\right|^2$. The total probability $P_{i_{ex}}(t)$ of the exciting of test atom (1) is the sum of
 107 condition probabilities

$$P_{i_{ex}}(t) = \sum_j \left| c_{i_{ex}j}^{(1)}(t) \right|^2 + \sum_{j \in \mathbf{N}(\mathbf{k})} \left| c_{i_{ex}j}^{(2)}(t, \mathbf{N}(\mathbf{k})) \right|^2 \quad (3)$$

One may use the another way and look for probability under consideration as a mean number of excited atoms in the state with energy $\varepsilon_{i_{ex}}$ if in system only one atom is present

$$P_{i_{ex}}(t) = \langle \tilde{\Psi}(t) | \hat{b}_{i_{ex}}^+ \hat{b}_{i_{ex}} | \tilde{\Psi}(t) \rangle. \quad (4)$$

If Fermi used [1] the formula (3) then in paper [6] one utilizes the formula (4). Both calculations have to lead to one and the same result since the acquaintance (3) follows from acquaintance (4) after introduction in it of the expression (2). The reason of results discrepancy in papers [2] and [6] is another. It is analyzed further.

Let us say that the square of matrix element $\left| c_{i_{ex}j_g}^{(1)}(t) \right|^2$ describes the probability of the excitation of (1) atom in coherent channel of atoms interaction. In this channel as a result of coherent process of reaction in space the free photons do not appear. Let us name the other channels of (1) atom excitation as no coherent. It follows from (3) that coherent channel of (1) atom excitation gives opportunity to estimate from the low value the total excitation probability of (1) atom

$$P_{i_{ex}}(t) \geq \left| c_{i_{ex}j_g}^{(1)}(t) \right|^2.$$

In Fermi's paper [1] the right side of this inequality is calculated. As it has shown in [5] the result of such calculation includes inside it the superluminal signal. Such signal can't be compensated by more precisely calculations.

If the probability of (1) atom excitation is calculated using formula (4) and interaction representation than one came across the formula (3) describing the presence of superluminal forerunner. On the other words the interaction representation with necessity predicts the appearance of superluminal forerunner. According to the paper [5] in Heisenberg representation the superluminal signals never appear. We state the none-identity of Heisenberg and Schrödinger representations in quantum electrodynamics of nonstationary processes. The reason of such nonidentity is investigated later.

Later we shall use other arguments which also lead to the conclusion on nonidentity of these representations and permit at the same time to clean the reason of nonidentity appearance.

In order to solve such problem let us calculate scalar product (4) in both interaction representation and Heisenberg representation. At the same time we shall pay attention on the reason of the discrepancy in such calculation results.

3. Interaction representation

The probability of (1) atom excitation in a form of scalar product (4) permits to calculate of such product in any quantum electrodynamics representation. In this paragraph we use the interaction representation. The Schrödinger equation in Schrödinger representation using the Hamiltonian (1) has a view

$$i\hbar \frac{\partial \Psi(t)}{\partial t} = \hat{H} \Psi(t).$$

In interaction representation the same equation has a form

$$i\hbar \frac{\partial \tilde{\Psi}(t)}{\partial t} = \hat{H}'(t) \tilde{\Psi}(t), \quad (5)$$

were

$$\hat{H}'(t) = -\frac{e}{mc} \int \hat{\psi}_1^+(x_1) \hat{p}_{r_1}^{\nu_1} \hat{A}^{\nu_1}(x_1) \hat{\psi}_1(x_1) d\mathbf{r}_1 - \frac{e}{mc} \int \hat{\psi}_2^+(x_2) \hat{p}_{r_2}^{\nu_2} \hat{A}^{\nu_2}(x_2) \hat{\psi}_2(x_2) d\mathbf{r}_2 \theta(t-t_0), \quad (6)$$

$$\hat{\psi}_1(x_1) = \sum_i \psi_i(\mathbf{r}_1 - \mathbf{R}_1) \hat{b}_i e^{-i\frac{\varepsilon_i}{\hbar}t}, \quad \hat{\psi}_2(x_2) = \sum_j \psi_j(\mathbf{r}_2 - \mathbf{R}_2) \hat{b}_j e^{-i\frac{\varepsilon_j}{\hbar}t},$$

$$\hat{A}^\nu(x) = \sum_{\mathbf{k}\lambda} \sqrt{\frac{\hbar c}{2kV}} e_{\mathbf{k}\lambda}^\nu \left(\hat{\alpha}_{\mathbf{k}\lambda} e^{i\mathbf{k}\mathbf{r} - ikct} + \hat{\alpha}_{\mathbf{k}\lambda}^+ e^{-i\mathbf{k}\mathbf{r} + ikct} \right).$$

Here ε_i and ε_j are the atom internal energies in consequence quantum states, $x = \{\mathbf{r}, t\}$. The solution of equation (5) has a view

$$\tilde{\Psi}(t) = \hat{S}(t) \tilde{\Psi}^0, \quad \hat{S}(t) = \hat{T} \left(\frac{1}{i\hbar} \int_{-\infty}^t \hat{H}'(t') dt' \right),$$

154 \hat{T} being chronological operator. The transformation of excitation from one atom to another in
 155 lowest order of perturbation theory is defined by the forth order. For such goal due to (4) the
 156 matrix $\hat{S}(t)$ has to be evaluated in the third order

$$157 \quad \hat{S}(t) = 1 + \hat{S}^{(1)}(t) + \hat{S}^{(2)}(t) + \hat{S}^{(3)}(t), \quad (7)$$

$$158 \quad \hat{S}^{(1)}(t) = \frac{1}{i\hbar} \int_{-\infty}^t \hat{H}'(t') dt', \quad \hat{S}^{(2)}(t) = \frac{\hat{T}}{2!} \left(\frac{1}{i\hbar} \int_{-\infty}^t \hat{H}'(t') dt' \right)^2, \quad \hat{S}^{(3)}(t) = \frac{\hat{T}}{3!} \left(\frac{1}{i\hbar} \int_{-\infty}^t \hat{H}'(t') dt' \right)^3. \quad (8)$$

159 The operators $\hat{S}^{(1)}(t)$ and $\hat{S}^{(3)}(t)$ describe no-coherent channels of reactions in which in
 160 space the excited atom (1) and free photons are present. The coherent channel of atom (1)
 161 excitation is described by operator $\hat{S}^{(2)}(t)$. The introduction (7) into (4) shows that

$$162 \quad P_{i_{ex}}(t) = \left\langle \hat{S}^{(2)}(t) \left| \hat{b}_{i_{ex}}^+ \hat{b}_{i_{ex}} \right| \hat{S}^{(2)}(t) \right\rangle + \left\langle \hat{S}^{(1)}(t) + \hat{S}^{(3)}(t) \left| \hat{b}_{i_{ex}}^+ \hat{b}_{i_{ex}} \right| \hat{S}^{(1)}(t) + \hat{S}^{(3)}(t) \right\rangle.$$

163 Let us calculate $\hat{S}^{(2)}(t)$. The introduction (6) into (8) leads to

$$164 \quad \hat{S}^{(2)}(t) = \left(\frac{e}{i\hbar mc} \right)^2 \int \hat{\psi}_1^+(x_1) \hat{p}_{\mathbf{r}_1}^{v_1} \hat{\psi}_1(x_1) \hat{\psi}_2^+(x_2) \hat{p}_{\mathbf{r}_2}^{v_2} \hat{\psi}_2(x_2) \cdot$$

$$165 \quad \cdot \left[i\hbar D^{v_1 v_2}(x_1, x_2) + \hat{N} \hat{A}^{v_1}(x_1) \hat{A}^{v_2}(x_2) \right] dx_1 dx_2. \quad (9)$$

166 Here we omitted the terms described the atoms self-action, \hat{N} is the normal product operator,
 167 $dx = d\mathbf{r} dt$. They used the conventional identity

$$168 \quad \hat{T} \hat{A}^{v_1}(x_1) \hat{A}^{v_2}(x_2) = i\hbar D^{v_1 v_2}(x_1, x_2) + \hat{N} \hat{A}^{v_1}(x_1) \hat{A}^{v_2}(x_2).$$

169 In its turn

$$170 \quad D^{v_1 v_2}(x_1, x_2) = D_r^{v_1 v_2}(x_1, x_2) + \Delta^{v_1 v_2}(x_1, x_2), \quad (10)$$

171 where $D_r^{v_1 v_2}(x_1, x_2)$ is the retarded Green function

$$172 \quad D_r^{v_1 v_2}(x_1, x_2) = \frac{1}{i\hbar} \left[\hat{A}^{v_1}(x_1); \hat{A}^{v_2}(x_2) \right] \theta(t_1 - t_2) = - \frac{\delta_{v_1 v_2} - n^{v_1} n^{v_2}}{4\pi |\mathbf{r}_1 - \mathbf{r}_2|} \delta \left(t_1 - t_2 - \frac{|\mathbf{r}_1 - \mathbf{r}_2|}{c} \right). \quad (11)$$

173 One supposes that the points \mathbf{r}_1 and \mathbf{r}_2 are divided by the wave radiation zone,

174 $n^\nu = (\mathbf{r}_1 - \mathbf{r}_2)^\nu / |\mathbf{r}_1 - \mathbf{r}_2|$. Further

$$175 \quad \Delta^{\nu_1 \nu_2}(x_1, x_2) = \frac{1}{i\hbar} \langle 0 | \hat{A}^{\nu_1}(x_1) \hat{A}^{\nu_2}(x_2) | 0 \rangle = -\frac{ic}{4\pi^2} \frac{\delta_{\nu_1 \nu_2} - n^{\nu_1} n^{\nu_2}}{|\mathbf{r}_1 - \mathbf{r}_2|} \int_0^\infty e^{ikc(t_1 - t_2)} \sin k |\mathbf{r}_1 - \mathbf{r}_2| dk. \quad (12)$$

176 The term in (9) containing the operator \hat{N} describes the no-coherent channel of reaction. In
 177 this channel besides an excited atom (1) the two free photons appear in space. The probability
 178 of such reaction is described by one of terms in the late sum in (3). This process we omit. In
 179 coherent channel according to (9)

$$180 \quad \hat{S}^{(2)}(t) = \hat{S}_1^{(2)}(t) + \hat{S}_2^{(2)}(t). \quad (13)$$

181 The first term contains function $D_r^{\nu_1 \nu_2}$ while the second one contains the function $\Delta^{\nu_1 \nu_2}$. The
 182 introduction (11) and (12) and (9) yields

$$183 \quad \hat{S}_1^{(2)}(t) = \frac{1}{i\hbar} \left(\frac{e}{mc} \right)^2 \int_{-\infty}^t p_{i_{ex} i_g}^{\nu_1} \exp \left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} t_1 \right) dt_1.$$

$$184 \quad \hat{b}_{i_{ex}}^+ \hat{b}_{i_g}^+ \hat{b}_{j_g}^+ \hat{b}_{j_{ex}}^+ \int_{-\infty}^t p_{j_g j_{ex}}^{\nu_2} \exp \left(-i \frac{\mathcal{E}_{j_{ex}} - \mathcal{E}_{j_g}}{\hbar} t_2 \right) D_r^{\nu_1 \nu_2}(\mathbf{R}_1, \mathbf{R}_2, t_1, t_2) \theta(t_2 - t_0) dt_1 dt_2,$$

$$185 \quad \hat{S}_2^{(2)}(t) = -\frac{1}{\hbar} \left(\frac{e}{mc} \right)^2 \frac{c}{4\pi^2} \int_{-\infty}^t p_{i_{ex} i_g}^{\nu_1} \exp \left(\frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} t_1 \right) dt_1.$$

$$186 \quad \hat{b}_{i_{ex}}^+ \hat{b}_{i_g}^+ \hat{b}_{j_g}^+ \hat{b}_{j_{ex}}^+ \int_{-\infty}^t p_{j_g j_{ex}}^{\nu_2} \exp \left(-i \frac{\mathcal{E}_{j_{ex}} - \mathcal{E}_{j_g}}{\hbar} t_2 \right) \int_0^\infty \frac{\delta_{\nu_1 \nu_2} - n^{\nu_1} n^{\nu_2}}{|\mathbf{R}_1 - \mathbf{R}_2|} \sin k |\mathbf{R}_1 - \mathbf{R}_2| e^{ikc(t_1 - t_2)} dk \theta(t_2 - t_0) dt_1 dt_2$$

187 , (14)

$$188 \quad p_{i_g i_{ex}}^\nu = \int \psi_{i_g}^*(\mathbf{p}) \hat{p}_\mathbf{p}^\nu \psi_{i_{ex}}(\mathbf{p}) d\mathbf{p}.$$

189 The introduction (14) in (4) shows that

$$190 \quad P_{i_{ex}}(t) = \left\langle \hat{S}_1^{(2)} + \hat{S}_2^{(2)} \left| \hat{b}_{i_{ex}}^+ \hat{b}_{i_{ex}} \right| \hat{S}_1^{(2)} + \hat{S}_2^{(2)} \right\rangle_0 + \left\langle \hat{S}^{(1)} + \hat{S}^{(3)} \left| \hat{b}_{i_{ex}}^+ \hat{b}_{i_{ex}} \right| \hat{S}^{(1)} + \hat{S}^{(3)} \right\rangle_0. \quad (15)$$

191 The quantum averaging process in this equality is performed over initial state of system. The
 192 operator $\hat{S}_1^{(2)}(t)$ does not contain superluminal forerunner while in operator $\hat{S}_2^{(2)}(t)$ such fore-
 193 runner is present.

194 4. Heisenberg representation

195 The transposition from Schrödinger representation to the Heisenberg representation is per-
 196 formed by operator $\hat{U}(t)$ satisfying the equation

$$197 \quad i\hbar \frac{\partial \hat{U}(t)}{\partial t} = (\hat{H}^0 + \hat{H}') \hat{U}(t). \quad (16)$$

198 The field operators in Heisenberg representation have a view

$$199 \quad \overset{\vee}{\psi}(x) = \hat{U}^+(t) \hat{\psi}(\mathbf{r}) \hat{U}(t), \quad \overset{\vee}{A}^v(x) = \hat{U}^+(t) \hat{A}^v(\mathbf{r}) \hat{U}(t), \quad \overset{\vee}{b}_{i_{ex}}(x) = \hat{U}^+(t) \hat{b}_{i_{ex}} \hat{U}(t).$$

200 The differential equation (16) may be transformed to the integral one

$$201 \quad \hat{U}(t) = \hat{U}^0(t) + \frac{1}{i\hbar} \hat{U}^0(t) \int_{-\infty}^t \hat{U}^0(t') \hat{H}'(t') \hat{U}(t') dt', \quad \hat{U}^0(t) = e^{-i \frac{\hat{H}^0}{\hbar} t}.$$

202 By using twice the iterative procedure we obtain [21] for the operator $\overset{\vee}{b}_{i_{ex}}(t)$

$$203 \quad \overset{\vee}{b}_{i_{ex}}(t) = \hat{b}_{i_{ex}}(t) + \frac{1}{i\hbar} \int_{-\infty}^t [\hat{b}_{i_{ex}}(t); \hat{H}'(t')] dt' + \frac{1}{(i\hbar)^2} \int_{-\infty}^t \int_{-\infty}^t \theta(t' - t'') [\hat{b}_{i_{ex}}(t); \hat{H}'(t')] \hat{H}'(t'') dt' dt'' + o(e^3)$$

204 ,

205 were

$$206 \quad \hat{b}_{i_{ex}}(t) = \hat{b}_{i_{ex}} e^{-i \frac{\hat{H}^0}{\hbar} t}.$$

207 By using the explicit form of operators $\hat{H}'(t)$, $\hat{\psi}(x)$ and $\hat{\psi}^+(x)$ in dipole approximation one
 208 yields

$$209 \quad \overset{\vee}{b}_{i_{ex}}(t) = \hat{b}_{i_{ex}} e^{-i \frac{\mathcal{E}_{i_{ex}}}{\hbar} t} - \frac{e}{i\hbar mc} e^{-i \frac{\mathcal{E}_{i_{ex}}}{\hbar} t} \int_{-\infty}^t p_{i_{ex} i_g}^{v_1} \exp\left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} t'\right) \hat{A}^{v_1}(\mathbf{R}_1, t') dt' \hat{b}_{i_g} + \left(\frac{e}{i\hbar mc}\right)^2 e^{-i \frac{\mathcal{E}_{i_{ex}}}{\hbar} t}.$$

$$\begin{aligned}
 & \cdot \int_{-\infty}^t \int_{t_0}^t p_{i_{ex}i_g}^{v_1} \exp\left(i \frac{\mathcal{E}_{ex} - \mathcal{E}_{i_g}}{\hbar} t'\right) p_{j_gj_{ex}}^{v_2} \exp\left(-i \frac{\mathcal{E}_{j_{ex}} - \mathcal{E}_{j_g}}{\hbar} t''\right) D_r^{v_1v_2}(\mathbf{R}_1, t', \mathbf{R}_2, t'') dt' dt'' \hat{b}_{i_g}^+ \hat{b}_{j_g}^+ \hat{b}_{j_{ex}} + o(e^3) \\
 & .(17)
 \end{aligned}$$

Now it is evident that

$$\begin{aligned}
 P_{i_{ex}}(t) = & \left\langle b_{i_{ex}}^+(t) \left[-\frac{e}{i\hbar mc} e^{-i \frac{\mathcal{E}_{i_{ex}}}{\hbar} t} \int_{-\infty}^t p_{i_{ex}i_g}^{v_1} \exp\left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} t'\right) \hat{A}^{v_1}(\mathbf{R}_1, t') dt' \hat{b}_{i_g} + \left(\frac{e}{i\hbar mc}\right)^2 e^{-i \frac{\mathcal{E}_{i_{ex}}}{\hbar} t} \right. \right. \\
 & \left. \left. \cdot \int_{-\infty}^t \int_{t_0}^t p_{i_{ex}i_g}^{v_1} \exp\left(i \frac{\mathcal{E}_{ex} - \mathcal{E}_{i_g}}{\hbar} t'\right) p_{j_gj_{ex}}^{v_2} \exp\left(-i \frac{\mathcal{E}_{j_{ex}} - \mathcal{E}_{j_g}}{\hbar} t''\right) D_r^{v_1v_2}(\mathbf{R}_1, t', \mathbf{R}_2, t'') dt' dt'' \hat{b}_{i_g}^+ \hat{b}_{j_g}^+ \hat{b}_{j_{ex}} + o(e^3) \right] \right\rangle_0 .
 \end{aligned}$$

Here the quantum averaging is performed over initial state of system.

5. The discussing of the results

The formulae (15) and (17) being calculated in different representations describe one and the same probability $P_{i_{ex}}(t)$. If in (15) the omitted term containing \hat{N} is reconstructed than in $\sim e^4$ approximation (15) and (17) evidentially would be equal. But in the present form they are senseless since they contain in infinite limits the integrals from oscillated functions. It is necessary to use the adiabatic hypothesis[22]. We stress that for the acquaintance (15) and (17) expressions it is necessary to take into account all the terms proportional to $\sim e^4$, and among them the term following from product of first order term on the third one. If they neglect of such term, that is necessary for coinciding with adiabatic hypothesis, than the results will be different.

The analysis in detail we began from formula (17) obtained in Heisenberg representation. The first term in this formula which is proportional to $\sim e^2$ describes the (1) atom excitation due to its interaction with electromagnetic vacuum. Such fact of not equality to zero the probability in question contradicts to the initial condition $\hat{b}_{i_g}^+ \hat{b}_{j_{ex}}^+ |0\rangle$. Besides this fact the electromagnetic vacuum cannot excite the atom being in its ground state according to the physical understanding. The probability of such processes has to be equal to zero. In conventional quantum electrodynamics such excitation is absent since it contradicts the law of energy conservation. The law of energy conservation follows from the adiabatic hypothesis that is additionally put-

234 ted on the solutions of quantum electrodynamics. Mathematically this hypothesis is expresses
235 by the equality

$$236 \quad \delta(\omega - \omega_0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(\omega - \omega_0)t} dt,$$

237 $\delta(\omega - \omega_0)$ being Dirac function. In its turn this equality demands the integration over the time
238 in infinite limits. Only the additional using of adiabatic hypothesis turns the set of perturba-
239 tion theory to the physically sense. But in the problem under consideration the using of adia-
240 batic hypothesis in its usual form is impossible since the variable t is finite. On the other
241 hand the atom (1) before the interaction with excited atom (2) was in its ground state the infi-
242 nitely long time interval $(-\infty \div t)$ permanently interacting with electromagnetic vacuum. The
243 time length of the interaction interval from the physically point of view is infinitely long. We
244 use this circumstance to investigate of the problematic right side term in (17)

$$245 \quad \int_{-\infty}^t \hat{A}^\nu(\mathbf{R}_1, t') \exp\left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} t'\right) dt' =$$

$$246 \quad = \sum_{\mathbf{k}\lambda} \int_{-\infty}^t \sqrt{\frac{\hbar c}{2kV}} e_{\mathbf{k}\lambda}^\nu \exp\left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} t'\right) \left(\hat{\alpha}_{\mathbf{k}\lambda} e^{i\mathbf{k}\mathbf{R}_1 - ikct'} + \hat{\alpha}_{\mathbf{k}\lambda}^+ e^{-i\mathbf{k}\mathbf{R}_1 + ikct'} \right) dt'. \quad (18)$$

247 It is necessary to pay attention to the fact that the probability of excitation transposition be-
248 tween (2) and (1) atoms does not depend on the time t but only on the time difference $t - t_0$.
249 Taken into account that the interaction of the atom (1) with electromagnetic field up to the
250 time t_0 has the infinitely long duration it necessary to pose that the physical mining the ex-
251 pression (17) has only in the limit $t \rightarrow \infty$. At the same time the difference $t - t_0$ rests constant
252 (general adiabatic hypothesis). Now from (18) yields

$$253 \quad \lim_{t \rightarrow \infty} \int_{-\infty}^t \hat{A}^\nu(\mathbf{R}_1, t') \exp\left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} t'\right) dt' =$$

$$254 \quad = 2\pi \sum_{\mathbf{k}\lambda} \sqrt{\frac{\hbar c}{2kV}} e_{\mathbf{k}\lambda}^\nu \left(\hat{\alpha}_{\mathbf{k}\lambda} e^{i\mathbf{k}\mathbf{R}_1} \delta\left(\frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} - kc\right) + \hat{\alpha}_{\mathbf{k}\lambda}^+ e^{-i\mathbf{k}\mathbf{R}_1} \delta\left(\frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} + kc\right) \right).$$

255 This expression carries in the result zero contributionsince the free photons are absent in
 256 space. The vacuum term transforms into zero due to the energy conservation law. Now it is
 257 evident that the product of the first term of perturbation theory by the third one also turns into
 258 zero. In approximation $\sim e^4$ only one term rests

259 $P_{i_{ex}}(t) =$

260
$$= \frac{1}{\hbar^2} \left(\frac{e}{mc} \right)^4 \left| \int_{-\infty}^t \int_{t_0}^t p_{i_{ex}i_g}^{V_1} \exp \left(i \frac{\mathcal{E}_{ex} - \mathcal{E}_{i_g}}{\hbar} t' \right) p_{j_gj_{ex}}^{V_2} \exp \left(-i \frac{\mathcal{E}_{j_{ex}} - \mathcal{E}_{j_g}}{\hbar} t'' \right) D_r^{V_1V_2}(\mathbf{R}_1, t', \mathbf{R}_2, t'') dt'' dt' \right|^2. \quad (19)$$

261 This result found in Heisenberg representation being equal to the result of paper [6] does not
 262 contain of the superluminal forerunners. This result may be explained as the one photon radi-
 263 ation by the atom (2) at time moment t'' and the absorption of this photon by the atom (1) at a
 264 time moment t' . The propagator

265
$$D_r^{V_1V_2}(\mathbf{R}_1, \mathbf{R}_2, t', t'') \sim \delta \left(t' - t'' - \frac{|\mathbf{R}_1 - \mathbf{R}_2|}{c} \right)$$

266 looks after carry out the condition $c(t' - t'') = |\mathbf{R}_1 - \mathbf{R}_2|$.

267 In the interaction representation we came across the same mathematical problem by calcula-
 268 tion the operator (8)

269
$$\hat{S}^{(1)}(t) = -\frac{e}{i\hbar mc} \int \hat{\psi}_1^+(x_1) \hat{p}_{r_1}^v \hat{A}^v(\mathbf{R}_1, t_1) \hat{\psi}_1(x_1) d\mathbf{r}_1 dt_1 =$$

270
$$= -\frac{e}{i\hbar mc} \int_{-\infty}^t p_{i_{ex}i_g}^v \exp \left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} t' \right) \hat{A}^v(\mathbf{R}_1, t') dt'.$$

271 In the limit $t \rightarrow \infty$ by condition $t - t_0 = const$ one gets $\hat{S}^{(1)}(t) \rightarrow 0$ if in space the free pho-
 272 tons are absent. Now in (15) one gets $\left\langle \hat{S}^{(1)}(t) \hat{b}_{i_{ex}}^+ \hat{b}_{i_{ex}} \hat{S}^{(3)}(t) \right\rangle_0 = 0$.

273 Let us consider now the operator $\hat{S}_2^{(2)}(t)$. In this operator according (14) integration over in-
 274 termedia variables t_1 captures the area $t_1 < t_0$. Let us divide the integral over t_1 in (14) by the
 275 sum of two integrals

$$\int_{-\infty}^{t_0} \exp\left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g} + kc\hbar}{\hbar} t_1\right) dt_1 + \int_{t_0}^t \exp\left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g} + kc\hbar}{\hbar} t_1\right) dt_1.$$

But the limit transition $t \rightarrow \infty$ if $t - t_0 = const$ demands the limit transition $t_0 \rightarrow \infty$. In this case the first integral transforms in Dirac δ -function $\delta\left(kc + (\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g})/\hbar\right)$ which is equal to zero due to the positive value of its argument. The expression (15) describing the probability of atom (1) excitation in approximation $\sim e^4$ is now rewritten in the following view

$$P_{i_{ex}}(t) = \frac{1}{\hbar^2} \left(\frac{e}{mc}\right)^4 \left| \int_{t_0}^t p_{i_{ex}i_g}^{v_1} \exp\left(i \frac{\mathcal{E}_{i_{ex}} - \mathcal{E}_{i_g}}{\hbar} t_1\right) \int_{t_0}^t p_{j_gj_{ex}}^{v_2} \exp\left(-i \frac{\mathcal{E}_{j_{ex}} - \mathcal{E}_{j_g}}{\hbar} t_2\right) dt_2 \right|^2. \quad (20)$$

$$\cdot \left[D_r^{v_1v_2}(\mathbf{R}_1, \mathbf{R}_2, t_1, t_2) + \frac{i}{8\pi^2} \frac{\delta_{v_1v_2} - n^{v_1}n^{v_2}}{|\mathbf{R}_1 - \mathbf{R}_2|} \left(\frac{1}{t_1 - t_2 - \frac{|\mathbf{R}_1 - \mathbf{R}_2|}{c} + i0} - \frac{1}{t_1 - t_2 + \frac{|\mathbf{R}_1 - \mathbf{R}_2|}{c} + i0} \right) \right] dt_1 dt_2 \Bigg|^2.$$

Here the first term coincides with the result (19) obtained in Heisenberg representation. The second one describes the signals placed in superluminal zone at a distance of the order of one wave length that coincides with corrective Fermi calculations. In the limit $t_0 \rightarrow -\infty, t \rightarrow \infty$ the second term turns into zero due to integrands analytical properties. By this reason in stationary problems the representations Schrödinger and Heisenberg are identical. In nonstationary conditions formulae (20) and (19) calculated in different representations are not coincide.

The formulae (19) and (20) arrived from formulae (15) and (17) if in the last one according to the general adiabatic hypothesis one misses the terms appearing from the products of the first order term of perturbation theory by the third one. By this reason these formulae cannot be equal. Other words the using of the general adiabatic hypothesis leads to non-equivalency of Schrödinger and Heisenberg representations in non-stationary quantum electrodynamics. We stress that the Schrödinger representation permits the appearance of superluminal forerunners.

6. Conclusion

In this work the non-stationary processes of transformation excitation from one atom to another is considered. The result of Fermi work [2,5] in which the matrix element for such

process was calculated permits to think about the principal presence in nature the superluminal signals. The repeated calculation [6] of this process probability performed by using Heisenberg representation led to the conclusion of the absence of superluminal signals in quantum theory. In the same work they postulated the no corrections of quantum transposition calculation as a square of corresponded matrix element. The other words they doubt about the Dirac theory of quantum transpositions [1].

It is shown in present work that the calculation of quantum transposition probability as matrix elements squared (Dirac's method) or as quantum average of corresponded quantum operators lead to identity results if last calculations are performed in Schrödinger representation.

Different results found in papers [2,5] and [6] are not the consequences of different probabilities definition. The results different is the consequences of non-identity Schrödinger and Heisenberg representation in quantum electrodynamics of nonstationary processes. As a proof of non-identity representations in present work the probability of one atom excitation by spontaneous radiation of another atom expressed through quantum averaging of corresponded operators is calculated. The calculations of such quantum averaging are performed by both Schrödinger and Heisenberg representations leading to the different results. The representations nonidentity follows finely from the no correct definition of scattering matrix $\hat{S}(t)$ creating the connection of interaction (Schrödinger) and Heisenberg representations. Since the product $\hat{S}(t)|\Phi\rangle$ where Φ is arbitrary wave function in quantum electrodynamic is represented as a divergent set that is non astonishing that the different summation set methods lead to different results. By using of the formal properties of $\hat{S}(t)$ operator the sets of perturbation theory obtained in Schrödinger and Heisenberg representations at first glance are equal. But such sets do not represent sensible solutions of quantum electrodynamics. In order to put them the physical sense it is necessary to use the adiabatic hypothesis which supposes switching and shutting off the interaction at $t \rightarrow \pm\infty$. This hypothesis mathematically expressed by using the following equality

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(\omega-\omega_0)t'} dt' = \delta(\omega-\omega_0) \quad (21).$$

By investigating of quantum transitions at finite time intervals it is not possible to use the conventional adiabatic hypothesis. Instead this hypothesis it is necessary to use its generation in the form

$$\lim_{t \rightarrow \infty} \frac{1}{2\pi} \int_{-\infty}^t e^{i(\omega - \omega_0)t'} dt' = \delta(\omega - \omega_0).$$

At the same time as in conventional quantum electrodynamics it is necessary to watch the order of carry out the mathematical operations. First of all it is necessary to carry out the limit transition (21) and only then to carry out the quantum operation of summation $\langle \dots \rangle$. After using the general adiabatic hypothesis the sets of perturbation theory lead to reasonable results. But such results obtained in Schrödinger and Heisenberg representations are different. The difference may appear already in the terms proportional to $\sim e^4$.

The representation nonidentity is worth in practical aspect. As is shown above the Schrödinger representation predicts the presents in the nature of specific quantum superluminal signals. The Heisenberg representation cannot describe the superluminal processes at all. In connection with experimentally observed superluminal phenomena such property of Schrödinger representation possesses the real interest which arises in connection with the work [12]. In this experimental work the authors investigate the superluminal signals on the example of radiation evolution primary formed in quantum compressed state.

Due to nonidentity of Schrödinger and Heisenberg representations the theories using these representations have to be considered as two mutual non-connecting theories. The physical systems in which the matrix $\hat{S}(t)$ is well definite are quasi-classical in the sense of non-possibility inside them the superluminal signals and Schrödinger and Heisenberg representations for such systems are identical. In general case the choice of one of these representations only the experiment may show. At present time only one such experiment is known [13] which shows on Schrödinger representation and predicts at the same time the existence in quantum electrodynamics the superluminal signals.

The existence of the superluminal signals does not break [23] the causality principle. It is necessary the causality principle to understand in the following form: the consequences can't act on their reasons. The Lorentz invariance of quantum electrodynamics equations is not the obstacle for superluminal signals appearance.

References

1. Fermi E.// Rev. Mod. Phys. 1932. V.4. P. 87-132.
2. Dirac P.A.M.// Proc.Roy.Soc. 1927. V.A114, P. 243-265.

- 359 **3.** Breit G.//Rev. Mod. Phys. 1933.V.5.P.91-140.
- 360 **4.** Milonni P.W., Knight P.L.// Phys.Rev. 1974. V.A10. P. 1096-1108.
- 361 **5.**ShirokovM.I. Sov.Phys. Usp.1978..V.21. P. 345-358.
- 362 **6.** BykovV.P., ZadernjvskiiA.A.// JETP1970.V.57.P.20-24.
- 363 **7.**BasovN.G., AmbartzumjanR.B., ZuevV.S., KryukovP.G., LetochovV.S. //JETP1966. V.50.
- 364 P.23-34(RF).
- 365 **8.** Chu S., Wong S. // Phys.Rev.Lett. 1982. V.48. P.738-741.
- 366 **9.** Steinberg A.M., Kwiat P.G., Chiao R.Y. //Phys. Rev.Lett. 1993. V,71. P.708-711.
- 367 **10.** AkulshinA.M., ChimmimoA.,OpatDj.I. Quantum Electronics. 2002. V. 32. P.567-569.
- 368 **11.** Wang L.J., Kuzmich A., Dogariu A.// Nature 2000. V.406. P.277-279.
- 369 **12.**Song K.Y., Herraez M.G., Thevenaz L.// Optics Express 2005. V.13. P.82-88.
- 370 **13.**Veklenko B.A., MslakhovYu.I.,NguenK.Sh.// Engineering Physics 2013. №5.P.25-
- 371 39.(RF).
- 372 **14.** Romanov G., Horrom**T.**, Novikova I., Mikhailov E.E.//Optics Lett.2014. V.39. P.1093-
- 373 1096.
- 374 **15.** Samson**A.M.**//DokladiBSSR1966. V.10.P.739-743 (RF).
- 375 **16.**KryukovP.G., LetokhovV.S. //Sov.Phys. Usp.1970. V.12. P.641-672.
- 376 **15.**ChiaoR.Y.// Phys. Rev. 1993. V.A48. R34-R37.
- 377 **16.**SteinbergA.M., ChiaoR.Y.// Phys.Rev. 1994. V.A49. P.2071-2075.
- 378 **17.** Garrett C.G.B., McCumber D.E.// Phys.Rev. 1970. V.A1. P.305-313.
- 379 **18.**Oraevskii A.N. //Phys.Usp.1998. V.41. P.1199-1209.
- 380 **19.** Zurita-Sanchez J.R., Abundis-Patino J.H., Halevi P.// Optics Express 2012. V.20.
- 381 P.5586-5600.
- 382 **20.** Veklenko B.A. //Applied Physics2010. №3.P.10-17 (RF).
- 383 **21.** Schwinger J.// Phys.Rev. 1948. V.47. P.1439-1461.
- 384 **22.** Akhiezer A.I., Berestetskii V.B. Quantum Electrodynamics. Moskow.Nauka. 1969. P.
- 385 217.
- 386 **23.**KadomtsevB.B. Phys.Usp. 1994.V.37. P.425-499.