

Variability of Horizontal Magnetic Field Intensity from some

Stations within the Equatorial Electrojet Belt

ABSTRACT

The variability of the horizontal component of magnetic field intensity from 7 stations within the Equatorial electrojet (EEJ) belt ($\pm 3^\circ$ of the magnetic equator) has been investigated using year 2008 data measured from the Magnetic Data Acquisition System (MAGDAS) magnetometers. The monthly mean hourly values show variation in the pre-sunrise hours, within the range of 1nT to 21nT. However, dH during the day has a much higher variation than those observed during the pre-sunrise and dusk hours. The presence of Counter electrojet (CEJ) in the morning and evening hours were also observed. Also, equinoctial maximum was observed in most stations, and the horizontal magnetic field intensity over ANC located in the South American sector displays the highest amplitude, while ILR in the African sector appears to have the lowest amplitude.

Keywords: *Equatorial electrojet, Magnetic field intensity, Variability*

1. INTRODUCTION

The regular daily fluctuation in the earth's magnetic field was first observed by Graham [1]. Stewarts [2] later posited that these regular daily fluctuations in the geomagnetic field originate from thermally forced motion of conducting air moving over the magnetic field in the ionosphere. Chapman [3] noticed the existence of solar quiet current system in the ionosphere around 100km altitude, which is caused by the dominant solar-driven wind and tidal motion in the ionosphere. This current produced by electromotive forces is frequently considered to be due to the action of dynamo, which results in ground perturbation.

In 1922, the geomagnetic observatory located in Huancayo Peru, the western part of South America, where the alignment of the geomagnetic field is almost along the geographic meridian, played a crucial role in equatorial electrojet discovery. The horizontal (H) component of the geomagnetic field at Huancayo was noticed to by McNish [4] to be abnormally large in comparison to Chapman [3] current system, which was based on data from mid-latitudes. Egedal [5], noted that there exist a daytime east-west overhead current within a narrow latitude belt of about 130km over the magnetic dip equator, where the geomagnetic field is horizontal. Chapman [6] called this, the Equatorial Electrojet.

Thus, Equatorial Electrojet is an intense ribbon of current flowing eastward in the low latitude ionosphere within the narrow region flanking the dip equator ($\pm 3^\circ$ of the magnetic equator). The infrequent reversal of the electrojet current at certain hours of the day is described as Counter electrojet, Gouin and Mayaud [7].

Several studies have been carried out on SqH and EEJ using different observation and experimental tools. The investigation of worldwide solar quiet of horizontal component (SqH) at different seasons by Owolabi et al. [8] showed that SqH exhibits transient variations, with the amplitude varying at different seasons. However, in the past, there has been an acute shortage of observational instrument for magnetic field study in Africa, which has limited the study of longitudinal variations of EEJ to Asia and American Sector only but the deployment and distribution of Magnetic Data Acquisition System (MAGDAS) magnetometers to Africa after the International Heliophysical Year in 2005, has provided more data for magnetic field study in the Africa. Hence, this study employs

44 data from Magnetic Data Acquisition System (MAGDAS) magnetometers located in stations along the
45 magnetic equator to study the variability of H field intensity.
46


47
48 **2. MATERIAL AND METHODS**

49 The year 2008 data of the Magnetic Data Acquisition System (MAGDAS) magnetometers of some
50 stations within the equatorial electrojet belt were used for this study. Figure.1. present the global
51 distribution of MAGDAS magnetometers while the coordinates of the stations used in the study are
52 given in Table. 1.

53 Magnetic data corresponding to five most magnetically quiet days of each month in year 2008
54 were selected and used in this study. The international quiet days were published on Geoscience
55 Australia website based on the magnetic activity index Kp. During quiet time, the horizontal
56 component of the magnetic field within the EEJ belt is a combination of Sq current and EEJ current.

57 The baseline of each day was defined as the mean of the H component of the four (4) hours
58 flanking the local midnight time of each station. By subtracting this baseline values from the H
59 component gives the hourly departure (dH).

60



61
62
63
64
65
66
67
68
69
70
71

Fig. 1. The global network of MAGDAS magnetometers

72
73
74
75
76
77

Table 1: Parameters of the stations used in the study

S/N	Station	Code	Geographic Latitude	Geographic Longitude	Geomagnetic Latitude	Geomagnetic Longitude	Dip Latitude
1	Addis Ababa	AAB	9.04	38.77	0.18	110.47	0.57
2	Ancon	ANC	-11.77	-77.15	0.77	354.33	0.74
3	Cebu	CEB	10.36	123.91	2.53	195.06	2.74
4	Davao	DAV	7.00	125.40	-1.02	196.54	-0.65
5	Ilorin	ILR	8.50	4.68	-1.82	76.80	-2.96
6	Langkawi	LKW	6.30	99.78	-2.32	171.29	-1.88
7	Yap Island	YAP	9.50	138.08	1.49	209.06	1.70

78

79 The baseline values were obtained using the expression:

80
$$H_b = \frac{H_{2300} + H_{2400} + H_{0100} + H_{0200}}{4} \quad (1)$$

81 Where H_b is the baseline value and, H_{2300} , H_{2400} , H_{0100} , and H_{0200} are the H values at 2300 LT, 2400
82 LT, 0100 LT and 0200LT respectively.

83 The hourly departures of H were obtained by deducting the baseline values for each station
84 on each quiet day from the hourly values (H_t) of the corresponding station and day.

85
$$dH_t = H_t - H_b \quad (2)$$

86

87 Where $t = 1 \dots 24$

88 The monthly mean values of each month were derived from the mean of the diurnal variations
89 for the five quietest days in each month, and were plotted in multiple series with respect to the local
90 time of each station.

91 The seasonal variations were also studied by classifying the year into three seasons, that is,
92 the Equinox Season (March, April, September, October), denoted as E- season, June Season (May,
93 June, July, August) , denoted as J-season and December Season (November, December, January,
94 February) which is denoted as D -season. The Seasonal values were deduced from the mean of all
95 the months in a particular season, after which the seasonal variations were presented in contour filled
96 form, in the order of increasing geomagnetic longitude. That is, from ILR to ANC. Also, we obtained
97 the noon time variation of dH across the stations for the seasons, using a group bar plot. Since dH
98 over the equatorial region is expected to reach its peak value at about local noon as a result of
99 increase in ionization by solar activity in accordant with atmospheric dynamo theory, Rabiou et al. [9];
100 Onwumechili [10].

3. RESULTS AND DISCUSSION

3.1 Monthly Variation

Fig.2. shows the monthly mean of the diurnal variation from January 2008 to December 2008, plotted in multiple series for some equatorial electrojet stations. The diurnal variation of all the station shows a regular and consistent variation throughout the year. It was observed that dH rises from 0700LT, reaching its peak at about local noon and leaning back to minimum values at around 1700LT. This result is in agreement with the works of Rigoti et al. [11], Chandra et al. [12] and Rastogi [13].

At pre-sunrise hours, dH is in the range of 1 to 21nT, where the maximum value of 21nT was noticed at LKW in August around 0500LT and the minimum value of ~ 1nT was observed at other longitudes for different months of the year. From the results in fig.2, some counter electrojet (CEJ) events were recorded between the pre-sunrise and early morning hours and also in the evening hours. This CEJ varies in amplitude (from -1nT to -42nT), the highest CEJ amplitude of -42nT was observed around 0800LT, May 2008 at AAB. The range in dH peak during the day is higher than the range observed in the pre-dawn and dusk hours. For instance, from Fig. 2, In March 2008, the maximum dH amplitude for the longitudes considered varies from about 59nT to 138nT (range ~ 79nT). In June 2008, dH varies from about 39nT to 85nT (range ~ 46nT). In October 2008, dH varies from 42nT to 121nT (range ~ 79nT). A quasi character is also noticed in dH amplitudes for other months. However, as observed from Fig.2 ANC was observed to have the highest midday peak than the other stations.





Fig. 2. Monthly mean daily variations of H field for five International Quiet days



129

130 Fig. 2 continued

131

132 3.2 Seasonal Variation

133 Figures 3, 4 and 5, shows that dH was observed to display seasonal variation across the
 134 longitudes. Fig. 3 gives the result for the variation of dH during E-Season. It was observed that the
 135 amplitude of dH is highest at ANC reaching about 116nT at 1200LT, followed by a magnitude of 83nT
 136 at DAV. A West African station, ILR display a lower variation in comparison to other stations used.
 137 This shows that the amplitude of dH increases with increasing longitude, except at YAP which has a
 138 lower amplitude than DAV. Some CEJ events were recorded at ILR and AAB in the pre-sunrise to
 139 sunrise.

140 From fig. 4, during J-Season, ANC was also observed to be the highest at about 1200LT with
 141 amplitude of 83nT, while ILR and AAB has the minimum dH, and a CEJ of about -23nT at 0800LT
 142 was observed at AAB station.

From fig. 5, during D-Season, ANC and DAV exhibits the maximum dH amplitude as both peaks at 1200LT with 101nT, where ILR and AAB has the minimum dH amplitude. It was observed that the latitudinal position of the stations also affects the magnetic field variability. For instance, in fig. 5 CEB and DAV are on a very close meridian, but DAV exhibits higher magnetic field amplitude than CEB, this is likely because DAV which has geomagnetic latitude of -1.02° is closer to the magnetic equator than CEB with geomagnetic latitude of 2.53° .




Fig. 3. dH during E-Season




Fig. 4. dH during J-Season




Fig. 5. dH during D-Season




Fig. 6. Seasonal variation of noon-time dH across the stations

The group plot of the noontime seasonal variation of dH across different longitude is presented in Fig. 6. It is shown that at noon time ANC exhibits maximum dH for all seasons, with the highest amplitude during the E-season and the minimum during J-Season. In comparison to other seasons, J-Season exhibits the weakest dH at noontime for all the stations, while E-Season is maximum for most stations. The equinoctial maximum in most stations can be ascribed to intensified electron density at equatorial region during equinox seasons, due to solar activity taking place directly on the equator.

4. CONCLUSION

This study has confirmed that the horizontal component of the magnetic field along the magnetic equator exhibits longitudinal variation. The range of variation is usually large and can be up to 79nT in some cases. ANC, in the South American sector has the most pronounced magnitude of dH in all seasons while the amplitude of dH appears to be weakest in ILR. Also, the longitudinal position of the stations affects the magnetic field intensity, as the horizontal component of the magnetic field intensity increases with increasing longitude, with respect to its closeness to magnetic equator. The seasonal variations can be ascribed to the reposition of the ionospheric current system with seasonal changes.

REFERENCES

1. Graham G. An account of observations made of the variation of the horizontal needle at London, in the latter part of the year 1722, and beginning of 1723, Philos. Trans. K Soc. London, 1724; 383, 96-107.
2. Stewart B., Hypothetical views regarding the connection between the State of the sun and terrestrial magnetism, in Encyclopedia Britannica, 9th ed., vol. 16, 1882; pp. 181-184.
3. Chapman S. The solar and lunar diurnal variations of terrestrial magnetism, Phil. Trans. Roy. Soc. 1919; A(218):1-118.
4. McNish A.G., Bull 10. Int. Assoc. Magn. Transaction of Edinburgh meeting, Copenhagen; 1937; pp 271-280.
5. Egedal J. Magnetism and Atmospheric Electricity, journal of Terrestrial physics, 1947; 52, 449-451.
6. Chapman S. The equatorial electrojet as detected from the abnormal electric current distribution above Huancayo and elsewhere, Arch. Meteorol. Geophys. Bioclimatol, 1951; A 4: 368-392.
7. Gouin P. and Mayaud P.N., A proposed e l'existence possibled 'un"contre electrojet"a ux latitudes magnetiques quatoriales, Ann. Geophys, 1967; 2, 3, 41-47.
8. Owolabi T. P., Rabiou A. B., Olayanju G. M. and Bolaji O. S., Seasonal Variation of Worldwide Solar Quiet of the Horizontal Magnetic Field Intensity, Applied Physics Research, 2014; Vol. 6, No. 2.
9. Rabiou A.B., Adimula I.A., Yumoto K., Adeniyi J.O., Maeda G., preliminary results from magnetic field measurements Using MAGDAS at Ilorin, Nigeria, Earth, Moon, and Planets. DOI:10.1007/s11038-008-9290-7. 2009; 104, 173.
10. Onwumechili C. A. and Ezema P. O., On the course of the geomagnetic daily variation in low latitudes. J. Atmos. Terr. Phys., 1977; 39, 1079-1086.

- 210 11. Rigoti A., Chamalaun F. H., Trivedi N. B. and Padilha A. L., Characteristics of the equatorial
211 electrojet determined from an array of magnetometers in N-NE Brazil, Earth Planet Space,
212 1999; 51, 115-128.
213
214 12. Chandra H., Sinha H. S. S., and Rastogi R. G., Equatorial electrojet studies from rocket and
215 ground measurements, Earth Planets Space, 2000; 52,. 111–120.
216
217 13. Rastogi R. G., Some aspects of equatorial electrojet in South America, Indian J. Rad. and
218 Space Phys, 2007; 36, 315-317.
219