2

3

4

5

6

7 8

9

1

Natural Convective Mass Transfer MHD Flow of Chemically Reactive Micropolar Fluid past a Vertical Porous Plate

Khan Enaet Hossain^{1*}, S.M. Maruf Hassan ², Sayed Auntashire Rahman³ Md. Mohidul Haque ⁴

¹Mathematics Discipline, Khulna University, Khulna, Gmail: enayetkhan09@gmail.com ²Mathematics Discipline, Khulna University, Khulna, Gmail: math.091264@gmail.com ³Mathematics Discipline, Khulna University, Khulna, Gmail: sam_maruf07@yahoo.com ³Mathematics Discipline, Khulna University, Khulna, Gmail: mmhaque@math.ku.ac.bd

16 **ABSTRACT**

Magnetic field effects on a free convective mass transfer flow of chemically reactive micropolar fluid over a vertical porous plate are investigated in this work. Suitable similarity transformations are used to derive a set of nonlinear ordinary differential equations governing the flow. Analytical solution of the dimensionless problem is obtained using perturbation technique. The velocity, angular velocity and concentration profiles, which are controlled by a number of parameters, are presented graphically. Based on these graphs the conclusions are depicted, and the obtained results are tested for their accuracy.

17 18 19

Keywords: MHD, Micropolar fluid, Porous plate, Chemical reaction, Perturbation technique.

20 **1. INTRODUCTION**

21 In recent years, the theory of micropolar fluid has fascinated momentous attention among the researchers. Indeed, the theory of micropolar fluids is introduced by Eringen(1960) can be used to explicate the flow characteristics of polymers, 22 paints, animal blood, lubricants, liquid crystals, hematological suspensions, geomorphological sediments and 23 geomorphological sediments etc. The theory described about local effect increasing due to microstructure and the 24 intrinsic motion of the fluid elements. The model of micropolar fluid represents fluids consisting of rigid, randomly 25 generated particles suspended in a viscous medium where the deformation of the particles is neglected. Kucaba-Pietal 26 (2004), Khedr (2009) investigated colloidal, Muthu (2008) analyzed human and animal blood, Lockwood et al. (1987) 27 driven liquid crystal as well as exotic lubricants. Kelson and Desseaux (2001) investigated the effect of surface 28 circumstance on micropolar fluid flow. The unsteady micropolar fluid flow between two parallel porous plates was driven 29 Srinivasacharya et al. (2001). Mixed convection micropolar fluid on a porous stretching sheet is investigated by Bhargava 30 (2003). Tripathy et al (2016) has investigated hydromagnetic micropolar fluid on a porous stretching sheet. Mohanty et al. 31 (2015) investigated heat and mass transfer effect on micropolar fluid on a porous stretching sheet. The mathematical 32 33 theory of equations of micropolar fluids and applications of these fluids in the theory of lubrication and in the theory of 34 porous media has been studied by Lukaszewicz (1999).

Fluid dynamics is the area of science which is related with the study of motion of fluids. Magnetohydrodynamics (MHD) is the study of the flow of electrically conducting fluids in the presence of a magnetic field. The study of Magnetohydrodynamics has numerous applications such as induction flow meter, cooling of nuclear reactors, structure of stars and planets. Recently, the application of Magnetohydrodynamics in the polymer industry and metallurgy has fascinated the attention of many scholars.

Free convection flow takes place frequently in nature, flows of fluid through porous media are of significant interest now days and have fascinated by many researchers owing to their applications in the science and technology. Study of fluid flow in porous medium is based upon the empirically Darcy's law. Such flows are assumed to be useful in diminishing the free convection.

Nomenclature			
и, v	Velocity component	D_m	Diffusion coefficient
8	Acceleration	C _r	Chemical reaction parameter
β^*	concentration expansion coefficient	C_{∞}	Concentration of uniform flow
B_0	Magnetic field component	Δ	Micro-rotational number
χ	Vortex viscosity	G_m	Modified Grashof number
γ	Spin gradient viscosity	K	Permeability of porous plate
υ	Viscosity	M	Magnetic force number
ρ	Density	^	Spin gradient viscosity number
S _c	Schmidt number	λ	Vortex viscosity

Along with the free convection flow, the phenomenon of mass transfer is also very significant in the theories of stellar
structure. The study of convective flow with mass transfer along a vertical porous plate is receiving paramount interest of
many scholars because of its numerous applications in the field of geophysical and cosmical sciences. Prathapkumer
(2010) studied on free convection flow of micropolar and viscous fluids through a vertical duct. Raptis (2011),
Samiulhaqet al. (2012) and Seth et al. (2015) studied on free convective oscillatory flow and mass transfer with ramped
temperature on a porous plate. Thereafter, Chamkha (2000), Chaudhary (2007), Samad and Mohebujjaman (2009),
Eldabe (2011) and Seth (2015) have paid attention to the study of MHD free convection and mass transfer flows.

52 The chemical reaction rate counts on the concentration of the species itself. In many chemical engineering systems, there 53 is the chemical reaction between a foreign mass and the fluid. These system occurs in many industrial applications such aspolymer production, manufacturing of ceramics and food processing. Mishra et al (2016) has analyzed the effect of 54 chemical reaction on hydromagnetic micropolar fluid flow. Raju et al. (2013) investigated an unsteady free convection and 55 chemically reactive MHD flow through infinite vertical porous plate. Bakr (2011) has driven the effect of chemical reaction 56 57 on micropolar fluid with oscillatory plate. Das et al. (1994) analyzed the effective of first order homogeneous chemical reaction of an unsteady micropolar fluid flow. Ibrahim et al. (2008), Anand Rao et al. (2012), Das (2012) and Raju et al. 58 (2013) investigated the effect of chemical reaction on an unsteady MHD free convection fluid through semi-infinite vertical 59 60 porous plate with heat absorption.Bakr (2011) analyzed the characteristic of a micropolar fluid velocity on oscillatory plate and constant heat source in a rotating frame. Kucaba-Pietal (2004), Khedret al. (2009) investigated the micro 61 62 inertia effects on the flow of a micropolar fluid past a semi-infinite plate.

Hence our main goal is to investigate a free convective mass transfer steady flow of a chemically reactive micropolar fluid past a semi-infinite porous plate.

65 2. FORMULATION OF THE PROBLEM

66 A natural convective mass transfer steady flow of a chemically reactive micropolar fluid through a semi-infinite vertical

porous plate is taken into account at the presence of magnetic field. The flow is considered vertically by x-direction and ydirection is represented horizontally. When the flow at rest, the species concentration level $C=C_{a}$ at all point, where C_{b} be

 c_{∞} direction is represented nonzontally. When the now at rest, the species concentration level $C=C_{\infty}$ at all point, where C_{∞} be

the concentration of uniform flow. It is also assumed that a magnetic field B of uniform strength is applied normal to the flow region. The physical configuration and co-ordinate system of the problem is presented in the following Fig.1.

44



Fig. 1. Physical configuration of the flow

Within the framework of the above stated assumption, the governing equations under the boundary-layer approximationsare given by,

74 Continuity Equation
$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} = 0$$
 (1)

Momentum Equation

71

76
$$u\frac{\partial u}{\partial x} + v\frac{\partial u}{\partial y} = g\beta^* \left(C - C_{\infty}\right) + \left(v + \frac{\chi}{\rho}\right)\frac{\partial^2 u}{\partial y^2} + \frac{\chi}{\rho}\frac{\partial\Gamma}{\partial y} - \frac{\sigma' uB_0^2}{\rho} - \frac{v}{K_1}u$$
(2)

Angular Momentum Equation

78
$$u\frac{\partial\Gamma}{\partial x} + v\frac{\partial\Gamma}{\partial y} = \frac{\gamma}{\rho j} \left(\frac{\partial^2\Gamma}{\partial y^2}\right) - \frac{\chi}{\rho j} \left(2\Gamma + \frac{\partial u}{\partial y}\right)$$
(3)

Concentration Equation

80
$$u\frac{\partial C}{\partial x} + v\frac{\partial C}{\partial y} = D_m \frac{\partial^2 C}{\partial y^2} - K_c (C - C_\infty)$$
(4)

81 with boundary condition,

82
$$u = 0, \ \Gamma = -s \frac{\partial u}{\partial y}, \ C = C_w \text{ at } y = 0$$

83
$$u = 0, \Gamma = 0, C = C_{\infty} \text{ at } y \to \infty$$

where *u* is the velocity component, Γ is the velocity acting in *z* - direction (the rotation of Γ is in the *x*-*y* plane), *B*₀ is the magnetic field component, *g* is local acceleration due to gravity, χ is the vortex viscosity, γ is the spin gradient viscosity, β^* is concentration expansion coefficient.

87 2.1 Non-dimensional Form

88 Since our goal is to attain analytical solutions of the problem so we introduce the following dimensionless variables,

89
$$\eta = y \sqrt{\frac{U_0}{2\upsilon x}}, \psi = \sqrt{2\upsilon U_0 x} f(\eta), u = \frac{\partial \psi}{\partial y} \text{ and } v = -\frac{\partial \psi}{\partial x}, \phi(\eta) = \frac{C - C_{\infty}}{C_w - C_{\infty}} \text{ and } \Gamma = \sqrt{\frac{U_0^3}{2\upsilon x}} g(\eta)$$

90 The dimensionless equations are,

91
$$(1+\Delta)f'''(\eta) + \Delta g'(\eta) + f(\eta)f''(\eta) + G_M \varphi(\eta) - (K+M)f'(\eta) = 0$$
 (5)

92
$$\wedge g''(\eta) + f'(\eta)g(\eta) + f(\eta)g'(\eta) - 2\lambda g(\eta) - \lambda f''(\eta) = 0$$
 (6)

93
$$\varphi''(\eta) + S_c f(\eta) \varphi'(\eta) - S_c C_r \varphi(\eta) = 0$$
 (7)

94 The associate boundary conditions

95
$$f(\eta) = f_w, f'(\eta) = 0, g(\eta) = -sf''(\eta), \varphi(\eta) = 1 \eta = 0$$

96
$$f'(\eta) = 0, g(\eta) = 0, \varphi(\eta) = 0, \eta \to \infty$$

97 Where, micro-rotational number $\Delta = \frac{\chi}{\rho v}$, modified Grashof number, $G_m = \frac{g\beta^* (C_w - C_{\infty})2x}{U_0^2}$ permeability of porous

98 plate, $K = \frac{2\nu x}{K_1 U_0}$, magnetic force number, $M = \frac{\sigma B_0^2 2x}{U_0 \rho}$, Spin Gradient number, $\Lambda = \frac{\gamma}{\nu \rho j}$, Vortex viscosity,

99
$$\lambda = \frac{2x\chi}{\rho j U_0}$$
, Schmidt number, $S_c = \frac{v}{D_m}$, chemical reaction parameter, $C_r = K_c \frac{2x}{U_0}$
100

101 3. ANALITICAL SOLUTION

102 Since the solution is sought for the large suction further transformation can be made Arifuzzaman (2015) as,

103 $\xi = \eta f_w$

1

104
$$f(\eta) = f_w F(\xi)$$
(8)

(7)

105
$$\phi(\eta) = f_w^2 G(\xi)$$
 (9)

06
$$g(\eta) = f_w^{3} H(\xi)$$
 (10)

107 Now the model with small quantity,

108
$$(1+\Delta)F'''+\Delta H'+FF''+\varepsilon G_M G-(K+M)\varepsilon F'=0$$
 (11)

$$109 \qquad \wedge H'' + F'H + FH' - 2\lambda \varepsilon H - \lambda \varepsilon F'' = 0 \tag{12}$$

$$110 \qquad G'' + S_c G' F - \varepsilon S_c C_r G = 0 \tag{13}$$

111 The associate boundary conditions

112
$$F(\xi) = 1$$
, $F'(\xi) = 0$, $G(\xi) = \varepsilon$, $H(\xi) = -sF''(\xi)$ at $\xi = 0$
113 $F'(\xi) = 0$, $G(\xi) = 0$, $H(\xi) = 0$ at $\xi \to \infty$

113 $F'(\xi) = 0, \ G(\xi) = 0, \ H(\xi) = 0 \text{ at } \xi \to \infty$

114 Now for the large suction $(f_w > 1)$, \mathcal{E} will be very small. Therefore following **Bestman** (1990), F, G and H can be 115 expended in terms of the small perturbation quantity ε ,

116
$$F(\xi) = 1 + \varepsilon F_1(\xi) + \varepsilon^2 F_2(\xi) + \varepsilon^3 F_3(\xi) + \dots$$
 (14)

117
$$G(\xi) = \varepsilon G_1(\xi) + \varepsilon^2 G_2(\xi) + \varepsilon^3 G_3(\xi) + \dots$$
(15)

118
$$H(\xi) = \varepsilon H_1(\xi) + \varepsilon^2 H_2(\xi) + \varepsilon^3 H_3(\xi) + \dots$$
 (16)

The dimensionless equations (5) - (7) transform into the following first order, second order and third order equations with
 their associated boundary conditions,

$$\begin{array}{c} (1+\Delta) F_{1}^{\prime\prime\prime} + \Delta H_{1}^{\prime} + F_{1}^{\prime\prime} = 0 \\ 121 \qquad \wedge H_{1}^{\prime\prime} + H_{1}^{\prime} = 0 \\ G_{1}^{\prime\prime} + S_{c} G_{1}^{\prime\prime} = 0 \end{array} \right\}$$
(17)

122

123 $F_1 = 0, \ G_1 = 1, \ H_1 = 0 \ \text{at} \ \xi = 0$

124
$$F_1' = 0, \ G_1 = 0, \ H_1 = 0 \text{ at } \xi \to \infty$$

125

$$(1+\Delta) F_{2}''' + \Delta H_{2}'(\xi) + F_{1}F_{1}'' + F_{2}'' + G_{M}G_{1}(\xi) - (K+M)F_{1}' = 0 126 \qquad \wedge H_{2}'' + F_{1}'H_{1} + F_{1}H_{1}' + H_{2}' - 2\lambda H_{1} - \lambda F_{1}'' = 0 G_{2}'' + S_{c}G_{2}' + S_{c}G_{1}'F_{1} - S_{c}C_{r}G_{1} = 0$$

$$(18)$$

127 128

 $F_2 = 0, \qquad G_2 = 0, \quad H_2 = 0 \text{ at } \xi = 0$

For the purpose of the applicability of the present mathematical model, the analytical solution are driven using the perturbation method and the discussion is made for various values of parameters just as Modified grashof number G_m ,Suction parameter f_w , Magnetic force number M, permeability of porous plate K, Micro-rotational number Δ , Vortex viscosity λ , Spin gradient viscosity number (\wedge), Schmidt number S_c and Chemical reaction parameter C_r . The fluid velocity, angular velocity and concentration versus the non-dimensional coordinate variable η are displayed in Figures.

The velocity profiles are illustrated in Fig. 2. The increase values of magnetic parameter create a drag force known as 160 Lorentz force. As it is observed, the velocity profiles curve climb up at the increase of magnetic force number. Modified 161 Grashof number signifies the effect of buoyancy force to the viscous hydrodynamic force. The effect of buoyancy force, 162 velocity curves increase with modified Grashof number (G_m). It is observed that, the effect of permeability parameter (k) at 163 the present of porous medium causes higher restriction to the flow. Due to increase of permeability parameter, the velocity 164 165 profiles decrease which agrees with Krishnamurthy et al (2016). Afterwards, the suction parameter (f_{w}) stabilizes the boundary layer growth. So the velocities profiles curve decline with go up suction parameters. Schmidt number decreases 166 167 the molecular diffusivity. Then with increase of vortex viscosity, the velocity profiles plunge.

The effect of different physical parameters on angular velocity is demonstrated in Fig. 3. The microrotational effect distinguishes the micropolar fluid. It is notice that, the Modified Grashof number and micro-rotational number enhance angular velocity profiles. The effect of Schmidt number is not remarkable but suction parameter, spian gradient number, vortex viscosity decrease angular velocity profiles. The presence of porous matrix, vortex viscosity decrease angular velocity which agrees with Mishra *et al* (2016).

Fig. 4 describes the concentration profiles. It is observed that concentration profiles affected by chemical reaction which is good agreement with the previous result of Krishnamurthy *et al* (2016), Tripathy *et al* (2016). The chemical reaction parameter declines the concentration of the spices in the boundary layer. The concentration profile decreases for Schmidt number which agrees with Mishra *et al* (2016). Concentration profiles calculated for higher values of Suction parameter where concentration is more decrease for $f_w = 29.0$ rather than $f_w = 25.0$. The effect of micro-rotational parameter on concentration profile is not remarkable.







(c)
 Fig. 2. Velocity profiles for different values of (a) modified Grashof number (b) suction parameter (c) magnetic force number (d) permeability of porous plate (e) spin gradient viscosity number.



(e)



Fig. 3. Angular velocity profiles for different values of (a) modified Grashof number (b) Schmidt number (c) suction parameter (d) micro-rotational number (e) spin gradient viscosity number (f) vortex viscosity.



Fig. 4. Concentration profiles for different values of (a) suction parameter (d) Schmidt number (c) microrotational number (d) chemical reaction parameter.

181 4. CONCLUSIONS

Some of the important findings of the present work obtained from the graphical representation of the results are listed 182 183 below:

- 1. The fluid velocity and angular velocity profile decreases with the increase of Modified Grashof number. 184
- 2. The velocity and angular velocity profile decreases with the increase of Suction parameter and also the 185 186 concentration profile decreases with the increase of Suction parameter.
- 187 The velocity profile decreases and angular velocity profiles decreases with the increase of Schmidt number and also the concentration profile decreases with the increase of Schmidt number. 188
- 4. The velocity profile increase with Magnetic force number. 189
- 5. The velocity profiles decreases with the increase of Permeability of porous plate. 190
- 191 6. The concentration profile decreases with the increase of Chemical reaction parameter.
- 7. The angular velocity profile increases with the increase of Micro-rotational number and the concentration profile 192 decreases with the increase of Micro-rotational number. 193
- 8. The angular velocity profile decreases with the increase of Spin gradient viscosity number. 194
- 9. The angular velocity profile decreases with the increase of Vortex viscosity. 195

REFERENCES 197

196

201 202

203

204

213

214

215

216

219

220

226

228 229

230

- 1. Eringen AC. Theory of micropolar fluids, Journal of Applied Mathematics and Mechanics. 1960; 6: 1–18. 198
- Kucaba-Pietal A. Microchannels flow modelling with the micropolar fluid theory. Bulletin of the Polish Academy of 199 2. 200 Science, 2004: 52(3): 209-214.
 - Khedr MEM, Chamkha AJ, Bayomi M. MHD Flow of a Micropolar Fluid past a Stretched Permeable Surface with 3. Heat Generation or Absorption. Nonlinear Analysis: Modelling and Control. 2009; 14(1): 27-40.
 - 4. Muthu P, Rathish Kumar V and Peeyush Chandra. A study of micropolar fluid in an annular tube with application to blood flow. Journal of Mechanics in Medicine and Biology. 2008; 8(4): 561-576.
- 205 Lockwood F. Benchaitra M and Friberg S. ASLE Tribology Transactions. Astrophys Space, 1987; 30: 539. 5.
- 206 6. Tripathy RS, Dash GC, Mishra SR, Mohammad Mainul Hoque, Numerical analysis of hydromagnetic micropolar fluid along a stretching sheet embedded in porous medium with non-uniform heat source and chemical reaction, 207 208 Engineering Science and Technology, 2016; 19: 1573–1581.
- 209 Lukaszewicz G., Micropolar Fluids: Theory and Applications. Basel, Switzerland: Birkhhauser, 1999. 7.
- 210 Mishra SR., S. Baag, DK. Mohapatra, Chemical reaction and Soret effects on hydromagnetic micropolar fluid 8. 211 along a stretching sheet, Engineering Science and Technology, 2016. 212
 - Krishnamurithy MR., Prasannakumara BC., Gireesha BJ., Rama Subba Reddy Golar, Effect of chemical reaction 9. on MHD boundary layer flow and melting heat transfer of Williamson nanofluid in porous medium. Engineering Science and Technology, an International Journal, 2016; 19: 53–61.
 - Prathap Kumar J, Umavathi JC, Chamkha J, Pop I. Fully developed free convective flow of micropolar and 10. viscous fluids in a vertical channel. Applied Mathematical Modelling, 2010; 34: 1175-1186.
- 217 Raptis A. Free convective oscillatory flow and mass transfer past a porous plate in the presence of radiation for 11. 218 an optically thin fluid. Thermal Science, 2011; 15: 849-857.
 - Samiulhag I, Khan F, Ali and Shafie S. MHD free convection flow in a porous medium with thermal diffusion and 12. ramped wall temperature. Journal of the Physical Society of Japan, 2012; 81: 044401.
- 221 Seth GS, Sharma R and Sarkar S. Natural convection heat and mass transfer flow with Hall current, rotation, 13. radiation and heat absorption past an accelerated moving vertical plate with ramped temperature. Journal of 222 Applied Fluid Mechanics, 2015; 8: 7-20. 223
- 224 14. Chamkha AJ and Khaled ARA. Hydromagnetic combined heat and mass transfer by natural convection from a 225 permeable surface embedded in a fluid saturated porous medium. International Journal Numerical Methods Heat Fluid Flow, 2000; 10: 455-477. 227
 - Kelson NA, Desseaux A. Effect of surface condition on flow of micropolar fluid driven by a porous stretching 15. sheet. International Journal of Engineering Science. 2001: 39: 1881–1897.
 - 16. Chaudhary RC. and Jain A. Combined heat and mass transfer effects on MHD free convection flow past an oscillating plate embedded in porous medium. Romanian Journal of Physics, 2007; 52: 505-524.
- Samad MA, Mohebujjaman M. MHD heat and mass transfer free convection flow along a vertical stretching sheet 231 17. 232 in presence of magnetic field with heat generation. Research Journal of Applied Sciences, Engineering and 233 Technology, 2009; 1(3): 98-106.

- 18. Eldabe NTM, Elbashbeshy EMA, Hasanin WSA and Elsaid EA. Unsteady motion of MHD viscous incompressible
 fluid with heat and mass transfer through porous medium near a moving vertical plate. International Journal of
 Energy Technology and policy, 2011; 3: 1–11.
- 19. Das UN, Dekha R and Soungalgekar VM. Effects on mass transfer on flow past an impulsively started infinite
 vertical plate with constant heat flux and chemical reaction. Forschungim Ingenieurwesen. 1994; 60: 284-287.
- 20. Ibrahim FS, Elaiw AM and Bakr AA. Effect of the chemical reaction and radiation absorption on the unsteady
 MHD free convection flow past a semi-infinite vertical permeable moving plate with heat source and suction.
 Communication Nonlinear Science and Numerical Simulation. 2008; 13: 1056–1066.
- 242 21. Annad Rao J, Sivaiah, S. and Srinivasa Raju, R. "Chemical Reaction Effects on an Unsteady MHD Free
 243 Convection Fluid Flow past a Semi-infinite Vertical plate Embedded in a Porous Medium with Heat Absorption."
 244 Journal of Applied Fluid Mechanics. 2012; 3: 63-70.
- 245 22. Das K. Magnetohydrodynamics free convection flow of a radiating and chemically reacting fluid past an
 246 impulsively moving plate with ramped wall temperature. Journal of Applied Mechanic. 2012; 79: 17-25.
- 247 23. Raju MC, Varma SVK, Rao RRK. Unsteady MHD free convection and chemically reactive flow past an infinite
 248 vertical porous plate. Journal of Future Engineering and Technology. 2013; 8(3): 35-40.
- 249
 24. Bakr AA. Effects of chemical reaction on MHD free convection and mass transfer flow of a micropolar fluid with
 250 oscillatory plate velocity and constant heat source in a rotating frame of reference. Communication Nonlinear
 251 Science and Numerical Simulation. 2011; 16: 698–710.
- 252 25. Srinivasacharya D, Ramana murthy JV, Venugopalam D. Unsteady stokes flow of micropolar fluid between two 253 parallel porous plates. International Journal of Engineering Science. 2001; 39:1557–1563.
- Bhargava R, Kumar L, Takhar HS. Finite element solution of mixed convection microploar fluid driven by a porous stretching sheet. International Journal of Engineering Science. 2003; 41: 2161–2178.
- 256 27. Mohanty B, Mishra SR, Pattanayak HB. Numerical investigation on heat and mass transfer effect of micropolar
 257 fluid over a stretching sheet through porous media. Alexandria Engineering Journal. 2015; 54: 223–232.
- 258 28. Ahmed N. and Das KK. Unsteady MHD mass transfer flow past a suddenly moving vertical plate in a porous 259 medium in rotating system. International Journal of Engineering Science and Technology. 2013; 5: 1906-1923.
- Arifuzzaman SM. Manjirul Islam, Mohidul Haque. Combined Heat and Mass Transfer Steady Flow of Viscous
 Fluid over a Vertical Plate with Large Suction. International Journal of Engineering Science, Technology and
 society. 2015; 3(5): 236-242.

APPENDIX

264 265

266
$$A_1 = \frac{1}{\Lambda}, \qquad A_2 = \frac{1}{1+\Delta}, \qquad A_3 = \frac{-A_2 G_M}{\left(-S_c^3 + A_2 S_c^2\right)}, \qquad A_5 = \frac{S_c}{A_2 \varepsilon^2}, \qquad A_6 = \frac{S_c}{\varepsilon^2} \qquad A_7 = \frac{S_c^2}{A_2 \varepsilon^2 \left(A_2^2 + S_c A_2\right)},$$

267
$$A_{20} = \frac{A_3}{2}, \qquad A_8 = \frac{-\frac{A_2N}{\varepsilon^2}}{\left(A_2^2 - A_1A_2\right)}, \qquad A_9 = \frac{A_3\lambda S_c^2}{S_c^2 - A_1S_c}, \qquad A_{10} = \Delta A_2^2 A_8 \qquad A_{11} = \Delta A_2 A_9 S_c, \qquad A_{12} = G_m C_r A_2,$$

268
$$A_{13} = \frac{A_2(K+M)}{\varepsilon^2}, \quad A_{14} = A_2 S_c A_3(K+M), \quad A_{15} = \frac{A_{10}}{A_2^2}, \quad A_{16} = \frac{A_{11}}{\left(-S_c^3 + S_c^2 A_2\right)}, \quad A_{17} = \frac{A_{12}}{\left(-S_c^3 + S_c^2 A_2\right)},$$

269
$$A_{18} = \frac{A_{13}}{A_2^2}, \quad A_{19} = \frac{A_{14}}{\left(-S_c^3 + S_c^2 A_2\right)}$$

A J