

**STRUCTURE OF THE ATOMIC NUCLEI IN THE UNIVERSE MODEL WITH  
MINIMAL INITIAL ENTROPY**

**Abstract**

In this paper we show on the base of new ideas about the origin and evolution of the Universe, that in three-dimensional space the fundamental particles should have electric charges equal to 0,  $\pm e$ ,  $\pm 2e$ ,  $\pm 3e$ , what corresponds to the neutron and three pairs of light stable nuclei (hydrogen, helium, lithium). All heavy ( $Z \geq 4$ ) cores are presented in the form of molecular structures consisting of light nuclei; there are shown the reasons of instability of the nuclei in the ground and excited states. The hierarchy of bosons which are responsible for the interaction between particles in different hierarchical layers of fiber space Super-Universe is given.

**Keywords:** heavy cores, light stable nuclei, excited states, hierarchy of bosons, three-dimensional space.

Pacs 23.20.Nx; 24.10.-i; 98.80 k

The process of our Universe origin as the part of the Super-Universe was described in detail in the article [1] on the basis of the Law of similarity [2] and the Law of unity.

Our Universe is a part of the Super-Universe. At the same time, the Super-Universe is layered space [3] where adjacent layers change the space dimensionality by one. Usual for us three-dimensional space (four-dimensional (3 + 1) Universe) has a border with two-dimensional space of quarks [4]. Two-dimensional space has a border with one-dimensional space of diones, that are found to be Planck particles. Finally, one-dimensional space has a border with zero-dimensional space of scalar Field-time. There is an information interaction among adjacent spaces through a single delocalized point. A zero-dimensional space of Field-time can interact with all other spaces and can dictate the program of the Universe evolution.

Field brings energy subsequently first into one-dimensional, then into two-dimensional and finally, after  $\Delta t = 3 \cdot 10^{-5}$  s, into three-dimensional space. As a result in three-dimensional space cold neutron energy is created, which has initial density similar to nuclear density. Neutrons decay leads to protons and electrons creation in equal quantity, making Universe electrically neutral.

In this paper, the mechanisms of particles and atomic nuclei creation in our four-dimensional (3 + 1) Universe will be revealed using the same Laws.

A lot of models of atomic nucleus structure have been discussed in the literature. One of these models included in all textbooks on nuclear physics, presents the nucleus as the set of protons and neutrons with the configuration which provides the minimum energy of nucleus. It is assumed that, despite of the  $\alpha$ -activity of heavy nuclei,  $\alpha$ -particles are absent in the nuclear structure as defined clusters. Among these models there is also cluster (molecular) model [2, 5-6].

Cluster model (or the model of nucleon associations) treats the structure of some nuclei as a kind of molecule consisting of  $\alpha$ -particles, deuterium (D), tritium (T), and others. For example,  $^{12}\text{C} = 3\alpha$ ,  $^{16}\text{O} = 4\alpha$ ,  $^6\text{Li} = \alpha + D$ ,  $^7\text{Li} = \alpha + T$  and so on.

Model of nucleon associations is a model of atomic nucleus based on nucleus representation as a system of clusters, or nucleon associations of a certain type, usually,  $\alpha$ -clusters. The simplest version of this model ( $\alpha$ -cluster model) was formulated in 1937 by J. A. Wheeler. This model has arisen from the fact that the stability of nuclei is increased if the



core has an even number of protons and neutrons, like in  $\alpha$ -particle. Therefore such nuclei were described as clusters of  $\alpha$ -particles. Among these nuclei there are  ${}^8\text{Be}$ ,  ${}^{12}\text{C}$ ,  ${}^{16}\text{O}$ ,  ${}^{20}\text{Ne}$  and similar nuclei (at  $n = 2, 3, 4, 5$ ). For these nuclei an enormous amount of energy  $E_n$  is needed to remove a neutron. For nearest nucleus with odd number of neutrons this energy decreases by 10-15 MeV. Meanwhile the energy which is needed to remove an  $\alpha$ -particle ( $E_\alpha$ ) is rather small.  ${}^8\text{Be}$  nucleus is unstable as for the decay into two  $\alpha$ -particles ( $E_\alpha < 0$ ), and as a result this nucleus does not exist. For other nuclei of this row the binding energy of the  $\alpha$ -particles increases (in a nucleus of  ${}^{12}\text{C}$  the energy  $E_\alpha = 7$  MeV, in  ${}^{16}\text{O}$   $E_\alpha = 16$  MeV).

There was experimentally established the following law: nuclei consisting of  $\alpha$ -particles can easily emit them in nuclear reactions. Moreover, it has been shown that these nuclei have excited states with abnormally large width of  $\alpha$ -transitions. This means that  $\alpha$ -particles **exist** on nucleus surface as separate clusters.

For such nuclei, the nucleus wave function can be written as a product of the antisymmetrized wave functions  $\psi_\alpha$ , describing the internal motion of the nucleons in the individual  $\alpha$ -cluster, and the wave function  $\chi$ , describing the motion of the clusters with respect to each other.

$$\psi({}^8\text{Be}) = \hat{A} \psi_{\alpha 1}(r_1) \psi_{\alpha 2}(r_2) \chi_L(R_1 - R_2),$$

where  $R_i = \sum_{i=1}^4 r_i / 4$  is the radius-vector of the center of mass of the  $\alpha$ -cluster,  $L$  is total

the orbital angular momentum of the nucleus,  $\hat{A}$  is the antisymmetrization operator on the nucleons belonging to different clusters.

However, it has been found that such wave function can satisfactorily describe the behavior of  ${}^8\text{Be}$  and  ${}^{12}\text{C}$ , but it can not describe  ${}^{16}\text{O}$ ,  ${}^{20}\text{Ne}$ , etc.

The cluster model is used to describe the nuclear reactions. The most common approach here is the so-called resonating group method which is similar to the method of valence bonds for the description of the molecules [7].

The cluster model of heavy clusters is frequently used to describe nuclei. For example,  ${}^{24}\text{Mg}$  nucleus is described as a "molecule", consisting of two  ${}^{12}\text{C}$  nucleus separated in space. In this case, wave functions  $\psi_{{}^{12}\text{C}}$  instead of  $\psi_\alpha$  are written for nucleus.

It is interesting, that a quark model of nucleons is analogous to the cluster model of nucleus (nucleon is considered as a 3-quark cluster and it is also assumed the existence of multi-quark configurations: 6- and 9-quark clusters).

The cluster model proved to be useful for the description of a nucleon fragmentation processes in the nuclear reactions taking place under an action of high-energy heavy ions.

Thus, we have a confirmation of the molecular structure of nuclei. The only difference between cluster models used in experimental and theoretical studies from our model is that they are empirical, unproved. Our presentation naturally arises from the new methodological basis of the World cognition.

## Particles of four-dimensional Universe

According to the statements of Victor Kulish [2] our Manifested World has 4 dimensions and Hidden World has only 3 ones. Together we have 7 dimensions: 3 dimensions for the quarks and four dimensions for the nucleons, electrons, atoms, matter, fields.



Four-dimensional World of particles is produced by the three-dimensional World by quarks gluing by gluon into particles. These quarks are in the Hidden World, while corresponding particles are in the Manifest World. [1-2]

The charges of quarks are  $-(\frac{1}{3})e$  and  $+(\frac{2}{3})e$  (opposite signs for antiquarks),  $e$  is the minimum charge of the particle in four-dimensional space-time.

It follows that quarks charges are formed by the dimension of the World:  $\pm(\frac{1}{3})e$  for each coordinate. Thus, all types of quarks are two-dimensional (since the space has two dimensions, all the particles in this space should move only in two directions) which is allowed by the dimensionality of space. So, it can be assumed, that in Hidden space the charges 0,  $\pm(\frac{1}{3})e$  and  $\pm(\frac{2}{3})e$  can exist.

Comparing these conclusions with the data in Table 1 it can be concluded that for the quarks only charge  $-(\frac{1}{3})e$  and  $+(\frac{2}{3})e$  are realized, and opposite charges are for antiquarks. This result can be understood taking into account that the birth of the Universe is presented as a vortex (and as a result tightening in gravity [8] and time [9] takes place). At the same time, the 3 projections of charge are realized as stationary states in the World-3, see Figure 1.




Fig.1. Three projections of charge in the World-3. Two types of charge  $-(\frac{1}{3})e$  should be different by additional quantum numbers (helicity). Mirror reflection with respect to the vertical y-axis (or in inversion point) gives the charges of antiparticles.

To determine the charge in the World-4 it is necessary to use a sphere rotation (Fig. 2).




Fig.2. Four projections of a charge in the World 4. Mirror reflection in the plane xy (or in reversal point) will give the charges of antiparticles.

Table 1. Quarks.

Type (flavor) of quark	Electric charge	Spin	Colour	Mass (calculation)
<b>d</b>	$-(1/3)e$	$\hbar/2$	yellow, blue, red	$\sim 7 \text{ MeV}/c^2$



<b>u</b>	$+(2/3)e$		–"–	$\sim 5 \text{ MeV}/c^2$
<b>s</b>	$-(1/3)e$		–"–	$\sim 150 \text{ MeV}/c^2$
<b>c</b>	$+(2/3)e$		–"–	$\sim 1,5 \text{ GeV}/c^2$
<b>b</b>	$-(1/3)e$		–"–	$\sim 4,5 \text{ GeV}/c^2$
<b>t</b>	$+(2/3)e$		–"–	$\sim 175 \text{ GeV}/c^2$

It is necessary to note another important detail: all particles of World-4 have been formed due to the transfer of information from a quarks cluster while the heavy nuclei are formed from the particles of World-4 whose quarks do not have a border. This is worth also to remember during the consideration of the fusion reaction of helium nucleus formation from the nuclei of hydrogen and lithium or deuterium, when the quarks of complex core do not border each other. And only due to virtual pairs participation (a proton-antiproton, etc.) the  $\alpha$ -particle of World-4 is formed from a complex helium nucleus.

The dimensionality of the World is changing during the transition from the Hidden World to the Manifested World, and hence the magnitude of a charge. The dimensionality of the Manifested World requires a combination of quark charges to create a charge  $\pm e$ .

On the other hand, charges  $0, \pm Q/4, \pm 2Q/4, \pm 3Q/4$  should exist in the Manifested World. Here, the value of  $\pm Q$  corresponds to the charge of the next five-dimensional World where our space is generating (and probably hidden).

As a result,  $Q = 4q = 4e$  is an elementary charge of the next Manifested World where our particles will be quarks-4.

It is also should be noted, that according to Figure 2 stable charges  $\pm e, \pm 2e, \pm 3e$ , and  $0$  should exist in our space.

Nuclei of hydrogen correspond to the first particle (proton and deuteron), nuclei of helium correspond to the second particle ( ${}^3_2\text{He}$  and  ${}^4_2\text{He} = {}^4_2\alpha$ ), the third particle corresponds to lithium nuclei<sup>1</sup>. Of course, particles and antiparticles corresponding to particles with opposite charges should exist. However, the Manifested World has electrons with charge  $-e$  to stabilize atoms and to provide the electrical neutrality of the Universe.

Table 2. Particles of four-dimensional World (World-4)<sup>2</sup>.

Charge	Particles	The total content of isotopes in the Universe
$-e$	$E$	
$0$	${}_0^1n$	
$+e$	${}_1^1H, {}_1^2D$	0,65 [2,10]
$+2e$	${}_2^3He, {}_2^4He$	0,24 [2,10]
$+3e$	${}_3^6Li, {}_3^7Li$	$6,5 \cdot 10^{-5}$ [11]

<sup>1</sup> This suggests an interesting parallel: three pairs of quarks and three pairs of particles of World-4. For the second and third pairs of quarks the top quark is more massive, and for the first pair an opposite situation takes place. Similar relationships can be observed for propagation of particles in World-4.

<sup>2</sup> Using the presentation of the particles of the World-4, shown in Figure 2, we can find a product of the length of corresponding circle on the height of the segment of a circle, and then can find the ratio of these values. It turns out that it is equal to 1: 0.6285: 0.2484: 0; i.e. corresponds to the concentration of the relevant charges in the Universe.



Note: The concentrations of charged particles in the World-4 are following:

$$[{}^2_1D] = 1,56 \cdot 10^{-4} \cdot [{}^1_1H], \text{ the last one is 65\% of the mass of Universe,}$$

$$[{}^3_2He] = 1,38 \cdot 10^{-6} \cdot [{}^4_2He], \text{ the last one is 24\% of the mass of Universe,}$$

$$[{}^6_3Li] = 8,1 \cdot 10^{-2} \cdot [{}^7_3Li], \text{ the last one is } 2 \cdot 10^{-10} \text{ of the mass of Universe, } 6,5 \cdot 10^{-5} \text{ in the Earth.}$$

Since among the particles of the World-4 a rapid process of exchange interaction ( $p \leftrightarrow p$ ,  $n \leftrightarrow n$ ,  $n \leftrightarrow p$ ) takes place, it is necessary to assume that each element of these particles is a result of averaging, i.e. all the elements of particles are identical, and the particles of World-4 are indivisible. In such way they will perform for particles of brane in the World -5.

Since the particles of the World-4 act as indivisible, it is better to present them using a quark structure:

$${}^1_1H = 2u + d \equiv u^2d,$$

$${}^2_1D = 3u + 3d \equiv u^3d^3,$$

$${}^3_2He = 5u + 4d \equiv u^5d^4,$$

$${}^4_2He = 6u + 6d \equiv u^6d^6,$$

$${}^6_3Li = 9u + 9d \equiv u^9d^9,$$

$${}^7_3Li = 10u + 11d \equiv u^{10}d^{11}.$$

So, we have a stable structure containing of three quarks, 6 quarks, 9 quarks, 12 quarks, 18 quarks and 21 quarks. There are no structures containing of 15 quarks ( ${}^5_2He$  or  ${}^5_3Li$ ) in the World-4 (negative binding energy of proton or neutron with  ${}^4_2He$  [12]).

Thus, during the transition from the Hidden World-3 into the Manifested World-4 a formation of particles from quarks takes place, i.e. the real Manifested World. That is why quarks are in the Hidden World, and hadrons are in the Manifested World and there is an information interaction between them.

As other nucleus and atoms of our World are formed as a result of a combination of a family of particles of World-4, it should be assumed that with the formation of other nuclei and atoms the Manifest World-4 received the fifth coordinate (it becomes the brane of four-dimensional space), which began to increase in time, leading to the birth of matter, planets, stars, etc., causing the expansion of the Universe.

So, we are living in the swelling brane of the World-5.

Moving in the opposite direction, we shall understand that a generating two-dimensional World should exist for the hidden for us three-dimensional World providing the possibility of  $\pm 1/2$  charges of a quark-3. For the World-4 these charges are equal to  $\pm e/6$ . These quarks-2 will generate all possible quarks-3. It was shown in [1] that quarks-2 should be **diones** having both electric and magnetic charges. During the transition to the spaces of higher dimensionality magnetic charges cause the appearance of the spin of elementary particles.

A lot of quantum numbers of quarks are lost at the birth of the World-4, and in particular color. Therefore, we can assume that in the two-dimensional World there are some characteristics which are lost at the transition to the World-3 (including the abovementioned magnetic charges). Thus, two particles of World -2 have a wide set of quantum numbers, which are lost during the transitions to the Worlds of higher dimensionality.

The Fields-time coordinates are common for all spaces, so two spatial dimensions of quarks and one spatial dimension constituent quarks (diones) from the previous World should be added for our four-dimensional World. Totally it will be 7 dimensions. However, 3 of



them have various degrees of secrecy (2 for the nearest Hidden World and 1 for the remote one).

There are photons in the World-4. They appear, in particular, during particles-antiparticles annihilations. But there are particles (for example,  $\pi^0$ ), which have quark-antiquark type structure. This results in a disintegration of such particles into  $\gamma$ -quants in the World-4, while the quark-antiquark annihilation has to give 2 two-dimensional photons specific for World-3. Types of these photons are discussed in [1].

### Particles of World-5. The hierarchy of bosons.

In our World-5, all other nuclei ( $Z \geq 4$ ) are combinations of "elementary" particles of World-4 and can decay into these "basic" particles. These combinations are:

${}^3_1T \rightarrow {}^2_1D + {}^1_0n$ , - unstable ( $\beta^-$  - active) due to the contribution of the neutron nucleus;

${}^8_3Li \rightarrow {}^7_3Li + {}^1_0n$ , - the nucleus is  $\beta^-$  - active;

${}^9_4Be \rightarrow {}^7_3Li + {}^2_1D$ , - the nucleus is stable, but quite rare because there are not enough lithium and deuterium;

${}^{10}_4Be \rightarrow {}^7_3Li + {}^2_1D + {}^1_0n \leftrightarrow {}^7_3Li + {}^3_1T$ , - the nucleus is  $\beta^-$  - active;

${}^{10}_5B \rightarrow {}^4_2\alpha + {}^6_3Li$ , - the nucleus is stable; but less than  ${}^{11}_5B$ , because  $[{}^6_3Li] < [{}^7_3Li]$ ,

${}^{11}_5B \rightarrow {}^4_2\alpha + {}^7_3Li$ , - the nucleus is stable, but quite rare because there is not enough lithium;

${}^{12}_5B \rightarrow {}^4_2\alpha + {}^7_3Li + {}^1_0n$ , - nucleus is  $\beta^-$  - active,  ${}^{12}_6C$  is formed in an excited state, which decays into three  $\alpha$ -particles.

${}^{11}_6C \rightarrow 2 {}^4_2\alpha + {}^2_1D + {}^1_1H$ , or  ${}^{11}_6C \rightarrow 2 {}^4_2\alpha + {}^3_2He$ , the contribution second is small, the first one is  $\beta^+$  - active (the proton in the field of nuclear forces is unstable),

${}^{12}_6C \rightarrow 3 {}^4_2\alpha$ , or  ${}^{12}_6C \rightarrow 2 {}^6_3Li$ , - the nucleus is stable, but the probability of the

reaction of the second type is very small, because of the lack of  ${}^6_3Li$  in the nature;

${}^{13}_6C \rightarrow {}^6_3Li + {}^7_3Li$ , - nucleus is stable, but its abundance is very small (1%).

${}^{14}_6C \rightarrow 2 {}^7_3Li$ , or  ${}^{14}_6C \rightarrow {}^7_3Li + {}^4_2\alpha + {}^2_1D + {}^1_0n$ , or  ${}^{14}_6C \rightarrow 3 {}^4_2\alpha + 2 {}^1_0n$ , - unstable nucleus ( $\beta^-$ -active) as a result of the contribution of the neutron, due to big amount of  ${}^4_2\alpha$ ;

${}^{12}_7N \rightarrow 2 {}^4_2\alpha + {}^3_2He + {}^1_1H$ , - - the nucleus is  $\beta^+$ - active,  ${}^{12}_6C$  is formed in an excited state, with following decay into three  $\alpha$ -particles.

${}^{13}_7N \rightarrow 3 {}^4_2\alpha + {}^1_1H$ , - the nucleus is  $\beta^+$  - active,

${}^{14}_7N \rightarrow 3 {}^4_2\alpha + {}^2_1D$ , or  ${}^{14}_7N \rightarrow 2 {}^4_2\alpha + {}^6_3Li$ , - the nucleus is stable,

${}^{15}_7N \rightarrow 2 {}^4_2\alpha + {}^7_3Li$ , - the nucleus is stable, but its abundance is very small (0,365%),

${}^{16}_7N \rightarrow 2 {}^4_2\alpha + {}^7_3Li + {}^1_0n$ , - the nucleus is  $\beta^-$  - active, it transforms into  ${}^{16}_8O$  in the excited state, which emits one  $\alpha$ -particle,

${}^{16}_8O \rightarrow 4 {}^4_2\alpha$ , or  ${}^{16}_8O \rightarrow 2 {}^6_3Li + {}^4_2\alpha$ , - the nucleus is stable,<sup>3</sup>; the second contribution is small, due to the lack of  ${}^6_3Li$ .

${}^{17}_8O \rightarrow {}^4_2\alpha + {}^7_3Li + {}^6_3Li$ , - the quantity of such nuclei is small, since

$N({}^4_2\alpha) \gg N({}^7_3Li) \gg N({}^6_3Li)$ .

<sup>3</sup> Below much more structures for the core  ${}^{16}_8O$  providing its stability in the ground state will be presented.



222  $^{18}_8O \rightarrow ^4_2\alpha + 2\ ^7_3Li$ , - the nucleus is stable; but the quantity of these nuclei is less than of  
223  $^{16}_8O$ , because  $N(^4_2\alpha) \gg N(^7_3Li)$ , but is bigger than of  $^{17}_8O$  in 6 times.  
224  $^{19}_8O \rightarrow ^4_2\alpha + 2\ ^7_3Li + ^1_0n$ , - the nucleus is  $\beta^-$  - active,  
225  $^{18}_9F \rightarrow 4\ ^4_2\alpha + ^2_1D$ , or  $^{18}_9F \rightarrow 3\ ^4_2\alpha + ^6_3Li$ ,  $^{18}_9F \rightarrow 2\ ^4_2\alpha + ^7_3Li + ^2_1D + ^1_1H$ , - only the later  
226 combination provides  $\beta^+$ - activity, so the reaction is slow (109,7 min),  
227  $^{19}_9F \rightarrow 3\ ^4_2\alpha + ^7_3Li$ , - the nucleus is stable,  
228  $^{20}_9F \rightarrow 3\ ^4_2\alpha + ^7_3Li + ^1_0n$ , - the nucleus is  $\beta^-$  - active (11,56 s),  
229  $^{19}_{10}Ne \rightarrow 4\ ^4_2\alpha + ^2_1D + ^1_1H$ ,  $^{19}_{10}Ne \rightarrow 4\ ^4_2\alpha + ^3_2He$ , - the nucleus is  $\beta^+$  - active, small contribution  
230 of the second combination,  
231  $^{20}_{10}Ne \rightarrow 5\ ^4_2\alpha$ , - the nucleus is stable <sup>4</sup> (90,92%),  
232  $^{21}_{10}Ne \rightarrow 3\ ^4_2\alpha + ^7_3Li + ^2_1D$ , - the nucleus is stable, but the amount of these nuclei is small  
233 (0,257%)  
234  $^{22}_{10}Ne \rightarrow 2\ ^4_2\alpha + 2\ ^7_3Li$ , - the nucleus is stable (8,82%),  
235  $^{23}_{10}Ne \rightarrow 2\ ^4_2\alpha + 2\ ^7_3Li + ^1_0n$ , - the nucleus is  $\beta^-$  - active.  
236  $^{22}_{11}Na \rightarrow 5\ ^4_2\alpha + ^2_1D$ ,  $4\ ^4_2\alpha + ^6_3Li$ ,  $3\ ^4_2\alpha + ^7_3Li + ^3_2He$ ,  $^4_2\alpha + 2\ ^7_3Li + ^3_2He + ^1_1H$ ,  $2\ ^7_3Li + ^6_3Li + 2\ ^1_1H$ ,  
237 the nucleus is  $\beta^+$  - active taking into account last configurations,  
238  $^{23}_{11}Na \rightarrow 4\ ^4_2\alpha + ^7_3Li$ , the nucleus is stable,  
239  $^{24}_{11}Na \rightarrow 4\ ^4_2\alpha + ^7_3Li + ^1_0n$ , the nucleus is  $\beta^-$  - active.  
240 \* \* \* \* \*  
241  $^{55}_{25}Mn \rightarrow 5\ ^4_2\alpha + 5\ ^7_3Li$ ,  
242  $^{54}_{26}Fe \rightarrow 10\ ^4_2\alpha + 2\ ^7_3Li$ ,  $7\ ^4_2\alpha + 2\ ^7_3Li + 2\ ^6_3Li$ , the nucleus is stable (5,84%), small  
243 contribution of the second combination,  
244  $^{55}_{26}Fe \rightarrow 10\ ^4_2\alpha + 2\ ^7_3Li + ^1_0n$ ,  $7\ ^4_2\alpha + 3\ ^7_3Li + ^6_3Li$ ,  $6\ ^4_2\alpha + 4\ ^7_3Li + ^3_2He$ , in this case, the  
245 experiment shows *K*-electron capture with a conversion into a stable nucleus  $^{55}_{25}Mn$ . Thus, it  
246 is necessary to assume that the contribution of the last configuration is a main one, while the  
247 first is very small. In the field of nucleus  $^3_2He$  there is a reduction of the number of neutrons,  
248 which results in *K*-electron capture with a conversion it into tritium nucleus and its  
249 combination with  $\alpha$ -particle gives  $^7_3Li$ , nucleus, corresponding to the  $^{55}_{25}Mn$  nuclei  
250 configuration.  
251 Proceeding in the same manner to the heavy nuclei, we draw an attention to the fact that  
252 the protons-neutrons number relation for nuclei with number up to No. 50 can be described  
253 by combinations of  $^7_3Li$ ,  $^4_2He$  etc. But after this number the contribution of neutrons  
254 increases. Moreover, at the transition from  $^{208}_{82}Pb$  to  $^{238}_{92}U$  10 protons and 20 neutrons should  
255 be added [13]. So,  $^3_1T$ ,  $^6_2He$ ,  $^9_3Li$  should be included into the consideration. Such nuclei really  
256 exist, but, they are  $\beta^-$  - active with the lifetime  $3.87 \cdot 10^8$  s = 12.262 years, 0.797 s and 0.176 s  
257 respectively.  
258 Neutrons in a free state are also  $\beta^-$  - active, but, all nuclei are containing them. The  
259 interaction between nucleons much faster makes a transformation of a neutron into a proton,  
260 than it would decay.

<sup>4</sup> In fact, this nucleus has much more structures, how it will be shown below for  $^{16}_8O$ .



So, these three heavy nuclei can stably exist in nuclei, where the number of neutrons is twice higher than the number of protons. The need for such nuclei should be grounded on intranuclear interaction.

The bosons are always responsible for the interaction between particles. The gluons are responsible for strong interaction between quarks; the bosons which are partly in the World-3, and partly in the World-4 are responsible for weak interaction  $W(\pm)$  и  $Z^0$  [14].  $\pi(\pm)$  and  $\pi^0$  bosons can not be neglected in the consideration. They are responsible for the transfer of the interaction between nucleons in three groups of particles in the four-dimensional World. Bosons of the World-4 should provide the interaction between the particles of the World-4.  $\alpha$ -particle and boson, consisting of two coupled neutron  $Y(2n)$  can play this role. For example:






Fig. 3. Intranuclear interactions due to  $Y(2n)$  boson transfer.



In this case, it becomes clear a presence of doubled number of neutrons (in comparison with protons) in heavy nuclei.

Since it is considered that boson  $X(\alpha)$  is much heavier than boson  $Y(2n)$ , it should provide a much stronger interaction. However, a reality shows that  $\alpha$ -particles are poorly connected to the rest of the nucleus fragments, because they have a large electrical charge. As a result,  $\alpha$ -particle is not able to provide interaction between components of a nucleus. Moreover, if  $\alpha$ -particle participate in the formation of nuclei with  $Z > 50$ , the protons-neutrons number relation 1:2 can not be satisfied. So the interaction via  $X(\alpha)$  bosons has to be excluded from the consideration.

So, it is clear now that  ${}^8_4\text{Be} \rightarrow 2 {}^4_2\alpha$  can not exist and should immediately decay into two  $\alpha$ -particles. In the present case, it is impossible to arrange the transfer of two neutrons. A resonant exchange by  $Y(2n)$ -bosons is only possible. But, in this situation it is necessary to take off two neutrons from  $\alpha$ -particles, and then put two other neutrons on their place. If the last reaction seems simple enough, the first one requires a lot of effort and its implementation looks problematic.

In the case of  ${}^{12}_6\text{C} \rightarrow 3 \cdot {}^4_2\alpha$  the nucleus can be imagined only in excited state, which leads to its decay on 3  $\alpha$ -particles. The ground state can be provided by the configuration  ${}^{12}_6\text{C} \rightarrow {}^9_3\text{Li} + 3 {}^1_1\text{H} \leftrightarrow {}^7_3\text{Li} + 2 {}^1_1\text{H} + {}^3_1\text{T}$ . The "Molecular" structure is following (Figure 4):






Fig.4. The “Molecular structure” of carbon nucleus  $^{12}_6\text{C}$ .




Fig. 5. A particle surrounded by virtual bosons: *a* - proton, *b* -  $^9_3\text{Li}$ .

Since the boson barring the interaction is virtual, the particle can emit it and absorb it at once (Figure 5). This phenomenon is described in detail in quantum electrodynamics.

There is leak of information in the literature concerning bineutron, which is treated in this manuscript as boson of World-4. It is only known, that neutrons have huge interactions between each other by exchange of neutral pions. The same processes should take place in biproton. But in such case electrostatic repulsion between protons ( $\approx 1$  MeV) leads to resulting binding energy in biproton equal to -0.5 MeV. So, strong interaction energy, caused by transfer of neutral pion between neutrons, is equal to 0.5 MeV [15, 16]. But, neutron decays because of processes of weak interaction after  $\sim 881$  s [17]. After comparing half-decay periods of  $\beta^-$  - active nucleus (e.g.  $T_{1/2}(^{16}\text{N}) = 7.14$  s i  $T_{1/2}(^{18}\text{N}) = 0.63$  s;  $T_{1/2}(^{20}\text{F}) = 11.56$  s and  $T_{1/2}(^{22}\text{F}) = 4.0$  s [13]), we could make following conclusion: period of neutron half-decay could be decreased on 1-2 orders with increasing neutrons quantity in cluster. But this time is much longer than period of half-decay of strong interaction bosons (pions).

So, as a result of the processes of boson radiation-absorbing a spatial orientation or shape of nucleus components can continuously change. This is important in the cases when a nucleus of 5-dimensional World brane contains more than two particles-4. For example,  $^{12}_6\text{C}$



nucleus contains 4 particles-4 ( ${}^9_3\text{Li}+3{}_1^1\text{H}$  or  ${}^7_3\text{Li}+2{}_1^1\text{H}+{}_1^3\text{T}$ ). In this case, the transfer of Y(2n)-boson is equally probable for all three nucleus protons. Thus, after transfer Y(2n)-boson, the wave function of the nucleus will contain equal contributions from all three protons.

Similarly, for oxygen-16 nucleus:  ${}^{16}_8\text{O} \rightarrow 4{}_2^4\alpha$  - such state is a highly excited. The presence of the four  $\alpha$ -particles provides more opportunities for the organization of ground and lower excited states, the lowest excited state emitting only one  $\alpha$ -particle, turning into a carbon-12 nucleus.

$${}^{16}_8\text{O} \rightarrow {}^9_3\text{Li} + {}_2^4\alpha + 3{}_1^1\text{H} ,$$

$${}^{16}_8\text{O} \rightarrow {}^9_3\text{Li} + {}_1^3\text{T} + 4{}_1^1\text{H} ,$$

$${}^{16}_8\text{O} \rightarrow {}^7_3\text{Li} + 2{}_1^3\text{T} + 3{}_1^1\text{H} ,$$

$${}^{16}_8\text{O} \rightarrow {}^6_2\text{He} + {}^7_3\text{Li} + 3{}_1^1\text{H} ,$$

$${}^{16}_8\text{O} \rightarrow {}^6_2\text{He} + {}_2^4\alpha + {}_1^3\text{T} + 3{}_1^1\text{H} .$$

According to the **principle of similarity**, the nucleus has to be built as a set of three pairs of particles of World-4 like molecules are built of atoms. A virtual photon acts as the boson, which determines the interaction between electron and nucleus in atom [18]. At the same time, a *pair of electrons in singlet state, being surrounded by a coat of virtual photons, plays a role of boson, which defines the interaction of atoms in a molecule*. This pair of electrons is in continuous motion around the interacting atoms.

Similarly, **bineutron** (i.e. two neutrons) **in a coat of neutral pions** acts as a boson, which is glo for the interaction between particles of the World-4 in nuclei of chemical elements. Therefore it is logical to assume that complex nuclei have a certain geometric structure which is similar to structure of atoms in molecules. In this case, for  ${}^{16}_8\text{O} \rightarrow {}^9_3\text{Li} + {}_2^4\alpha + 3{}_1^1\text{H}$   ${}^9_3\text{Li}$  nucleus is surrounded along three sides by protons, and the interaction in this structure is due to Y(2n)-bosons. The interaction of this structure with a boson  ${}_2^4\alpha$  will be weakened, and as a result  $\alpha$ -particle will be emitted out of the nucleus, this is observed at the excitation of  ${}^{16}_8\text{O}$  nucleus.

The state  ${}^{16}_8\text{O} \rightarrow {}^6_2\text{He} + {}^7_3\text{Li} + 3{}_1^1\text{H}$  will be almost resonant with the previous state, if the both of them have the same geometric structure. However, in this state,  ${}_2^6\text{He}$  is assumed to be an active particle. Consequently, the structure may be different from the previous one and there are more variants of interaction via Y(2n) boson transfer. This can lead to corresponding reduction in the energy and stabilization of the nucleus.

The structure  ${}^{16}_8\text{O} \rightarrow {}^6_2\text{He} + {}_2^4\alpha + {}_1^3\text{T} + 3{}_1^1\text{H}$  where two transfer of Y(2n)-bosons takes place should have much lower energy. A little bit lower is situated the state corresponding to the structures  ${}^{16}_8\text{O} \rightarrow {}^9_3\text{Li} + {}_1^3\text{T} + 4{}_1^1\text{H}$  and its resonant (identity) state  ${}^{16}_8\text{O} \rightarrow {}^7_3\text{Li} + 2{}_1^3\text{T} + 3{}_1^1\text{H}$ , where two Y(2n)-bosons have transferred together. All these structures are stable states of  ${}^{16}_8\text{O}$  nucleus.

Let's come back to the virtual photons and gravitons. It is necessary to find a mechanism providing repulsion of two electric charges of the same sign and attraction of opposite sign charges. If the virtual particle is a usual plane-polarized photon, it is impossible to satisfy the specified requirements for the interaction between charges. So a virtual photon must be circularly polarized, (Figure 6-a).






Fig. 6. Circular right-hand polarized photon (a) and the graviton (b).

It should be taken into account that the virtual particle is coupled with emitting particle, i.e. a virtual particle is localized in a potential well.

Because a virtual boson can be presented as the boson coupled with a particle (Figure 7), the total energy of a particle with its virtual particles should be slightly higher (otherwise there will not be interaction between the particles) than the energy of the particles themselves<sup>5</sup>, but much smaller than the sum of the energies of the particles and released boson.

If we consider the electrically charged particles, it has to be considered that positive charges emit a circularly polarized photon of the first type (for instance, right-hand polarized one; but it is necessary to establish this), while negative charges emit photons of the second type. Absorption with the attraction between particles takes place, if the particle gets a virtual photon which is of different type than the particle emits. So the electron does not want to absorb the virtual photon emitted by other electrons. The scattering with repulsion will take place. Similar situation is observed for proton. Its own virtual photon after particles removal is reflected back to a potential well with a change of its direction of circular polarization (odd wave function). Then such photon will be absorbed by particle which was emitted it.

The proposed mechanism exhaustively describes electrostatic interaction experimental data.

Now take a look on gravitons. The main property of the gravitational field: there is an attraction between masses and there is no repulsion. However, according to the law of gravitational interaction, the mass will repel the negative mass (if there is a hypothetical negative mass). This is the first condition. And the second condition is: the graviton must be a boson with the spin  $s = 2$ .

These requirements can be met if a graviton is a double helix (Figure 6-b), like a DNA double helix. Because the wave function of the virtual graviton is supposed to be even, after a reflection it does not change the direction of circular polarization and can be absorbed by mass which was emitted it. If a graviton radiated by a negative mass, circular polarization changes a direction. Such negative graviton will be absorbed by a negative mass, but will be scattered by a usual mass. Thus, it will provide a repulsion of a usual mass from a negative mass.

<sup>5</sup> In this case a particle is in a coat of vacuum particles (bosons with zero energy).



Virtual pair of particles generated by the physical vacuum is different from a virtual photon near an electric charge because both particles in the pair (electron-positron or a virtual pair of other particles) are virtual, so they are situated in a deep potential well. This virtual pair annihilates without photon emission, because the total energy of a virtual pair is zero up to the uncertainty relation. However, such a virtual pair can interact with a real pair. As a result the wave function of a real particle can be complex leading to a strange behavior of particles.




Fig.7. Virtual particles in the vicinity of radiating particles.

## Conclusions

On the base of new ideas about the creation of the Universe and using of the Laws of similarity and unity in the Universe the description of the structure for the heavy ( $Z \geq 4$ ) cores and hierarchies of boson interaction is provide. In particular:

1. The classification of charges of elementary particles in different layers of the fiber space of Super-Universe was introduced. It was shown that diones with an electric charge  $\pm e/6$  should exist in the one-dimensional space, charges  $\pm e/3$  and  $\pm 2e/3$  should exist in the two-dimensional World (World of quarks), charges 0,  $\pm e$ ,  $\pm 2e$  and  $\pm 3e$  should exist in the three-dimensional space.

2. The model of the molecular structure of nuclei has been proposed and the reasons for unstability of nuclei in ground and excited states have been shown.

3. The hierarchy of bosons which are responsible for the interaction between particles in different hierarchical layers of the fiber space of Super-Universe has been analyzed.

4. New bosons have been proposed to explain the interaction between the elements of atomic nuclei. It has been shown that coupled neutron pairs (bineutrons) play the role of these bosons.

## References

- [1]. Petro O. Kondratenko. The Birth and Evolution of the Universe with Minimal Initial Entropy. // International Journal of Physics and Astronomy. December 2015, Vol. 3, No. 2, pp. 1-21. Published by American Research Institute for Policy Development DOI: 10.15640/ijpa.v3n2a1. URL: <http://dx.doi.org/10.15640/ijpa.v3n2a1>



- [2]. Victor V. Kulish. Hierarchic Electrodynamics and Free Electron Lasers: Concepts, Calculations, and Practical Applications. - CRC Press-Taylor & Francis Group. - 2011. – 697 p.
- [3]. D. Husemöller. Fibre Bundles. Springer Science & Business Media, 1994.- 353 p.
- [4]. Jean Letessier, Johann Rafelski, T. Ericson, P. Y. Landshoff. Hadrons and Quark-Gluon Plasma. — Cambridge University Press, 2002. — 415 p.
- [5]. K. Wildermuth, Ya. Tang, Unified theory of the nucleus. - Moscow: 1980 (in Russian, translated from English).
- [6]. O. F. Nemets. and others. Nucleon associations in atomic nuclei and nuclear reactions multi-nucleon transfer / Kyiv: 1988 .(in Russian).
- [7]. Peter W. Atkins and Ronald S. Friedman. Molecular Quantum Mechanics. Fifth Edition. - 2010. 560 p.
- [8]. S. G. Fedosin. Physics and Philosophy from the similarity to preons metagalaxies. - Perm: Style-MG. – 1999. - 544 pp.(in Russian).
- [9]. N. A. Kozyrev. Features of the physical structure of the components of binary stars. // Proceedings of the Main Astronomical Observatory in Pulkovo. - 1968. - №184. – c.108-115 (in Russian).
- [10]. Encyclopedia of astronomy and astrophysics [2002]. Copyright © Nature Publishing Group 2001. Brunei Road, Houndmills, Basingstoke, Hampshire, RG21 6XS, UK Registered No. 785998 and Institute of Physics Publishing 2001 Dirac House, Temple Back, Bristol, BS1 6BE, UK.
- [11]. Chemical Encyclopedia / Chief Editor I.L.Knunyants. - Moscow: Soviet Encyclopedia. – 1990 (in Russian).
- [12]. K. H. Beckurts and K. Wirtz. Neutron physics. / Translated from second Germany edition (1964). Springer-Verlag. – New York, 1964. – 444 pp.
- [13]. Tables of physical quantities / Directory ed. I.K Kikoin. – Moscow: Atomizdat. – 1976. – 1008 pp. (in Russian).
- [14]. Victor V. Kulish. Hierarchic Methods. Vol. I. Hierarchy and Hierarchical Asymptotic Methods in Electrodynamics. - Kluwer Academic Publishers. - 2002.
- [15]. J.D. Barrow, F.J. Tipler. The Anthropic Cosmological Principles. – Oxford: Clarendon Press. – 1986.
- [16]. Okun L B "The fundamental constants of physics" *Sov. Phys. Usp.* **34** (9) 818–826 (1991).
- [17]. Nakamura, K (2010). "Review of Particle Physics". *Journal of Physics G: Nuclear and Particle Physics*. **37** (7A): 075021. *Bibcode*:2010JPhG...37g5021N. *doi*:10.1088/0954-3899/37/7A/075021.
- [18]. R. Feynman. QED - a strange theory of light and substance. - Moscow: Nauka. – 1988. – 144 pp (in Russian, translated from English).