

Arbitrary l-state Solution of the Schrodinger Equation for q-deformed attractive radial plus Coulomb-like Molecular Potential within the framework of NU-Method.**Abstract**

The Schrodinger equation in 1-dimension for the q-deformed attractive radial plus coulomb-like molecular potential (ARCMP) is solved approximately to obtain bound states eigen solutions using the parametric Nikiforov-Uvarov (NU) method. The corresponding unnormalized eigen functions are evaluated in terms of Jacobi polynomials. Interestingly, the Klein-Gordon and Dirac equation with the arbitrary angular momentum values for this potential can be solved by this method.

Keywords: q-deformed potential, attractive radial, coulomb-like, Schrodinger

1 INTRODUCTION

An exact analytical solution of Schrodinger equation for central potentials has attracted enormous interest in recent years. So far, some of these potentials are the parabolic type potential [1], the Eckart potential [2, 3], the Fermi-step potential [2,3], the Rosen-Morse potential [4], the Ginocchio barrier [5], the Scarf barriers [6], the Morse potential [7] and a potential which interpolates between Morse and Eckart barriers [8]. Many researchers have investigated on exponential type potentials [9–12] and quasi-exactly solvable quadratic potentials [13–15]. Furthermore, Schrodinger, Dirac, Klein-Gordon, and Duffin-Kemmer-Petiau equations for a Coulomb type potential are solved by using different method [16–18]. Recently our group has also made significant progress in the use of combined or superposed molecular potentials to investigate the eigensolutions of relativistic and non-relativistic equations [19]. We have studied the eigensolutions (eigenvalues and eigenfunctions) of Klein-Gordon, Dirac and Schrodinger equations using superposed or mixed potentials. Some notable examples include Woods-Saxon plus Attractive Inversely Quadratic potential (WSAIQP) [19], Manning-Rosen plus a class of Yukawa potential (MRCYP) [20], generalized wood-Saxon plus Mie-type potential (GWSMP) [21], Kratzer plus Reduced Pseudoharmonic Oscillator potential (KRPHOP) [22], Inversely Quadratic Yukawa plus attractive radial potentials (IQYARP) [23], Modified Echart plus Inverse Square Molecular Potentials (MEISP) [24]

In nuclear and atomic physics, the shape form of a potential plays an important role, particularly when investigating the structure of deformed nuclei or the interaction between them. Therefore, our aim, in this present work, is to investigate approximate bound state solutions of the Schrodinger equation with q-deformed attractive radial plus coulomb-like molecular potential (qARCMP) using the parametric Nikiforov-Uvarov (NU) method. The solutions of this equation will definitely give us a wider and deeper knowledge of the properties of molecules moving under the sway of the superposed potential which is the goal of this paper. The parametric NU method is very convenient and does not require the truncation of a series like the series solution method which is more difficult to use. The organization of this work is as follows. In Section 2, we briefly introduce the basic concepts of the NU method. Section 3 is devoted to the solution of the Schrodinger problem to obtain the approximate bound-state energy of q-deformed attractive radial plus coulomb-like molecular potential (qARCMP) and their corresponding eigenfunctions

by applying the NU method. The results of special cases of potential consideration are discussed in Section 4. The scientific significance of this research paper includes giving an insight into possible eigensolutions of atoms and molecules moving under the influence of qARCOMP potential. Secondly, the resulting eigenenergy equations can be used to study the spectroscopy of some selected diatomic atoms and molecules.

2 REVIEW OF PARAMETRIC NIKIFAROV-UVAROV METHOD

The NU method is based on the solutions of a generalized second order linear differential equation with special orthogonal functions. The Nikiforov-Uvarov method has been successfully applied to relativistic and nonrelativistic quantum mechanical problems and other field of studies as well [25]. The hypergeometric NU method has shown its power in calculating the exact energy levels of all bound states for some solvable quantum systems.

$$\Psi_n''(s) + \frac{\tilde{\tau}(s)}{\sigma(s)} \Psi_n'(s) + \frac{\bar{\sigma}(s)}{\sigma^2(s)} \Psi_n(s) = 0 \quad (1)$$

Where $\sigma(s)$ and $\bar{\sigma}(s)$ are polynomials at most second degree and $\tilde{\tau}(s)$ is first degree polynomials. The parametric generalization of the N-U method is given by the generalized hypergeometric-type equation

$$\Psi''(s) + \frac{c_1 - c_2 s}{s(1 - c_3 s)} \Psi'(s) + \frac{1}{s^2(1 - c_3 s)^2} [-\epsilon_1 s^2 + \epsilon_2 s - \epsilon_3] \Psi(s) = 0 \quad (2)$$

Thus eqn. (1) can be solved by comparing it with equation (2) and the following polynomials are obtained

$$\tilde{\tau}(s) = (c_1 - c_2 s), \sigma(s) = s(1 - c_3 s), \bar{\sigma}(s) = -\epsilon_1 s^2 + \epsilon_2 s - \epsilon_3 \quad (3)$$

The parameters obtainable from equation (3) serve as important tools to finding the energy eigenvalue and eigenfunctions. They satisfy the following sets of equation respectively

$$c_2 n - (2n+1)c_5 + (2n+1)(\sqrt{c_9} + c_3\sqrt{c_8}) + n(n-1)c_3 + c_7 + 2c_3c_8 + 2\sqrt{c_8c_9} = 0 \quad (4)$$

$$(c_2 - c_3)n + c_3n^2 - (2n+1)c_5 + (2n+1)(\sqrt{c_9} + c_3\sqrt{c_8}) + c_7 + 2c_3c_8 + 2\sqrt{c_8c_9} = 0 \quad (5)$$

While the wave function is given as

$$\Psi_n(s) = N_{n,l} S^{c_{12}} (1 - c_3 s)^{-c_{12} - \frac{c_{13}}{c_3}} P_n^{(c_{10}-1, \frac{c_{11}}{c_3} - c_{10}-1)}(1 - 2c_3 s) \quad (6)$$

Where

$$c_4 = \frac{1}{2}(1 - c_1), c_5 = \frac{1}{2}(c_2 - 2c_3), c_6 = c_5^2 + \epsilon_1, c_7 = 2c_4c_5 - \epsilon_2, c_8 = c_4^2 + \epsilon_3,$$

$$c_9 = c_3c_7 + c_3^2c_8 + c_6, c_{10} = c_1 + 2c_4 + 2\sqrt{c_8}, c_{11} = c_2 - 2c_5 + 2(\sqrt{c_9} + c_3\sqrt{c_8})$$

$$c_{12} = c_4 + \sqrt{c_8}, c_{13} = c_5 - (\sqrt{c_9} + c_3\sqrt{c_8}) \quad (7)$$

and P_n is the orthogonal polynomials.

$$77 \quad \text{Given that } P_n^{(\alpha, \beta)} = \sum_{r=0}^n \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(\alpha+r+1)\Gamma(n+\beta-r+1)(n-r)!r!} \left(\frac{x-1}{2}\right)^r \left(\frac{x+1}{2}\right)^{n-r} \quad (8)$$

78 This can also be expressed in terms of the Rodriguez's formula

$$79 \quad P_n^{(\alpha, \beta)}(x) = \frac{1}{2^n n!} (x-1)^{-\alpha} (x+1)^{-\beta} \left(\frac{d}{dx}\right)^n ((x-1)^{n+\alpha} (x+1)^{n+\beta})$$

80

81 **3 EIGENSOLUTIONS OF THE SHRODINGER EQUATION WITH qARCMP**

82 The 1-State Schrodinger Equation with vector V(r), potential is given as [26-29]

$$83 \quad \frac{d^2 R(r)}{dr^2} + \frac{2\mu}{\hbar^2} \left[(E - V(r)) + \frac{l(l+1)\hbar^2}{2\mu r^2} \right] R(r) = 0 \quad (9)$$

84 Where E is the eigen energy value, l is the angular momentum quantum number

85 The q-deformed Attractive Radial Potential is given as [26]

$$86 \quad V(r) = - \left(\frac{V_1 e^{-4\alpha r} + V_2 e^{-2\alpha r} + V_3}{(1 - q e^{-2\alpha r})^2} \right) \quad (10)$$

$$87 \quad \text{Where } V_1 = \frac{\alpha^2}{4}, V_2 = \frac{(A-8)\alpha^2}{4}, V_3 = \frac{(4-A)\alpha^2}{4}$$

88 Where, screening parameter α determines the range of the potential, and V_1, V_2, V_3 are the
89 coupling parameters describing the depth of the potential well. In general q-deformed hyperbolic
90 functions are defined as

$$91 \quad \text{Sinh}_q(r) = \frac{1}{\text{Cosech}_q(r)} = \frac{e^{r-qe^{-r}}}{2}, \text{Cosh}_q(r) = \frac{e^r + qe^{-r}}{2}, \text{Coth}_q(r) = \frac{\text{Cosh}_q(r)}{\text{Sinh}_q(r)} \quad (11)$$

92

$$93 \quad \text{The Coulomb-like Potential, } V(r) = -\frac{A}{r} \quad (12)$$

94 Making the transformation $s = e^{-2\alpha r}$ the sum of the potentials (qARCMP) in equations (10) and
95 (12) becomes

96

$$97 \quad V(s) = \left(\frac{V_1 s^2 + V_2 s + V_3}{(1 - qs)^2} - \frac{2A\alpha}{(1 - qs)} \right) \quad (13)$$

98 By applying the Pekeris-like approximation [27, 28] to the inverse square term, $\frac{1}{r^2} = \frac{4\alpha^2}{(1-s)^2}$ to eq. (13)
99 enable us to completely solve eq. (9).

100 Again, applying the transformation $s = e^{-2\alpha r}$ to get the form that Nikiforov-Uvarov (NU)
101 method is applicable, equation (9) gives a generalized hypergeometric-type equation as

102

$$103 \quad \frac{d^2 R(s)}{ds^2} + \frac{(1-s)}{(1-s)s} \frac{dR(s)}{ds} + \frac{1}{(1-s)^2 s^2} [(2\beta^2 q^2 - B)s^2 + (-Hq - P - 4\beta^2 q)s + (2\beta^2 + H - J + \lambda)] R(s) = 0 \quad (14)$$

104

105 Where

$$106 \quad -\beta^2 = \left(\frac{\mu E}{4\alpha^2 \hbar^2}\right), \quad B = \left(\frac{\mu}{2\alpha^2 \hbar^2}\right) V_1, \quad \lambda = l(l+1), \quad P = \left(\frac{\mu}{2\alpha^2 \hbar^2}\right) V_2, \quad H = \left(\frac{\mu}{\alpha \hbar^2}\right) A, \quad J = \left(\frac{\mu}{2\alpha^2 \hbar^2}\right) V_3 \quad (15)$$

$$108 \quad c_1 = c_2 = c_3 = q, \quad c_4 = 0, \quad c_5 = -\frac{q}{2}, \quad c_6 = \frac{q^2}{4} + 2\beta^2 q^2 - B, \quad c_7 = -4\beta^2 q - P - Hq,$$

$$109 \quad c_8 = 2\beta^2 - J + H + \lambda, \quad c_9 = \frac{q^2}{4} + pq + Jq^2 + \lambda q^2 + B, \quad c_{10} = q + 2\sqrt{2\beta^2 - J + H + \lambda},$$

$$110 \quad c_{11} = 2 + 2\left(\sqrt{\frac{q^2}{4} + pq + Jq^2 + \lambda q^2 + B} + \sqrt{2\beta^2 - J + H + \lambda}\right), \quad c_{12} = \sqrt{2\beta^2 - J + H + \lambda},$$

$$111 \quad c_{13} = -\frac{1}{2} - \left(\sqrt{\frac{q^2}{4} + pq + Jq^2 + \lambda q^2 + B} + \sqrt{2\beta^2 - J + H + \lambda}\right), \quad \varepsilon_1 = 2\beta^2 q^2 + B,$$

$$112 \quad \varepsilon_2 = 4\beta^2 q + P + Hq, \quad \varepsilon_3 = 2\beta^2 + H - J + \lambda \quad (16)$$

113

114 Now using equations (6), (15) and (16) we obtain the energy eigen spectrum of the q-deformed
115 ARCMP as

116

$$117 \quad \beta^2 = \left[\frac{(2Jq - P - \lambda q) - q\left(n^2 + n + \frac{1}{2}\right) - (2n+1)\sqrt{\frac{q^2}{4} + pq + Jq^2 + \lambda q^2 + B}}{q\left(n + \frac{1}{2}\right) + 2\sqrt{\frac{q^2}{4} + pq + Jq^2 + \lambda q^2 + B}} \right]^2 - (J - H - \lambda) \quad (17)$$

118 The above equation can be solved explicitly and the energy eigen spectrum of q-deformed ARCMP
119 becomes

$$120 \quad E =$$

$$121 \quad \frac{4\alpha^2 \hbar^2}{\mu} \left\{ \left[\frac{\left(2q\left(\frac{\mu}{2\alpha^2 \hbar^2}\right)V_3 - \left(\frac{\mu}{2\alpha^2 \hbar^2}\right)V_2 - l(l+1)q\right) - q\left(n^2 + n + \frac{1}{2}\right) - (2n+1)\sqrt{\frac{q^2}{4} + q\left(\frac{\mu}{2\alpha^2 \hbar^2}\right)V_2 + q^2\left(\frac{\mu}{2\alpha^2 \hbar^2}\right)V_3 + l(l+1)q^2 + \left(\frac{\mu}{2\alpha^2 \hbar^2}\right)V_1}}{q\left(n + \frac{1}{2}\right) + 2\sqrt{\frac{q^2}{4} + q\left(\frac{\mu}{2\alpha^2 \hbar^2}\right)V_2 + q^2\left(\frac{\mu}{2\alpha^2 \hbar^2}\right)V_3 + l(l+1)q^2 + \left(\frac{\mu}{2\alpha^2 \hbar^2}\right)V_1}} \right] \right\} -$$

$$122 \quad \left(\left(\frac{\mu}{2\alpha^2 \hbar^2}\right)V_3 - \left(\frac{\mu}{\alpha \hbar^2}\right)A - l(l+1) \right) \quad (18)$$

123 In the standard case of the attractive radial potential where $q = 1$, our energy eigen spectrum
124 formula (eq. [18]) matches up with the results of parametric Nikifrov-Uvarov approach in ref.
125 [29]

126

127 We now calculate the radial wave function of the q-deformed ARCMP as follows

128

$$129 \quad \rho(s) = s^u (1 - qs)^v \quad (19)$$

$$130 \quad \text{Where } u = 2\beta^2 - J + H + \lambda, \text{ and } v = 2q\sqrt{\frac{q^2}{4} + pq + Jq^2 + \lambda q^2 + B}$$

$$131 \quad X_n(s) = p_n^{(u,v)}(1 - 2qs), \text{ where } p_n^{(u,v)} \text{ are Jacobi polynomials}$$

$$132 \quad \varphi(s) = s^{u/2} (1 - qs)^{1+v/2} \quad (20)$$

133 Radial wavefunction

$$134 R_n(s) = N_n \varphi(s) X_n(s) \quad (21)$$

$$135 R_n(s) = N_n s^{u/2} (1 - qs)^{1+v/2} P_n^{(u,v)}(1 - 2qs)$$

136

137 And using equation (16) we get

$$138 \varphi(s) = s^{u/2} (1 - s)^{v-1/2}, \quad (22)$$

139

140 We then obtain the radial wave function from the equation

$$141 R_n(s) = N_n \varphi(s) \chi_n(s),$$

142 As

$$143 R_n(s) = N_n s^{u/2} (1 - s)^{(v-1)/2} P_n^{(u,v)}(1 - 2s), \quad (23)$$

144

145 Where n is a positive integer and N_n is the normalization constant

146 4 DISCUSSION

147 We consider the following cases from equation (19)

148 CASE I: If we choose $V_1 = V_2 = V_3 = 0$ then the energy eigen values of the Coulomb-like molecular potential is
149 given as

$$150 E = \frac{4\alpha^2 \hbar^2}{\mu} \left\{ \left[\frac{l(l+1)q - q(n^2 + n + \frac{1}{2}) - (2n+1)\sqrt{\frac{q^2}{4} + l(l+1)q^2}}{q(n + \frac{1}{2}) + 2\sqrt{\frac{q^2}{4} + l(l+1)q^2}} \right] \right\} - \left(\left(\frac{\mu}{\alpha \hbar^2} \right) A - l(l+1) \right) \quad (24)$$

151

152 CASE II: If we choose $A = 0$ then the energy eigen values of the q-deformed Attractive Radial Potential

$$153 E =$$

$$154 \frac{4\alpha^2 \hbar^2}{\mu} \left\{ \left[\frac{\left(2q \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_3 - \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_2 - l(l+1)q \right) - q \left(n^2 + n + \frac{1}{2} \right) - (2n+1) \sqrt{\frac{q^2}{4} + q \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_2 + q^2 \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_3 + l(l+1)q^2 + \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_1}}{q \left(n + \frac{1}{2} \right) + 2 \sqrt{\frac{q^2}{4} + q \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_2 + q^2 \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_3 + l(l+1)q^2 + \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_1}} \right] \right\} -$$

$$155 \left(\left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_3 - l(l+1) \right) \quad (25)$$

156 CASE III: If we choose $l = 0$ then the eigen energy spectrum of the s-wave 1-dimensional Schrodinger
157 equation with q-deformed ARCMP

$$158 E = \frac{4\alpha^2 \hbar^2}{\mu} \left\{ \left[\frac{\left(2q \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_3 - \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_2 \right) - q \left(n^2 + n + \frac{1}{2} \right) - (2n+1) \sqrt{\frac{q^2}{4} + q \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_2 + q^2 \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_3 + \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_1}}{q \left(n + \frac{1}{2} \right) + 2 \sqrt{\frac{q^2}{4} + q \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_2 + q^2 \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_3 + \left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_1}} \right] \right\} -$$

$$159 \left(\left(\frac{\mu}{2\alpha^2 \hbar^2} \right) V_3 - \left(\frac{\mu}{\alpha \hbar^2} \right) A \right) \quad (26)$$

160

161 5 CONCLUSION

162 In this work, using the parametric generalization of the NU method, we have obtained
 163 approximately energy eigenvalues and the corresponding wave functions of the Schrodinger
 164 equation for q-deformed attractive radial plus Coulomb-like molecular potential. The
 165 corresponding unnormalized eigen functions are evaluated in terms of Jacobi polynomials. Interestingly,
 166 the Klein-Gordon and Dirac equation with the arbitrary angular momentum values for this
 167 potential can be solved by this method. The resulting eigen energy equations can be used to
 168 study the spectroscopy of some selected diatomic atoms and molecules.

169

170 6 REFERENCES

- 171 [1] G. Barton, Ann. Phys. **166**, 322 (1986).
- 172 [2] S. Flügge, Practical Quantum Mechanics I (Springer-Verlag, Berlin, 1974).
- 173 [3] L. D. Landau and E.M. Lifshitz, Quantum Mechanics (Pergamon Press, London, 1958).
- 174 [4] P.M Morse and H. Feshbach, Methods of Theoretical Physics (McGraw-Hill Book Company
175 Ltd., New York,1953).
- 176 [5] B. Sahu, S. K. Agarwalla, and C. S. Shastri, J. Phys. A **35**, 4349 (2002).
- 177 [6] A. Khare and U.P. Sukhatme, J. Phys. A **21**, L501 (1988).
- 178 [7] Z. Ahmed, Phys. Lett. A **157**, 1 (1991).
- 179 [8] Z. Ahmed, Phys. Rev. A **47**, 4757 (1993).
- 180 [9] M. Znojil, Phys. Lett. A **264**, 108 (1999).
- 181 [10] D.T. Barclay et al., Phys. Lett. A **305**, 231 (2002).
- 182 [11] G. Lévai and M. Znojil, J. Phys. A **35**, 8793 (2002).
- 183 [12] C. S. Jia, X. L. Zeng, and L.T. Sun, Phys. Lett. A **294**, 185 (2002).
- 184 [13] Z. Ahmed, Phys. Lett. A **290**, 19 (2001).
- 185 [14] C.M. Bender and S. Boettcher, J. Phys. A **31**, 1273 (1998).
- 186 [15] M. Znojil, J. Phys. A **33**, 4203 (2000).
- 187 [16] F.Yasuk, C. Berkdemir, A. Berkdemir, and C. Oñem, Physica Scripta **71**, 340 (2005).
- 188 [17] M. S. İm, İ. Şek and H. Eşgrifes, J. Phys. A **37**, 4379 (2004).
- 189 [18] H. Eşgrifes and R. Sever, Phys. Lett. A **344**, 117 (2005).
- 190 [19]B.I. Ita, H. Louis, T.O. Magu and N.A. Nzeata-Ibe (2017). WSN 74 (2017) 280-287
- 191 [20]H. Louis, B.I. Ita, T.O. Magu and N.A Nzeata-Ibe (2017) WSN 70(2) (2017) 312-319
- 192 [21]B.I. Ita, B. E. Nyong., H. Louis, T.O. Magu (2017): *J. Chem. Soc. Nigeria, Vol. 41, No. 2, pp21-26*
- 193 [22]H. Louis*, B.I. Ita., B.E. Nyong., T.O.Magu, S. Barka and N.A. Nzeata-Ibe. *NAMP* vol.36, No. 2,
194 (July, 2016) pp.193-198
- 195 [23]B.I. Ita, B.E. Nyong, N.O. Alobi, H. Louis and T.O. Magu (2016). *Equatorial Journal of*
196 *Computational and Theoretical Sciences*, Volume 1, Issue 1, (2016), pp. 55-64.
- 197 [24]B.I. Ita, A.I. Ikeuba, H. Louis and P. Tchoua (2015):. *Journal of Theoretical Physics and*
198 *Cryptography. IJTPC, Vol. 10, December, 2015. www.IJTPC.org*
- 199 [25]M. K. Bahar and F. Yasuk. PRAMANA J. Phys.80 (2013) 187-197
- 200 [26]Eshghi, M. and Hamzavi, M. Commun. (2012) Theor. Phys. 57 355-360
- 201 [27] Louis H., Ita B.I., Nelson N.A., I. Joseph., Amos P.I and Magu T.O; ...*International Research*
202 *Journal Of Pure and Applied Physics*. Vol. 5, No. 3, pp 27-32, 2017

203 [28] Louis Hitler, Benedict Iserom Ita, Pigweh Amos Isa, Nzeata-Ibe Nelson, Innocent Joseph, Opara
204 Ivan, Thomas Odey Magu. *World Journal of Applied Physics*. Vol. 2, No. 4, 2017, pp. 109-112.

205 [29] Louis Hitler, Benedict Iserom Ita, Pigweh Amos Isa, Innocent Joseph, Nzeata-Ibe Nelson, Thomas
206 Odey Magu, Opara Ivan.. *World Journal of Applied Physics*. Vol. 2, No. 4, 2017, pp. 101-108.

207

208

209

210

211

212

213

214

215

216