

Dense Phase Carbon dioxide: An emerging Non thermal technology in Food Processing

Abstract

DPCD is an emerging non-thermal process which evidently inactivates certain micro organisms and enzymes at lower temperatures and high pressure combination thereby maintaining fresh like characteristics as well as providing convenient shelf life, nutritional value and the product. DPCD ensures minimal nutrient loss and better preserved quality attributes as compared to any thermal treatment. This technology has been under investigation **into** past few decades and various studies have been carried on to demonstrate its mechanism and effect on various microorganisms such as bacteria , yeast and moulds. Many liquid foods have shown promising results. This article is a review of effect of DPCD on the quality and safety of foods with special application **of** juices, beverage and dairy industry.

Keywords : DPCD, Non thermal treatment, shelf life, nutritional value

Introduction:

Non-thermal food preservation techniques have been recent trend owing to consumer demand for fresh-like food products. Thermal pasteurization is the most common method to prevent microbial spoilage of highly acidic juices. However, it can cause degradation of flavor, nutrient, color and texture. Dense phase CO₂ is a promising non-thermal process to preserve mainly liquid foods. CO₂ at ambient pressure can inhibit microorganisms but, when it is applied **for** elevated pressures, it can effectively inactivate a number of microorganisms. Fraser (1951) proposed a technique for collecting the content of Escherichia coli by bursting cells in liquid culture with a sudden release of pressurized Ar, N₂, N₂O or CO₂, and suggested that

pressurized CO₂ could be used for the inactivation of E. coli.

Several authors worked on use of DPCD for inactivation of micro-organisms, enzymes in liquid foods and its effect on physico-chemical qualities. DPCD has been shown to eliminate vegetative forms of spoilage and pathogenic bacteria, moulds, yeasts, and can inactivate some enzymes. **Damar (2006)** These enzymes include polyphenol oxidase, which causes browning of fruits, juices, seafood and vegetables; pectinesterase in fruit juice, which causes cloud loss; peroxidase, which causes food discolouration; and lipoxygenase, which destroys chlorophyll and contributes to the development of off-flavours in frozen vegetables. Gui (2005) reported that high-

pressure CO₂ had a significant effect on the inactivation of polyphenol oxidase in apple juice maintaining only 38.5% residual activity at 30 MPa and 55 °C for 60 min.

Microbial inactivation by DPCD is dependant on different parameters like exposure time, pressure, temperature, pressure cycling, initial pH of medium, water activity, cell growth phase or age, species of microorganisms, and type of treatment system. (Kamihira et al., 1987; Arreola et al., 1991; Lin et al., 1993; Dillow et al., 1999; Hong et al., 1999; Spilimbergo et al., 2003b). Most of DPCD inactivation studies currently available in the scientific literature have been performed on inoculated or spoiled foods (Arreola et al., 1991; Ferrentino et al., 2013a). High pressure facilitates CO₂ solubilization in water and penetration through cell walls, and increases density and therefore extraction power. Higher temperature enhances deactivation by increasing the fluidity of cell membranes, making them easier to penetrate, and increasing the diffusivity of CO₂. However, higher temperatures may reduce the ability of CO₂ to extract low-volatility materials and decrease CO₂ solubility in aqueous media. Depending on temperature and pressure, CO₂ exists on the gas, liquid or supercritical fluid state. CO₂ exists in its supercritical state above 31 °C and 7.34 MPa. Below this critical temperature–pressure combination, CO₂ exists on the subcritical liquid or gaseous state depending on the specific temperature–pressure combination. Supercritical CO₂ has properties of both liquid and gaseous CO₂ with altered viscosity, diffusivity and

solubility, resulting in improved dissolving power. The gas-like diffusivity allows supercritical CO₂ to quickly diffuse through complex matrices; and the liquid-like density confers high extraction power. CO₂ is having relatively low critical pressure (74 bar) and temperature (32 °C), relatively non-toxic, non-flammable, available in high purity at relatively low cost, and is readily available. Kamihira et al. (1987) reports that Baker's yeast, E. coli and S. aureus were completely sterilized by SC-CO₂ at 200 atm and 35° C while no sterilizing effect was detached with G- and L-CO₂. Experimental results have been also published showing DPCD effects on other products as dry cured ham (Ferrentino et al., 2013b), fresh cut pear (Valverde et al., 2010), fresh cut carrot (Spilimbergo et al., 2013), spinach leaves (Zhong et al., 2008), alfalfa sprouts (Jung et al., 2009).

Inactivation kinetics:

Lin et al. observe a two-stage kinetic curve. The first stage was characterized by a slow deactivation rate, and the second stage by fast linear deactivation. Two-stage kinetics were also observed by several other researchers (Fig. 3(a) and (b)). Some studies showed a fast initial rate followed by a slow deactivation stage (Fig. 3(d)). Interestingly, both kinds of kinetic curves appeared in Hong and Pyun [65] (Fig. 3(e)). One reason for variation in kinetic behavior may be the difference in the efficiency of contact between CO₂ and the microorganism. That depends on process temperature, pressure, state of CO₂, type of suspension medium etc.




Fig. 1. Different kinds of kinetic curves reported in publications (J. Zhang et al., 2006).

Deactivation mechanism:

Physical cell rupture was the earliest proposed mechanism. Cells were presumed to rupture because of the explosive expansion of high-pressure CO₂ during flash depressurization. However, based on electron microscopy, Enomoto et al. proposed that, the rupture of cells happens *during* pressurization when the cells are swollen by CO₂, rather than during the depressurization stage. The physiological

deactivation mechanisms include several different scenarios,

Step 1: solubilization of pressurized CO₂ in the external liquid phase

Water in contact with pressurized CO₂ generally becomes acidic due to the formation and dissociation of H₂CO₃, which liberates H⁺ ions. This lowered extracellular pH may inhibit microbial growth (Valley and Rettger, 1927; Daniels et al., 1985;

Hutkins and Nannen, 1993). However, a reduction in pH is not enough to account for the lethal effect of CO₂. Lin et al. (1993, 1994) therefore suggested that the lowered pH contributes to an increase in cell permeability, which facilitates the penetration of CO₂ into microbial cells.

Step 2: cell membrane modification

Aqueous CO₂ may diffuse into the cellular membrane and may accumulate into its lipophilic (phospholipid) inner layer (Isenschmid et al., 1995). This accumulated amount of CO₂ in the lipid phase may then structurally and functionally disorder the cell membrane due to an order loss of the lipid chain (a process known as “anesthesia”) (Jones and Greenfield, 1982), which may increase the fluidity and hence permeability of the membrane (Isenschmid et al., 1995).

Step 3: intracellular pH decrease

Due to the increased membrane permeability, pressurized CO₂ may easily penetrate through the bacterial cell membrane and accumulate in the cytoplasmic interior of bacterial cells. There, the relative concentrations of both aqueous CO₂ and HCO₃⁻² are in first instance controlled by internal pH buffering as a result of pH homeostasis in order to maintain a more or less constant cytoplasmic pH_i. A more important homeostatic system is a membrane-bound H⁺-ATPase (Hutkins and Nannen, 1993), which expels protons from the cytoplasm against the prevailing pH gradient and electrochemical gradient. If too much dissolved CO₂ enters the cytoplasm, the cells may be unable to expel

all the resulting protons and pH_i will start to decrease. If pH_i is lowered too much, cell viability will seriously be impaired.

Step 4: key enzyme inactivation/cellular metabolism

Enzymes have maximal activity at the optimum pH, and their activity declines sharply on either side of the optimum. So lowering of the cytosolic pH_i might cause inhibition and/or inactivation of key enzymes essential for metabolic and regulating processes (Hutkins and Nannen, 1993).

Step 5: direct (inhibitory) effect of molecular CO₂ and HCO₃⁻² on metabolism

The reaction rate of each enzymatic reaction is not only a function of the pH but also of the intracellular concentrations of its substrate(s), product(s), and cofactor, which are primary elements in the regulation of enzymatic activity. The concentration of HCO₃⁻ appears to be central to the regulation of enzymatic activity (and hence cellular metabolism). Dissolved CO₂ can inhibit decarboxylation reactions.

Step 6: disordering of the intracellular electrolyte balance

Lethal damage to the biological system of the cells may also be produced when the applied CO₂ pressure accumulates in the cytoplasmic interior of the bacterial cells. This may convert HCO₃⁻² to CO₃⁻², which could precipitate intracellular inorganic electrolytes (such as Ca²⁺, Mg²⁺ and similar ions) from cells and cell membranes (Lin et al., 1993). Since these inorganic electrolytes aid in maintaining the osmotic relationships

between cells and their surrounding media, this could have deleterious effects on the volume of cells.

Step 7: removal of vital constituents from cells and cell membranes

The pressurized CO₂ first penetrates into the cells to build up the density of a critical level within the cells, after which it removes intracellular constituents (such as phospholipids and hydrophobic compounds) to disturb or alter the structure of the biomembrane and/or the balance of the

biological system, thus promoting inactivation (Lin et al., 1992a, 1993). This removal process appeared to be stimulated by a sudden release of the applied pressure, leading to a rapid transfer of intracellular materials out of the biological system into the extracellular environment (Lin et al., 1992a, 1993). Lin et al. (1992a) also suggested that the inactivation rate could be improved by repeating the release and recharge of pressurized CO₂ in the pressure vessel during the treatment to improve transfer of intracellular materials out of the bacterial cells.




Fig. 2. Proposed deactivation mechanisms (J. Zhang et al., 2006).

DPCD APPLICATIONS IN FOOD PROCESSING:

This process of cold pasteurization is applied mostly to liquid foods mainly fruit juices. At high pressure, the inactivation of *E. coli* exposed to DPCD was dramatically increased as the temperature increased. (Valley et al., 1977). Similar results were earlier observed (Hong et al., 1999) Higher

temperatures stimulated the diffusivity of supercritical CO₂ and also increased the fluidity of cell membrane to make its penetration of the cell easier (Hong et al., 1997). DPCD increases the susceptibility of apple. It had a noticeable effect on apple PME activity. Z values representing temperature increase needed for a 90% reduction of the D value and the activation energy E_a of the labile fraction of 30 MPa

was found to be 22.32 °C and 86.88 kJ/mol (Zhi et al., 2008). DPCD has been added to cottage cheese, ice cream, yogurt and ricotta cheese because CO₂ is highly soluble in lipids and aqueous solutions. The gas decreases the growth rate of microorganisms, displaces oxygen thereby minimizing rancidity, and can be combined with barrier packaging to extend shelf life, in some cases by triple (Morris et al., 2007). A continuous high-pressure carbon dioxide system, running at ambient conditions, was tested for its performance in reducing both natural and inoculated microbial loads. The prototype system continuously processes orange juice with carbon dioxide (CO₂) at high pressures. The unit was able to cause a 5-log reduction of the natural flora in spoiled juice, and could attain a 5-log decrease in numbers of pathogenic *Escherichia coli*, *Salmonella typhimurium*, and *Listeria monocytogenes*. (Kincal et al., 2005). The influence of thermal and dense-phase carbon dioxide (DPCD) pasteurization on physicochemical properties and flavor compounds in melon juice was investigated. Melon juice was pasteurized using DPCD treatment and compared to a conventional high-temperature-short-time (HTST) method. The DPCD treatment was carried out using a DPCD unit (55 °C, 60 min, and 35 MPa). The thermal pasteurization was performed at 90 °C for 60 s with an adapted laboratory setup. The changes of pH and organic acid and sugar concentrations were not significant. There were significant differences between treatments of microbial count, vitamin C, β -carotene, and volatile

compound concentrations. In general, DPCD treatment had less of an effect on the measured variables than the thermal treatment but significant enough to cause microbial inactivation and nutrient preservation. (Zhang et al., 2009). Red grapefruit juice was treated with continuous dense phase carbon dioxide (DPCD) equipment to inactivate yeasts and molds and total aerobic microorganisms. A central composite design was used with pressure. (13.8, 24.1, and 34.5 MPa) and residence time (5, 7, and 9 min) as variables at constant temperature (40 °C), and CO₂ level (5.7%) after experimentally measuring CO₂ solubility in the juice. Five log reduction in yeasts and moulds and total elimination of aerobic microorganisms occurred at 34.5 MPa and 7 min of treatment (Ferrantino et al., 2009). DPCD technology was investigated for its effects on milk processing (Tomasula 1997; Hofland 1999; Tisi 2004). Depending on the purpose of the process, the effects of DPCD on milk can be desirable or undesirable. Tisi (2004) showed that DPCD-treated milk had higher lipolytic activity during storage compared with the untreated milk because of the homogenization effect of DPCD on the fat micelles. Tomasula (1997) and Hofland (1999) showed that DPCD can be used for casein production, due to the pH lowering effect. The use of DPCD has advantages over a traditional process using lactic or mineral acids for casein precipitation. CO₂ is removed from the system after DPCD treatment, eliminating additional treatment steps (Tomasula 1997).

Table 1 —Summary of the studies on inactivation of vaious microorganisms by dense phase CO2 (DPCD).

S No	Solution	Microorganism	Pressure MPa	Time	Temperature (°C)	System	Log reduction	Reference
1.	Physiological saline	Saccharomyces cerevisiae	20	2h	35	Batch	7.5 (C)	Kamihira and others 1987
2.	Herbs	Total bacteria count	5.52	2h	45	Batch	6.5 (C)	Haas and others 1989
3.	Apple juice Orange juice Nutrient broth	Total bacteria count E. coli	5.52	30 min	45 55	Batch	>3 (C) 4 (C)	
4.	Distilled water	Listeria monocytogenes	6.18	2h	35	Batch	9	Wei and others 1991
5.	Egg yolk Orange juice	Salmonella thyphimurium	2 h	2h 33 mins	8 35	Batch	2	Arreola and others 1991b
6.	Growth medium	S. cerevisiae	6.9 to 20.7	15 mins	35	Batch	7 (C)	Lin and others 1992
7.	Growth medium	Leuconostoc dextranicum	4	15-20 mins	40	Batch	>8	Lin and others 1993
8.	Sterile water	S. cerevisiae	25	< 3hrs	35	Batch	8 (C)	Nakamura and others 1994
9.	Physiological saline	Lactobacillus brevis	15	30 mins	35	Batch	6 (C)	Ishikawa and others 1995b
10.	Physiological saline	S. cerevisiae	25	30 mins	35	Micro bubble	6 (C)	Ballestra and others 1996
11.	Sterile water	S. cerevisiae	15	1 hr	30	Batch	8 (C)	Kumagai and others 1997
12.	Whole milk	Aerobic plate count	14.5	5 hrs	25	Batch	>8	Shimoda and others 1998

CONCLUSION:

In some years, DPCD treatment could become one of the most available emergent technologies. However, to meet this high expectation, consumers and stakeholders must be convinced about the improvements this new technology represents. This will require convincing data, and provision of clear, objective and unbiased information also including the potentially negative aspects of the technology and their limitations. HPP (pressures above 100 MPa) has been studied intensively for a number of years and continues to the present (Yuste, Capellas et al. 2001). It is currently being used commercially to process higher value products that are perishable and have few processing alternatives such as guacamole and salsas.

Although the FDA has not issued a final ruling, an additional benefit of using pressure to preserve foods might be to label foods as “fresh”, a term highly valued by consumers without having any adverse effects while handling and usage. Impact on green house effect can be considered to be minimal yet a study needs to be ascertained.

Also, for being able to replace other preservation techniques, DPCD treatment must not only improve food quality, but also promote shelf life and (long-term) safety by inactivating spoilage and pathogenic microorganisms. Therefore, further research is essential to demonstrate and explain the effect of DPCD preservation on the shelf life and safety of food products. In addition, it is important that the effect of a DPCD treatment on the sensory and nutritional quality of both liquid and solid foods is more thoroughly investigated. Also the economics of the process must be assessed. Finally, although this review clearly shows that DPCD has great potential for improving the safety and quality of foods, some technological and regulatory hurdles (such as further optimization of the process, proper scale-up, acquisition of more sensory and shelf life data, quality certification to be obtained, etc.) still need to be overcome before the supply chain can receive these benefits.

References:

- A.G. Arreola, M.O. Balaban, C.I.Wei, A. Peplow, M. Marshall, J. Cornell, Effect of supercritical carbon dioxide on microbial populations in single strength orange juice, *J. Food Qual.* 14 (1991) 275.
- A.K. Dillow, R.S. Langer, N. Foster, J.S. Hrkach, Supercritical fluid sterilization method, United States Patent, US 6,149,864 (2000).
- Ballestra, P., Dasilva, A.A., Cuq, J.L., 1996. Inactivation of *Escherichia coli* by carbon dioxide under pressure. *Journal of Food Science* 61, 829–836.
- Chen, J., Zhang, J., Feng, Z., Song, L., Wu, J., & Hu, X. (2009). Influence of thermal

and dense-phase carbon dioxide pasteurization on physicochemical properties and flavor compounds in Hami melon juice. *Journal of agricultural and food chemistry*, 57(13), 5805-5808

D. Fraser, Bursting bacteria by release of gas pressure, *Nature* 167 (1951) 33.

Damar S, Balaban MO. Review of dense phase CO₂ technology: microbial and enzyme inactivation, and effects on food quality. *J. Food Sci.* 2006; **71**(1): R1–R11.

Daniels, J.A., Krishnamurthi, R., Rizvi, S.S.H., 1985. A review of effects of carbon dioxide on microbial growth and food quality. *Journal of Food Protection* 48, 532–537.

Ferrentino, G., Balzan, S., Spilimbergo, S., 2013a. Optimization of supercritical carbon dioxide treatment for the inactivation of the natural microbial flora in cubed cooked ham. *Int. J. Food Microbiol.* 161, 189e196

Ferrentino, G., Balzan, S., Spilimbergo, S., 2013b. Supercritical carbon dioxide processing of dry cured ham spiked with *Listeria monocytogenes*: inactivation kinetics, color, and sensory evaluations. *Food Bioprocess Technol.* 6, 1164e1174

Ferrentino, G., Plaza, M. L., Ramirez-Rodrigues, M., Ferrari, G., & Balaban, M. O. (2009). Effects of dense phase carbon dioxide pasteurization on the physical and quality attributes of a red grapefruit juice. *Journal of food science*, 74(6).

Gui, F., Wu, J., Chen, F., Liao, X., Hu, X., Zhang, Z., Wang, Z., 2005. Change of

polyphenol oxidase activity, color, and browning degree during storage of cloudy apple juice treated by supercritical carbon dioxide. *European Food Research and Technology* 223, 427–432.

H. Ishikawa, M. Shimoda, K. Tamaya, A. Yonekura, T. Kawano, Y. Osajima, Inactivation of *Bacillus* spores by the supercritical carbon dioxide micro-bubble method, *Biosci. Biotechnol. Biochem.* 61 (1997) 1022.

H.M. Lin, Z.Y. Yang, L.F. Chen, Inactivation of *Saccharomyces cerevisiae* by supercritical and subcritical carbon dioxide, *Biotechnol. Progress* 8 (1992) 458.

Hong, S.I., Park, W.S., Pyun, Y.R., 1999. Non-thermal inactivation of *Lactobacillus plantarum* as influenced by pressure and temperature of pressurized carbon dioxide. *International Journal of Food Science & Technology* 34, 125–130.

Hutkins, R.W., Nannen, N.L., 1993. pH homeostasis in lactic-acid bacteria. *Journal of Dairy Science* 76, 2354–2365.

Jones, R.P., Greenfield, P.F., 1982. Effect of carbon dioxide on yeast growth and fermentation. *Enzyme and Microbial Technology* 4, 210–223.

Jung, W.Y., Choi, Y.M., Rhee, M.S., 2009. Potential use of supercritical carbon dioxide to decontaminate *Escherichia coli* O157:H7, *Listeria monocytogenes*, and *Salmonella typhimurium* in alfalfa sprouted seeds. *Int. J. Food Microbiol.* 136, 66e70.

K. Nakamura, A. Enomoto, H. Fukushima, K. Nagai, M. Hakoda, Disruption of microbial-cells by the flash discharge of

high-pressure carbondioxide, *Biosci. Biotechnol. Biochem.* 58 (1994) 1297.

Kincal, D., Hill, W. S., Balaban, M. O., Portier, K. M., Wei, C. I., & Marshall, M. R. (2005). A continuous high pressure carbon dioxide system for microbial reduction in orange juice. *Journal of Food Science*, 70(5).

Lin, H.-M., Cao, N.J., Chen, L.-F., 1994. Antimicrobial effect of pressurized carbon dioxide on *Listeria monocytogenes*. *Journal of Food Science* 59, 657–659.

Lin, H.-M., Chan, E.-C., Chen, C., Chen, L.F., 1991. Disintegration of yeast cells by pressurized carbon dioxide. *Biotechnology Progress* 7, 201–204.

Lin, H.-M., Yang, Z.Y., Chen, L.-F., 1992a. Inactivation of *Saccharomyces cerevisiae* by supercritical and subcritical carbon dioxide. *Biotechnology Progress* 8, 458–461.

Lin, H.-M., Yang, Z.Y., Chen, L.-F., 1992b. An improved method for disruption of microbial cells with pressurized carbon dioxide. *Biotechnology Progress* 8, 165–166.

Lin, H.-M., Yang, Z.Y., Chen, L.-F., 1993. Inactivation of *Leuconostoc dextranum* with carbon dioxide under pressure. *Chemical Engineering Journal and the Biochemical Engineering Journal* 52, B29–B34.

M. Kamihira, M. Taniguchi, T. Kobayashi, Sterilization of microorganisms with supercritical carbon dioxide, *Agric. Biol. Chem.* 51 (1987) 407.

Morris, C., Brody, A. L., & Wicker, L. (2007). Non-thermal food processing/preservation technologies: a review with packaging implications. *Packaging Technology and Science*, 20(4), 275-286.

O. Erkmen, H. Karaman, Kinetic studies on the high pressure carbon dioxide inactivation of *Salmonella typhimurium*, *J. Food Eng.* 50 (2001) 25.

P. Ballestra, A.A. Dasilva, J.L. Cuq, Inactivation of *Escherichia coli* by carbon dioxide under pressure, *J. Food Sci.* 61 (1996) 829.

S. Spilimbergo, F. Dehghani, A. Bertucco, N.R. Foster, Inactivation of bacteria and spores by pulse electric field and high pressure CO₂ at low temperature, *Biotechnol. Bioeng.* 82 (2003) 118.

S.I. Hong, Y.R. Pyun, Membrane damage and enzyme inactivation of *Lactobacillus plantarum* by high pressure CO₂ treatment, *Int. J. Food Microbiology*. 63 (2001) 19.

senschmid, A., Marison, I.W., von Stockar, U., 1995. The influence of pressure and temperature of compressed CO₂ on the survival of yeast cells. *Journal of Biotechnology* 39, 229–237.

Spilimbergo, S., Komes, D., Vojvodic, A., Levaj, B., Ferrentino, G., 2013. High pressure carbon dioxide pasteurization of fresh-cut carrot. *J. Supercrit. Fluids* 79, 92e100

Valley, G., Rettger, L.F., 1977. The influence of carbon dioxide on bacteria. *Journal of Bacteriology* 14, 101–137.

Valverde, M.T., Marín-Iniesta, F., Calvo, L., 2010. Inactivation of *Saccharomyces cerevisiae* in conference pear with high pressure carbon dioxide and effects on pear quality. *J. Food Eng.* 98, 421e428.

Zhi, X., Zhang, Y., Hu, X., Wu, J., & Liao, X. (2008). Inactivation of apple pectin methylesterase induced by dense phase carbon dioxide. *Journal of agricultural and food chemistry*, 56(13), 5394-5400.

Zhong, Q., Black, D.G., Davidson, P.M., Golden, D.A., 2008. Nonthermal inactivation of *Escherichia coli* K-12 on spinach leaves, using dense phase carbon dioxide. *J. Food Prot.* 71, 1015e1017.