

# An Experimental Study to Examine the Curved Spacetime Using Magnetic Fields

## ABSTRACT

The curvature of spacetime represented by Einstein field equation has many physical implications, including gravity. As light is deflected by the curvature of spacetime, a magnetic field will also be influenced by the curved spacetime. A permanent magnet is generally known to maintain its own persistent magnetic field on the ground as long as there is no external magnetic interference. However, a series of experiments find that there are noticeable changes in the magnetic fields distribution while the permanent magnet rotates. The magnetic field lines of the permanent magnet are deflected towards Earth's center, implying a possibility that we can use magnetic field, a more efficient tools than a satellite, to measure the curvature of spacetime.

*Keywords: curvature of spacetime; magnetic field; Einstein field equation*

## 1. INTRODUCTION

The Einstein field equation [1,2] that mass warps spacetime was announced in 1916. Yet, the theory of general relativity was not known to the public until Arthur Eddington observed the deflection of starlight around the Sun in 1919[3, 4]. Besides the Arthur Eddington's discovery, several experimental studies have verified the Einstein field equation through recession of Mercury's orbit, a gravitational redshift in the Pound–Rebka[5], gravitational lensing[6-9] around the galaxies, and so on. Despite many studies confirming the Einstein field equation, however, not many have directly measured the curvature, which is the key to the Einstein field equation. In order to verify the geodetic effect or the frame-dragging effect predicted in Einstein field equation, it is critical to measure the curvature of spacetime around us. Yet, given that the deflection of solar light by the local spacetime of Earth is not more than 40 micro arcseconds[10], examining the curvature of spacetime still remains challenging. NASA has been trying to measure Earth's curvature of spacetime with a precise gyroscope since 1964. Recently, NASA has successfully detected the precession caused by the geodetic effect with four gyroscopes in Gravity Probe B (GP-B) experiments [11-13]. However, it is hard to say that the experiment was efficient, because it required (1) expensive satellites and (2) perfect gyroscopes to measure the minuscule angle.

Here, I am suggesting an alternative and more efficient method to measure the curvature of spacetime – that is, using magnetic fields on the ground. As light, a type of electromagnetic wave, is deflected by the curvature of spacetime, a magnetic field is influenced by the curved spacetime [14-17]. Maxwell's equations are generally used on the assumption that the spacetime is flat. However, it is necessary to use Maxwell's equations in curved spacetime in order to observe magnetic fields on the ground more precisely[17]. Thus, if there are noticeable changes in the magnetic fields due to the curvature of spacetime on the ground, a magnetic field will be a good tool to measure the curvature of spacetime.

## 37 2. EXPERIMENTAL APPROACH TO EXAMINE THE CURVED SPACETIME USING 38 MAGNETIC FIELDS

39

40 To test the above ideas on the effect of curvature in spacetime on magnetic field, this study  
41 conducts a series of experiments using a permanent magnet attached on a rotating table. A  
42 permanent magnet is used, because it maintains its own persistent magnetic field as long as there is  
43 no external magnetic interference. A disc-type neodymium permanent magnet having a hole in the  
44 center is used in the experiments. The neodymium magnet has 30mm of outer diameter, 3mm of  
45 thickness and 7mm of the inner diameter (Fig. 1). The magnetic measurement instrument is a gauss  
46 meter made by keuwlsoft. The permanent magnet and the magnetic field sensor are fastened to the  
47 rotating table (Fig. 2). The magnetic field sensor is located 4mm away from the center line of the  
48 permanent magnet, facing the S pole (Fig. 1). The sensor measures the magnitude of Y-directional  
49 magnetic flux density of the permanent magnet (Fig. 1). Two experiments have been done in Jeju  
50 volcanic island.

51 In the first experiment, the plane of the rotating table is set to be parallel to the horizontal plane (Fig.  
52 2). Changes in the magnetic flux density of the permanent magnet are measured while the table  
53 rotates 360°. In this case, I expect to see no changes in the magnetic fields strength, because the  
54 magnetic field distribution is symmetric at the table's plane according to Maxwell's equation in curved  
55 spacetime (Fig. 3).

56 In the second experiment, the plane of the rotating table is set to be vertical to the horizontal plane  
57 (Fig. 4). Again, changes in the magnetic flux density of the permanent magnet are measured while the  
58 table rotates 360°. In this case, I expect to see changes in the magnetic fields strength, because the  
59 magnetic field distribution does not maintain symmetry at the table's plane when the table rotates  
60 vertically. The magnetic field lines will be deflected to the center of Earth, as expected in Maxwell's  
61 equations in curved spacetime (Fig. 3).

62 Fig. 5 and Fig. 6 show the measured and the net magnitude of magnetic flux density from the  
63 experiments. The net magnitude of magnetic flux density represents the measured magnitude of  
64 magnet flux density excluding Earth's magnetic field (Fig. 7 and Fig. 8). I repeated both the first  
65 (horizontal rotation) and second (vertical rotation) experiments four times and recorded the average  
66 measures.

67 The results are consistent with my expectations. When the table rotates horizontally, the net  
68 magnetic flux density has the average value of 2599.85 and the standard deviation value of 1.24  
69 throughout the table's 360° rotation (Fig. 5). The changes in the net magnetic flux density during the  
70 rotation of table are negligible, as the small standard deviation implies.


71 On the other hand, the results from the vertical rotation are different. There are noticeable changes  
72 in the net magnetic flux density during the 360° rotation of the table. The net magnetic flux density has  
73 the average value of 1303.35 and the standard deviation of 2.96 (Fig. 6). The net magnetic flux  
74 density increases when the S pole faces toward the center of the Earth. The maximum change is  
75 about 10 during the rotation of table. The results are worth our attention, given that magnetic field  
76 distribution of a permanent magnet is widely known to be not changeable on the ground.

77 Here, I am analyzing and discussing the results from the vertically rotating experiments in details.  
78 When the magnetic sensor is located in zone A during the rotation of table (Fig. 6 and Fig. 9), the  
79 magnetic field lines are moving away from the sensor, because the strength of the magnetic field is  
80 decreasing. On the other hand, when the magnetic sensor is located in zone B during the rotation of  
81 table (Fig. 6 and Fig. 9), the magnetic field lines are moving closer to the sensor, because the strength  
82 of the magnetic field is increasing. The increase and decrease we see in zone A and B indicate that  
83 magnetic field distribution is no more symmetric. Rather, the magnetic field lines are deflected toward  
84 Earth's center as shown in Fig.9.

85


86

87  
88  
89  
90  
91




**Fig. 1. Neodymium permanent magnet and the magnetic sensor location.**

92  
93  
94  
95




**Fig. 2. In the first experiment, the permanent magnet and the magnetic sensor are fastened to the rotational table which is set to be parallel to the horizontal plane.**

96  
97  
98  
99



**Fig. 3. The conceptual diagram of magnetic field line in curved space both horizontally and vertically rotating table.**

100



101

102

**Fig. 4. In the second experiment, the permanent magnet and the magnetic sensor are fastened to the rotational table which is set to be vertical to the horizontal plane.**

103


104

105

106

107

108



109

110

**Fig. 5. Net magnetic flux density during 360 degree of horizontal rotation of table.**

111


112

113

114

115

116



117

118


119 **Fig. 6. Net magnetic flux density during 360 degree of vertical rotation of the table**

120

121

122

123



124 **Fig. 7. The observed Earth's magnetic flux density during 360 degree of horizontal rotation**  
 125 **of table.**

126

127

128

129

130


131

132

133

134

135



136

137


138

139

140

141

**Fig. 8. The observed Earth's magnetic flux density during 360 degree of vertical rotation of the table.**



142

143

144

145

146

**Fig. 9. The conceptual diagram of the deflected magnetic field lines during the 360 degree of vertical rotation of the table, which are obtained from analyzing the changes in the measured magnetic flux density.**

147

### 3. DISCUSSIONS

148

149

150

The results of the vertical rotation experiments show that the magnetic field distribution of a permanent magnet is changing with the rotation of the table. The solar winds ejected by Sun affects the magnetic field on the ground. However, the solar wind's effects are taken into account in the

151 measured Earth's magnetic flux density. The net magnetic field density excluding Earth's magnetic  
152 field is thought to be entirely that of the neodymium permanent magnet. The deflections observed in  
153 the vertical rotation experiments are caused by the curvature of the local spacetime by Earth. In other  
154 words, this study shows the possibility of using magnetic fields to measure the curvature of spacetime.  
155 Future studies could examine the vectors of deflections in the magnetic field lines in order to verify the  
156 geodetic effect. Compared to the gyroscopes, magnetic field has many advantages in measuring the  
157 curvature of spacetime as it is easier to handle and manufacture  
158

#### 159 **4. CONCLUSIONS**

160 Many studies [3-9] have verified the distortion in spacetime. However, not many tools are available  
161 to measure the curvature of spacetime, which is the key to describe gravity in the Einstein field  
162 equation. As shown in NASA's GP-B experiments [11-13], precise gyroscopes are the only available  
163 tools to measure the curvature of spacetime at this time. Yet, a precise gyroscope needs a lot of effort  
164 in handling and manufacturing. This study shows the possibility of using a magnetic field to measure  
165 the curvature of spacetime, which could be a more efficient tool than gyroscopes. In a series of  
166 experiments, it is found that there are noticeable changes in the magnetic field distribution of the  
167 permanent magnet fastened to rotating table. Magnetic field lines of a permanent magnet are  
168 deflected towards the Earth's center. The deflections are considered to be caused by the curvature of  
169 local spacetime of the Earth, as predicted in the Maxwell equations in curved spacetime. In the future,  
170 the results in this study should be compared with the theoretically obtained values of curvature of  
171 spacetime from Einstein field equation.  
172

#### 173 **REFERENCES**

- 174 [1] Einstein A., *Annalen der Physik* 1916; 354(7), 769.  
175 [2] Turner MSW, *International Journal of Modern Physics A* 2001; 17 (S1):180–196.  
176 [3] Dyson FW, Eddington AS, Davidson CA, *Philosophical Transactions of the Royal Society* 1920;  
177 220A: 291–333.  
178 [4] Kennefick D, *Physics Today* 2009: 37-42.  
179 [5] Holberg JB, *Journal for the History of Astronomy* 2010; 41(1): 41-64.  
180 [6] Einstein A, *Science* 1936; 84 (2188), 506–507.  
181 [7] Dyson FW, Eddington AS, Davidson CA, *Philosophical Transactions of the Royal Society* 1920;  
182 220A, 291–333.  
183 [8] van der Wel A, et al, *Astrophysical Journal Letters* 2013; 777: L17.  
184 [9] Wong K, et al, *Astrophysical Journal Letters* 2014; 789, L31.  
185 [10] Gould A., *Astrophysical Journal*, Part 2 – Letters 1993; 414(1): 37-40.  
186 [11] NASA - Gravity Probe B, [https://www.nasa.gov/mission\\_pages/gpb/](https://www.nasa.gov/mission_pages/gpb/).  
187 [12] NASA Gravity Probe Confirms Two Einstein Theories, <https://www.space.com/11570-nasa-gravity-probe-einstein-theory-relativity.html>.  
188 [13] Everitt CWF, et al, *Physical Review Letters* 2011;106, 221101.  
189 [14] Red'kov VM, Tokarevskaya NG, Bychkouskaya EM, *Nonlinear Dynamics and Applications* 2006;  
190 13: 207–228.  
191 [15] Lichnerowicz A, *Waves and Shock Waves in Curved Space-Time* (Springer, Netherlands); 1994.  
192 [16] Leonhardt U. Philbin T, *Geometry and Light: The Science of Invisibility* (Courier Corporation, U.S);  
193 2012.  
194 [17] [https://en.wikipedia.org/wiki/Maxwell%27s\\_equations\\_in\\_curved\\_spacetime](https://en.wikipedia.org/wiki/Maxwell%27s_equations_in_curved_spacetime)  
195