## Variation of the intensity of the gamma radiation in the period from May 28 to September 25 of 2017 in Sao José dos Campos, SP, Brazil Abstract

The variation of gamma radiation intensity between 200 keV to 10.0 MeV was measured, in the period from June 28 to September 25 of 2017. These measurements were taken 25 meters above ground level in a tower in São José dos Campos, SP, Brazil. A minute interval was used from each point of measurements. During this period, there were weak and moderate rains with net intensity of 27 mm. The variation in temperature during this period was between 15°C and 32°C, reproducing a desert-like climate. By monitoring the gamma radiation it was possible to observe the arrival of cold fronts from Southern Brazil and the (day/night) cycles due to the greater or lesser amount of radon gas present in the region. The main feature of this work is that it indicates the dynamics of gamma radiation in a simple way according to the variation of meteorological parameters in that location, which is very important for environmental studies in the tropical region.

# 1) Keywords: gamma radiation; cold fronts; radon gas; meteorological parameters; weak and moderate rains; day / night cycles

# 23 1. Introduction

At the (ground / air) interface of Earth's surface, the ionizing radiation is composed mainly of radon gas, the telluric radiation of the soil and the radiation of the primary and secondary cosmic rays. However, it is difficult to separate over time the intensity of the ionizing radiation emanating from each component as the energies overlap. The telluric radiation is given by <sup>238</sup>U, <sup>235</sup>U, <sup>40</sup>K, <sup>232</sup>Th and is constant for each region [1]. The radon gas that comes from the disintegration of <sup>238</sup>U of Earth's crust [2] into Ra-226 to Rn-222 arriving at the isotopes <sup>214</sup>Pb, <sup>214</sup>Po and <sup>214</sup>Bi giving alpha and gamma radiation. The primary cosmic radiation consists mainly of galactic and extragalactic protons and from the Sun with very high energy that interacts with Earth's atmosphere producing the EAS (Extensive Air Showers) [3]. The efficiency of this interaction is maximum when it occurs at altitudes between 15 and 17 km in the tropics, which form secondary cosmic rays with muonic, mesonic, and neutronic components that reach the Earth's surface in the region [4]. These radiations can cause health problems for the crew and passengers of civil/military aviation in this altitude that are present at the beginning of the stratosphere called Pfotzer maximum. However, this component contributes less to the concentration of ionizing radiation on the Earth's surface. Another possible source of ionizing radiation in the Earth's lower atmosphere is produced by electrical discharges between cloud-earth, earth-cloud and cloud-cloud. X-rays, gamma rays, neutrons and beta particles are all formed by the lightning cone [5]. Other sources of ionizing radiation are those produced in industry, medical, dental clinics and hospitals, but these radiations are mainly controlled in small and specific areas. 

# **2.** Materials and Methods

The gamma ray detector for the energy range of 200 keV to 10.0 MeV consists of a 3-inchby-3-inch-diameter and high sodium iodide scintillation crystal (3" x 3"), doped with thallium. This crystal is directly coupled to a photomultiplier (PM), which registers the pulses coming from the scintillator and with amplification and an analog digital converter (ADC) these digitized signals are recorded by a computer [6]. The detector was in a closed room and air exchange was ensured during the measurements. The experimental set (Figure 1) is located in the inner room of a tower (23°12'45.0"S 45°52'00.3"W), 25 meters high in relation to the ground surface.



Figure-1. View of the gamma scintillator associated with electronics and computer. Source: Project Atmosrad 2017 [9]

55

61

56 The scintillator coupled to photomultiplier is wrapped in a thin layer of aluminum to make it 57 portable. Scintillator and associated electronics were calibrated in terms of energy and counting 58 intensity per minute, at the laboratory of experimental teaching physics of ITA (Technological 59 Institute of Aeronautics), using radioactive sources and a spectral analyzer of counts versus energy in the range of 0.2 to 10 MeV (Million electron Volt), [7,8]. 60

#### **3. Results and Discussions** 62

63 Gamma radiation measurements were carried out during the period of June <mark>28</mark> to September 25 of 64 2017, in the inner room above the tower, seen in Figure 2 below. During the interval described above, 65 on the roof of the tower was placed the rain gauge that reported the intensity of rains in (mm / min). 66



Figure-2. Exterior view of the tower with the room 25 meters above the ground level. Source: Project Atmosrad 2017 [9]

Figure 3 shows the serie of measured gamma radiation intensity between June 28 to September 69 25 of 2017, with uninterrupted monitoring minute by minute during this net time.



### Figure-3. Monitoring of gamma radiation serie between the start time and 130 x 10<sup>3</sup> minutes after. Source: Project Atmosrad 2017 [9]



70 71

74 75 The smooth tool was used to transform data from minutes to days as seen on the green line. Analyzing 76 the dynamics of the radiation measurements, there are 4 large variations occurring in the whole period 77 analyzed. Between the beginning of monitoring and close to 70 x 10<sup>3</sup> minutes, the mean intensity of the 78 measured radiation was 37.5 x 10<sup>3</sup> (counts / minute). In this period it was possible to see small 79 variations which indicate passages of cold fronts without rain. See this dynamic in the expanded period 80 in Figure 4, taken from the graph in Figure 3.



Figure-4. Monitoring of gamma radiation between the start time and 70 x 10<sup>8</sup> minutes after. Source: Project Atmosrad 2017 [9]

82 83 84

85

81

Figure 5 shows the radiation monitoring between 70 to 80 x  $10^3$  minutes after the start of the measures. It was a rainy week with intensities varying according to Figure 5 and Figure 8.



Figure-5. Monitoring of gamma radiation between 70 X 10<sup>3</sup> and 80 x 10<sup>3</sup> minutes after. Source: Project Atmosrad 2017 [9]

86

87 In the beginning between 70 and 71 x  $10^3$  minutes there was intense rain, where the level of 88 radiation count reached the order of  $40 \times 10^3$  (counts / min). On the other days, there were always less 89 intense rains as shown by the gamma radiation peaks caused by the rains. In Figure 6, taken during the 90 measurement time of 80 to  $100 \times 10^3$  minutes, there are variations in the dynamics of the radiation with 91 passages of two cold fronts in the region, but without causing rains. However, the terrestrial surface 92 was wet and with very little exhalation of radon gas. The arrival of the cold front causes an increase in 93 the radiation due to the accumulation of radon gas that arrives with theses fronts.

3



Figure-6. Monitoring of gamma radiation between 80 x 10<sup>3</sup> and 100 x 10<sup>3</sup> minutes after. Source: Project Atmosrad 2017 [9]

95 In Figure 7, the gamma radiation intensity measured between the times of 100 to 130 x 10<sup>3</sup> minutes 96 with average intensity of 37.3 x 10<sup>3</sup> (counts / min), undergoes an influence of soil in the region that was 97 very dry and high pressure that was 970 mbar when local background value is 940 mbar. This occurs in 98 the afternoon where the temperature varied between 20<sup>o</sup> to 30 °C and the night between 12<sup>o</sup> to 20 °C. 99 This dynamics in temperature during this period facilitates greater and lesser exhalation of the radon 90 gas, as shown in the figure 7 below with cycles of exactly 24 hours. The two largest radiation peaks, 91 shown in Figure 7, are caused by heavy fog in the mornings of those days.



103 104 105

94

Figure 8 shows the rainfall spectrum in (mm / min), varying in time. During the whole period, only27 mm of net rain accumulated in the region in the course of a week.



108

Figure-8. Spectrum in time of the rain that occurred in the region during this period of measurements.
Source: Project Atmosrad 2017 [9]

Fig

### "Historically, the mean rainfall value for the period mentioned above is around 170 mm".

# 112 4. Conclusion

113

124

126 127

128

129

130

131 132 133

134 135

136 137

138

139 140

141

142 143

144

145

146 147

148 149

150

114 In the period of June to September of 2017, the intensity of rain was monitored every minute and in 115 the same place. In the same time the intensity of low energy gamma rays was also measured. The analysis show that during one single week of moderate and light rains, there was a noticeable increase in 116 the intensity of gamma rays. The net rainfall in the period measured was 27 mm scattered in time. 117 118 Analysis of all figures plotted in this work shows clearly the difference caused by the rains in the measurement of gamma rays. Also in this work, the perfect oscillation of the gamma rays (day / night) 119 in the dry period is evidenced, without clouds, fog or lightning. This oscillation is caused by the 120 121 exhalation of radon gas (Rn-222) in the region that is larger during the local solar zenith time with dry soil. The periodicity is partially or totally destroyed with the presence of fog, continuous rain and 122 123 drizzle in the region.

# 125 **References**

- [1] Gomes, Marcelo Pego; Martin, Inácio Malmonge; Silva, Franklin Andrade; Sismanoglu, Bogus Nubar; Monitoring of gamma radiations and meteorological parameters at ground level in São José dos Campos, Brazil. Impact: International Journal of Research in Engineering and Technology. Vol. 4, no. 8, 2016.
- [2] Martin, I.M.; Douglas Carlos Vilela ; Marcelo P. Gomes . Dynamics in Times of Ionizing Radiation and Rainfalls in Tropical Region of Brazil. Asian Review of Environmental and Earth Sciences, v. 4, p. 7-11, 2017.
- [3] Gusev, A. A., U. B. Jayanthi, I. M. Martin, G. I. Pugacheva, and W. N. Spjeldvik, Nuclear reactions on rarest atmosphere as a source of magnetospheric positron radiation belt, J. Geophys. Res., 106(A11), 26,111–26,116, 2001.
- [4] Gusev, Anatoli. A.; Martin, Inacio. M.; Alves, Mauro A.; de Abreu, Alessandro. J. . Simulation of the radiation fallout from gamma-ray measurements. Modeling Earth Systems and Environment, v. 1, p. 18, 2015.
- [5] Martin, Inácio Malmonge ; Gomes, Marcelo P . Intensity variation of gamma radiation on ground level interface in São josé dos Campos, SP, Brazil.. Environmental Science: An Indian Journal , v. 8, p. 79-82, 2013.
- [6] Silva, F. A.; Martin, I. M.; Gomes, M. P.; Monitoring of ionizing radiation and rain intensity during May to October 2015 in São José dos Campos, Brazil. Davi: Journal of Environmental Science and Engineering B 2 941-944, 2016.
- [7] Gomes, M. P.; Martin, I. M. Simultaneous Measurements of Rainfall Intensity, Low Energy Neutrons and Gamma Radiation in São Jose dos Campos, SP, Brazil. Journal of Environmental Science and Engineering , v. 09, p. 161-167, 2013.
- [8] Martin, I.M.; Marcelo P Gomes ; Bogos Nubar Sismanoglu ; Nicolas Cruvinel Lindo . Daily Variability of Radon Gas in Brazilian Tropics Near Ground Level Surface. Journal of Environmental Science and Engineering , v. A4, p. 516-521, 2015.
- [9] Project Atmosrad. Research Group of CNPq <u>http://dgp.cnpq.br/dgp/espelhogrupo/4663704786429697</u>.
- 151 152 153