

Adiabatic Lapse Rate of Water Is Negative Due to Its Negative Compressibility

ABSTRACT

When the pressure of a fluid changes without the addition of heat, the temperature of the fluid changes; the rate at which the temperature changes with pressure is called the adiabatic lapse rate. According to thermodynamic equations, the adiabatic lapse rate is positive if the thermal expansion coefficient is positive, and negative if this coefficient is negative. Experiments show, however, that the adiabatic lapse rate is also positive for substances with negative thermal expansion, although for water it is negative when it has negative thermal expansion. In the present paper, a theory is developed which shows that the adiabatic lapse rate must always be positive, but is negative for water because it has negative compressibility in that temperature–pressure region. Numerous substances with negative compressibility have already been identified.

Keywords: adiabatic lapse rate; adiabatic temperature gradient; ocean thermodynamics; water; negative thermal expansion; negative compressibility; internal energy; enthalpy

1. INTRODUCTION

The adiabatic lapse rate of water is the rate at which its temperature changes with pressure at constant entropy S , and salinity. According to conventional theory, the adiabatic lapse rate $(\partial T/\partial P)_S$ is positive if the thermal expansion coefficient α is positive, and negative for negative values of α [1–4]:

$$\left(\frac{\partial T}{\partial P}\right)_S = \frac{\alpha TV}{C_P}, \quad (1)$$

where T is the absolute temperature, P is the pressure, V is the volume, and C_P is the isobaric heat capacity. In the present paper, this phenomenon is explained for water with zero salinity. Taking the salinity into account does not change the conclusions of the theory, because the salinity of water is not of primary importance to this problem [1]. However, the derivative of the internal energy U , $(\partial U/\partial P)_S$, is always positive [1,5]:

$$\left(\frac{\partial U}{\partial P}\right)_S = \frac{\beta C_V PV}{C_P}, \quad (2)$$

where C_V is the isochoric heat capacity, and $\beta = -1/V(\partial V/\partial P)_T$ is the isothermal compressibility. The increase in the internal energy dU due to pressure is equal to $C_S dT$, where C_S is the heat capacity at constant entropy, $(\partial U/\partial T)_S$, and dT is the increase in temperature due to the pressure [5]:

$$\left(\frac{\partial U}{\partial T}\right)_S = \frac{\beta C_V P}{\alpha T}. \quad (3)$$

Therefore, for negative values of α , C_S and dT must be negative. In this paper it is shown that C_S and dT must be positive regardless of the sign of α ; however, if dT is negative for water, then this is due to its negative compressibility. These results are confirmed by experiment.

2. THEORY

If $C_S < 0$ at $\alpha < 0$, then by virtue of Eqs. (1)–(3), the temperature decreases when the internal energy increases. This cannot be true for a substance with positive compressibility, since if a quantity of heat δQ is introduced into water at $\alpha < 0$, both its internal energy and its temperature increase. According to Joule's principle of equivalence of heat and work, it makes no difference whether the internal energy is increased by introducing a quantity of heat or by compression.

The internal energy is the sum of the kinetic and potential energies:

$$U = K + \Pi. \quad (4)$$

In accordance with traditional theory, for substances with negative thermal expansion, when internal energy increases in Eq. (4), kinetic energy decreases and potential energy increases [1]. Under compression, the internal energy changes by dU :

$$dU = \left(\frac{\partial U}{\partial K} \right)_{\Pi} dK + \left(\frac{\partial U}{\partial \Pi} \right)_K d\Pi. \quad (5)$$

The first differential on the right-hand side of Eq. (5) always increases with U . This means that the temperature also increases because the derivative is positive and is equal to 1. This derivative is also greater than zero at $\Pi \approx \text{CONSTANT}$, due to the continuity of the energy function. Therefore, the derivative $(\partial U / \partial K)_S$ is also positive in the same interval where $\Pi \approx \text{CONSTANT}$ (if it were negative, then the energy function would not be continuous). Let us suppose that outside that interval, $(\partial U / \partial K)_S$ is negative. This means that at the border of this interval, this derivative equals zero, and therefore $(\partial U / \partial T)_S$ is zero. This cannot happen, because in this case it follows from Eq. (3) that α tends to infinity; hence from Eq. (1) it follows that $(\partial T / \partial P)_S$ tends to infinity. This contradicts the definition: α is finite. The conclusion can be drawn that K and T always increase with U , independent of the sign of α .

It can also be argued that under compression, internal energy increases; part of it becomes a change in kinetic energy, and another part becomes a change in potential energy:

$$dU = dU_1 + dU_2 = \frac{dK}{dU_1} dU_1 + \frac{d\Pi}{dU_2} dU_2.$$

Here, dU_1 and dU_2 are greater than zero and less than dU , and hence dK and dT are always greater than zero. Thus, heat capacity C_S cannot be negative due to the reasons given above.

In the traditional equation for adiabatic compression, $(\partial U / \partial T)_S$ is positive for positive values of α , and negative for negative values of α (Eq. (3)). For some pressures and temperatures, however, α is equal to zero, and hence this derivative is infinite. This is an unphysical result: a small change in the temperature causes an infinite increase in the internal energy. It also follows from Eq. (1) that

80

$$\left(\frac{\partial P}{\partial T} \right)_S = \frac{C_p}{\alpha TV} .$$

82 One can see that for $\alpha = 0$, a small change in the temperature causes an infinite increase in the
83 pressure, and this is unrealistic. The reason for this discrepancy is that the first law of
84 thermodynamics for the *heat exchange* is the following:

$$\delta Q = dU + PdV . \quad (6)$$

87
88 A quantity of heat δQ is introduced into the system, and this becomes a change in the internal energy
89 and work produced by the system. However, for the heating of a substance *by compression*, the first
90 law of thermodynamics cannot be derived from Eq. (6) and must be derived independently [6–9]:

$$PdV = -dU . \quad (7)$$

93
94 The substance is compressed by the work PdV , and this results in a change in internal energy. Heat
95 losses are not taken into account. The sign before dU in Eq. (7) cannot be determined theoretically; it
96 can only be determined by experiment. Equations (1), (3), (9) and (10) were derived for heat
97 exchange, but not for compression, and cannot be used for the description of adiabatic compression.

98
99 Numerous experiments on substances with negative thermal expansion have been performed, and
100 these have shown that heat is expressed under compression [7,9–11]. The same experiments with
101 water showed, however, that heat is absorbed under compression in the temperature–pressure
102 interval where water has negative thermal expansion [12–14]. The following explanation can be
103 proposed. Recently, many substances with negative compressibility have been discovered (i.e. they
104 expand under compression) [15–23]. Some studies have supposed that water has negative
105 compressibility when it has negative thermal expansion (for pressures of less than 30–60 MPa)
106 [7,24,25]. This may be the reason why its temperature decreases as the pressure increases.

107
108 It is necessary to note that Fine and Millero determined the compressibility of water from 273 to 373 K
109 and 0 to 100 MPa from sound velocity measurements and found it to be always positive [26]. This was
110 an indirect determination using the following equation:

$$\beta = \frac{V}{c^2} + \frac{T\alpha^2 V}{C_p} \quad (8)$$

113
114 where c is the velocity of sound. However, according to this equation the compressibility of all
115 substances is always positive, which is not the case. Equation (8) is derived from the Mayer's relation
116 and in [27] it was shown that the derivation of the Mayer's relation is wrong.

117 3. DISCUSSION

118
119
120 It is interesting to consider how enthalpy (heat content) H changes under adiabatic pressure in
121 traditional theory [5]:

$$\left(\frac{\partial H}{\partial P} \right)_S = V \quad (9)$$

124
125 and

$$\left(\frac{\partial H}{\partial T} \right)_S = \frac{C_p}{\alpha T} . \quad (10)$$

128

At $\alpha < 0$, enthalpy decreases with temperature. This contradicts the physical sense of the process: if the heat content is increased, the temperature must also increase.

Another interesting result follows from Eq. (10) and $(\partial H / \partial T)_P = C_P$. Due to the continuity of H ,

$$\text{sign} \left(\frac{\partial H}{\partial T} \right)_{P \approx \text{CONST}} > 0.$$

Therefore, in this pressure interval

$$\text{sign} \left[\left(\frac{\partial H}{\partial T} \right)_S \right]_{P \approx \text{CONST}} > 0.$$

Let us suppose that outside of this pressure interval where $P \approx \text{CONSTANT}$, the derivative $(\partial H / \partial T)_S$ is less than zero. This means that at the border of this interval, this derivative is equal to zero, and from Eq. (10) it follows that α tends to infinity. This cannot be true, so it must be concluded that $(\partial H / \partial T)_S$ is always positive. As mentioned in the previous section, Eqs. (9) and (10) cannot be used for the description of mechanical compression.

4. CONCLUSION

It has been shown that the traditional thermodynamic equations cannot be used to describe the adiabatic compression of substances because they are derived from the equation which describes heat exchange, Eq. (6). Adiabatic compression is a different process from heat exchange and must be described by other equations. The traditional equations predict that substances with negative thermal expansion absorb heat under compression, while numerous experiments show that they express heat. However, water does absorb heat when it has negative thermal expansion. A possible reason could be that water has negative compressibility in this region, as proposed earlier by the author. As many substances with negative compressibility have recently been found, this explanation appears to be plausible. Direct measurements of compressibility of water were performed at pressures higher than 60 MPa, where thermal expansion and compressibility are positive. Negative compressibility is supposed to take place at pressures lower than 30 MPa for $T = 270$ K and lower than 60 MPa for $T = 260$ K [25]; therefore, precise experiments in this low-pressure region are necessary. Taking the salinity of water into account does not change the results of the theory.

REFERENCES

1. McDougall TJ, Feistel R. What causes the adiabatic lapse rate? Deep-Sea Res, Pt I. 2003;50(12): 1523–35.
DOI: 10.1016/j.dsr.2003.09.007
2. Boehrer B, Schultze M. Stratification of lakes. Rev Geophys. 2008;46(2):RG2005.
DOI: 10.1029/2006RG000210
3. Björnbohm P. Temperature lapse rates at restricted thermodynamic equilibrium in the Earth system. Dynam Atmos Ocean. 2015;69:26–36.
DOI: 10.1016/j.dynatmoce.2014.10.001
4. Tailleux R. On the validity of single-parcel energetics to assess the importance of internal energy and compressibility effects in stratified fluids. J Fluid Mech. 2015;767:R2.
DOI: 10.1017/jfm.2015.77
5. Karapetyants M.Kh. Himicheskaja termodinamika. Moscow: Vysshaja Shkola; 1974. pp. 108–110. English translation: Karapetyants MKh. Chemical thermodynamics. (Translated by Leib G). Moscow: Mir; 1978.
6. Stepanov IA. The first law of thermodynamics for auxetic materials. J Non-Cryst Solids. 2013;367: 51–2.

181 DOI: 10.1016/j.jnoncrsol.2013.02.017
 182 7. Stepanov IA. Compression of substances with negative thermal expansion. *Mater Lett.* 2015;161:
 183 365–7. And references therein.
 184 DOI: j.matlet.2015.08.150
 185 8. Stepanov IA. The isochoric heat capacity of ZrW_2O_8 and $Sc_2W_3O_{12}$ is greater than their isobaric
 186 one. *Mater Lett.* 2016;177:112–5.
 187 DOI: j.matlet.2016.04.153
 188 9. Stepanov IA. Heating by heat exchange and by compression – asymmetry in the equations. *JP J*
 189 *Heat Mass Transf.* 2017;14(4):579–85.
 190 DOI: 10.17654/HM014040579
 191 10. Jayaraman A, Dernier P, Longinotti LD. Study of the valence transition in SmS induced by
 192 alloying, temperature, and pressure. *Phys Rev B.* 1975;11(8):2783–94.
 193 11. Stepanov IA. Thermodynamics of iodine under high pressure at low temperatures. *Physica B.*
 194 2004;349(1–4):251–3. And references therein.
 195 DOI: 10.1016/j.physb.2004.03.177
 196 12. Caldwell DR. The maximum density points of pure and saline water. *Deep Sea Res.*
 197 1978;25(2):175–81.
 198 DOI: 10.1016/0146-6291(78)90005-X
 199 13. Caldwell DR, Eide SA. Adiabatic temperature gradient and potential temperature correction in
 200 pure and saline water: an experimental determination. *Deep Sea Res.* 1980;27A(1):71–8.
 201 DOI: 10.1016/0198-0149(80)90073-4
 202 14. Rögner H, Soll P. Beitrag zum Thermodynamischen Messverfahren – Der Isentrope Temperatur-
 203 Druck-Koeffizient des Wassers. *Brennstoff-Wärme-Kraft.* 1980;32(10):472–8. German.
 204 15. Baughman RH, Stafström S, Cui C, Dantas SO. Materials with negative compressibilities in one or
 205 more dimensions. *Science.* 1998;279(5356):1522–4.
 206 DOI: 10.1126/science.279.5356.1522
 207 16. Kornblatt JA. Materials with negative compressibilities. *Science.* 1998;281(5374):143a.
 208 DOI: 10.1126/science.281.5374.143a
 209 17. Baughman RH. Avoiding the shrink. *Nature.* 2003;425(6959):667.
 210 DOI: 10.1038/425667a
 211 18. Moore B, Jaglinski T, Stone DS, Lakes RSs, Negative incremental bulk modulus in foams. *Phil*
 212 *Mag Lett.* 2006;86(10):651–9.
 213 DOI: 10.1080/09500830600957340
 214 19. Moore B, Jaglinski T, Stone DS, Lakes RS. On the bulk modulus of open cell foams. *Cell Polym.*
 215 2007;26(1):1–10.
 216 20. Lakes R, Wojciechowski KW. Negative compressibility, negative Poisson's ratio, and stability.
 217 *Phys Status Solidi B.* 2008;245(3):545–51.
 218 DOI: 10.1002/pssb.200777708
 219 21. Rozsa VF, Strobel TA. Triple guest occupancy and negative compressibility in hydrogen-loaded β -
 220 hydroquinone clathrate. *J Phys Chem Lett.* 2014;5(11):1880–4.
 221 DOI: 10.1021/jz5005895
 222 22. Rodney D, Gadot B, Martinez OR, Roscoat SR, Orgéas L. Reversible dilatancy in entangled
 223 single-wire materials. *Nat Mater.* 2016;15(1):72–8.
 224 DOI:10.1038/nmat4429
 225 23. Baughman RH, Fonseca AF. Architected materials: straining to expand entanglements. *Nat*
 226 *Mater.* 2016;15(1):7–8.
 227 DOI: 10.1038/nmat4436
 228 24. Stepanov IA. Thermodynamics of substances with negative thermal expansion and negative
 229 compressibility. *J Non-Cryst Solids.* 2010;356(23–24):1168–72.
 230 DOI: 10.1016/j.jnoncrsol.2010.03.013
 231 25. Stepanov I.A. Anomalous dependence of the heat capacity of supercooled water on pressure and
 232 temperature. *Results Phys.* 2014;4:28–30. (Free online).
 233 DOI: 10.1016/j.rinp.2014.02.003
 234 26. Fine RA, Millero FJ. Compressibility of water as a function of temperature and pressure. *J Chem*
 235 *Phys.* 1973;59(10):5529–36
 236 DOI: 10.1063/1.1679903

237 27. Stepanov IA. Exact calculation of the internal energy of the ideal gas in statistical mechanics.
238 Phys Sci Int J. 2017;14(2):1–5.
239 DOI: 10.9734/PSIJ/2017/32822