

# Phytochemical Analysis Of n- Hexane and Ethylacetate Extracts Of *Diodia scandens* Sw and Spectroscopic Identification Of an Omega- 6 Fatty acid and a Glyceryl trilinoleate.

## ABSTRACT

Traditional Medicine Practitioners in South East Nigeria use *Diodia scandens* Swartz for the treatment of different diseases such as venereal, cutaneous, subcutaneous, fungal infections, inflammatory, antiabortion, fertility management, antibiotic and as antidotes for snake bite. Cold maceration of n-hexane and Ethyl acetate solvents extracts for 48 hours yielded brownish oily solids which were subjected to phytochemical analysis. Column Chromatographic fractionation of the crude extracts using n-hexane: ethyl acetate in different v/v ratio of 95:05, 90:10, 80:20; down to 100 ml of ethyl acetate yielded different fractions. Eluent fractions with same or similar  $R_f$  values were pooled together and evaporated. These were subjected to spectroscopic analysis. They were labelled as CHDS11 and CHDS12 from the 95:05 n-hexane: ethyl acetate solvents mixture. The result of the Phytochemical analysis of the crude n-hexane extract showed the presence of steroids, glycoside, alkaloids, and saponins. Crude ethyl acetate extract showed the presence of steroid and saponin. The fraction CHDS11, was identified as cis,cis-9, 12-octadecadienoic acid (linoleic acid). The  $^1\text{H}$  NMR chemical shift of 5.05(t) confirmed the hydrogens attached to the double bonds at C-9 and 10; C-12 and 13 adjacent to the methylene group; CHDS12 was identified as 1, 2, 3-propanetriyltris (cis, cis - 9, 12-octadecadienoate), known as glyceryl trilinoleate with the chemical shifts 5.28 ppm showing the vinyl protons at C-9 and 10; C-12 and 13; 5.19 ppm showed protons in between trilinoleate chains. The presence of these compounds justified the use of this plant for treating arthritis and the diseases earlier mentioned.. To the best of our knowledge, this is the first time these compounds are isolated from this plant.

**Keywords:** *Diodia scandens*, n-hexane, ethylacetate, phytochemical analysis,  $^1\text{H}$  NMR.

## INTRODUCTION

Human beings from antiquity depended on plants either directly for foods and beverages, or indirectly as feed for animals or the flavoring of foods. Plants are also the source of beverages produced either by infusion, such as coffee and tea; by fermentation, such as beer and wine; or by distillation, such as whisky, vodka, rum, and other alcoholic spirits [1].

Plants are the source of many natural products such as essential oils, natural dyes, pigments, waxes, resins, tannins, alkaloids, amber and cork. Products derived from plants include soaps, shampoos, perfumes, cosmetics, paint, varnish, turpentine, rubber, latex, lubricants, linoleum, plastics, inks, and gums. Renewable fuels from plants include firewood, peat and many other biofuels. Coal and petroleum are fossil fuels derived from the remains of plants. Olive oil has been used in lamps for centuries to provide illumination [1].

Man has been engaged in the use of the plants and their different parts for treatment of various ailments. Generally, plants have been used throughout the world in folk medicine and as local cure for common ailments. Medicinal plants in particular have been in use for centuries as remedies for human diseases like malaria, dysentary, diarrhoea, typhoid, arthritis, infertility etc because they contain components of therapeutic value [2]. Folk medicine gave rise to traditional system of medicine in various diseases.

Medicinal herb is considered to be a chemical factory as it contains multitude of chemical compounds like alkaloids, glycosides, saponins, resins, oleoresins, sesquiterpene, lactones and oils (essential and fixed). India has one of the oldest, richest and most diverse cultural traditions associated with the use of medicinal plants. According to an estimate, 120 or so plant based drugs prescribed for use throughout the world come from just 95 plant species [3]. Natural antimicrobials can be derived from plants, animal tissues and microorganisms. The shortcoming of the drugs available today propelled the discovery of new pharmacotherapeutic agents from medicinal plant research [3]. The aim of this research is to carry out phytochemical analysis of n-hexane and ethylacetate extracts of *Diodia scandens* Sw and spectroscopic identification of an omega-6 fatty acid and a glyceryl trilinoleate.

## DESCRIPTION OF PLANT

*Diodia scandens* Sw (Rubiaceae) is an evergreen perennial herb, which has an alternate leaf arrangement, petiole is present. It has compound leaves, ovate to lanceolate in shape, replete venation, entire in margin, its apex is acute, its base is cuneate, it has glabrous surface and its texture is chartaceous. *Diodia scandens* Sw has a dark green coloration, tasteless, odorless and has solitary inflorescence. It is a straggling herb, which has been in use in the western Africa system of medicine. It has enormous usefulness and importance; whole parts of the plants are useful in curing various ailments [4].

The plant's medicinal value includes its use as antidotes, painkiller, treatment of venereal diseases and cutaneous and subcutaneous fungal infections. The different parts of the plants- sap, leaf, stem and root, are used for various medical purposes. The leaf is used for treating arthritis, rheumatism, cutaneous and subcutaneous parasitic infection, diarrhoea, dysentery and antiabortifacients. The leaf plus roots are used for dropsy, swellings, oedema, and gout and as lactation stimulants; while the sap is used for treating ear infections, paralysis, epilepsy, convulsions, spasm and pulmonary troubles [4]. The whole plant of *D. scandens* Sw is used for treating fibroid and uterine disorder [5].

## MATERIALS AND METHODS

### MATERIALS

The solid compounds used in this work were purified and solvents redistilled before use. The spectroscopic equipment were;

Infrared spectrometer, transmittance 4000-650.

Ultraviolet/ Visible spectrometer, 4.20(468).

Nuclear Magnetic Resonance, 400 MHz

### METHODS

## Sample collection and preparation.

*Diodia scandens* Sw was collected at St. Mary's Pro-Cathedral Parish Udi and the plant was identified by a qualified taxonomist, Prof. J C. Okafor, at No. 7 Dona drive, Off Ihiala Street, Independence Layout Enugu, Enugu state, Nigeria.

The plant leaves were washed and air dried.. The dried leaves were ground with a mechanical grinder. It was stored in large sample bottle and labeled.

## Extraction of the plant material.

Using cold maceration method n-hexane and ethylacetate were used. The pulverised sample (200 g) was macerated in 500 mL of n-hexane, stirred vigorously and was left for 48 hours. It was filtered and the filtrate was allowed to dry at room temperature .. This was repeated using ethylacetate.

## Preparation of reagents

**Meyer's reagent:** Meyer's reagent was prepared by adding 1.3 g of mercuric iodide in 10 mL of water and 5.0 g of potassium iodide in 20 mL of water and making it up to 100 mL solution in a flask [6].

**Marquis reagent:** Marquis reagent was prepared by mixing concentrated tetraoxosulphate (VI) acid and formalin in the ratio of 10:1 v/v respectively [6].

**Draggendorf's reagent:** Draggendorf's reagent was prepared by dissolving 0.6 g bismuth subnitrate in 2 mL concentrated hydrochloric acid and 10 mL of water and was mixed with 6 g of potassium iodide in 10 mL of water. Then 7 mL of concentrated hydrochloric acid 15 mL of water was added and the whole was diluted with 400 mL of water [6].

## PHYTOCHEMICAL ANALYSIS [6], [7]

## COLUMN CHROMATOGRAPHY

The crude n- hexane extract was column chromatographed on silica gel adsorbent using n-hexane: ethylacetate v/v in the ratio of 95:05, 90:10, 80:20,70:30 up to 100 mL of ethyl acetate. Fractions with same or similar  $R_f$  values were pooled together. Fractions CHDS11 and CHDS12 were collected from 95:05 n-hexane: ethyl acetate.

## RESULTS AND DISCUSSION

### Mass and percentage yield of crude extracts

Table .1 showed the mass and percentage yield of crude extracts from the solvents. Ethylacetate had a mass yield of 7.15 g (3.58%) and n-hexane had weight 3.18 g (1.59%). From the result, a more polar solvent ethylacetate extracted more than a non polar solvent. This shows that polar solvents are better extraction solvent than non-polar solvents

**TABLE 1: MASS and percentage yield of the crude extracts from the solvents**

Extracting solvents	Mass of crude extract	Percentage yield
---------------------	-----------------------	------------------

	(gram)	
Ds ethylacetate	7.15	3.58
Ds n-hexane	3.18	1.59

Ethylacetate extracted a higher percentage mass than n-hexane

#### Phytochemical analysis of the crude extracts of *D. scandens Sw*

Table 2. showed the phytochemical result for crude extract of *D. scandens Sw*. The phytochemical analysis of n-Hexane crude extract showed that glycoside, steroid, saponin and alkaloid were present. In the ethyl acetate, steroid and saponin were present. Properties of saponin containing herbs are many and varied and may include diuretic, expectorant, anti-catarthal, anti-inflammatory, antispasmodic, aphrodisiac, antioxidant, emmenagogue, cardiac stimulant, hormone modulating, hepatoprotective, and adrenal adaptogenic effects [8]. Steroids have been reported to have antibacterial properties and they are very important compounds especially due to their relationship with compounds such as sex hormones. Alkaloids have been associated with medicinal uses for centuries and one of their common biological properties is their cytotoxicity. Several workers have reported the analgesic, antispasmodic and antibacterial properties of alkaloids. Glycosides are known to lower the blood pressure according to many reports [8]. The results obtained in this study thus suggest that the identified phytochemical compounds may be the bioactive constituents, and this plant is proving to be an increasingly valuable reservoir of bioactive compounds of substantial medicinal merit.

**TABLE2: PHYTOCHEMICAL RESULT FOR CRUDE EXTRACT OF *Diodia scandens Sw***

	Ds n-Hexane	Ds ethylacetate
<b>Glycoside</b>	+	—
<b>Steroid</b>		
i) Salkowski test	+	+
ii) Lieberman-Buchard test	+	+
iii) Formaldehyde test	+	+
<b>Saponin</b>		
i) Emulsion test	+	+
ii) Frothing test	—	+
<b>Alkaloids</b>		
i) Maquis test	+	—
ii) Meyer's test	+	—
iii) Dragendorff's test	—	—
<b>Tannins</b>		
i) Test with 5% FeCl <sub>3</sub>	—	—
ii) Test with 10% KOH	—	—
<b>Phlobatanintest</b>	—	—
<b>Anthraquinonetest</b>	—	—
<b>Flavanoids</b>		
Test with concentrated H <sub>2</sub> SO <sub>4</sub>	—	—
Test with 10% NaOH	—	—
Test with Mg - HCl	—	—

**Key:** + = present, - = absent.

#### SPECTROSCOPIC ANALYSIS OF THE PURE SAMPLES

Table 3: Pure samples were analysed for structural elucidation using  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ).  
Compounds were got from CHDS11, and CHDS12.  
The compound CHDS11 was oily solid and has a wine red colour. Fraction CHDS12 was an oily solid.

**Table 3: SPECTROSCOPIC ANALYSIS OF THE PURE SAMPLES**

Fraction	Colour and nature
CH DS <sub>11</sub>	Wine red oil
CH DS <sub>12</sub>	Orange oil

Isolated fractions and their colours.

### **$^1\text{H}$ NMR spectrum of CHDS11 as (cis,cis-9,12-octadecadienoic acid) in $\text{CDCl}_3$ compared to literature**

Table 3 showed the experimental chemical shifts and multiplicity of CHDS11 compared to literature values..

The chemical shift at 0.8(qd) showed the presence of methyl group at C-18; 5.05(t) confirms the hydrogens attached to the 2 double bonds at C-9 and 10; C-12 and 13 of the compound.

The chemical shift 2.3 (t) at C-11 shows the methylene hydrogen linking the two double bonds, 1.97 (t) at C-2 indicates the methylene hydrogen attached to carboxylic acid group. The chemical shift 1.61(s) at C-14 shows methylene hydrogen preceding the vinyl carbons. The chemical shift 1.20 (m) indicates methylene groups in between the  $\text{CH}_2$  following the vinyl carbon and the carboxylic acid group and the one (3( $\text{CH}_2$ )) in between the methyl group and the  $-\text{CH}_2-$  group attached to vinyl carbon, given a total of 14 hydrogens. These chemical shifts as compared with literature confirm the presence of linoleic acid in the sample from *D. scandens* Sw.

The presence of fatty acids such as linoleic acid which balances female reproductive hormones and also helps to lubricate the mucous membrane.


Growing evidence indicates that the design and delivery of supplemental fatty acids to the lower gut may target reproductive tissues to improve reproductive function and fertility. Improvement in embryo survival may be associated with suppression of uterine prostaglandin secretion via linoleic acid or other longer chain unsaturated fatty acids. Changes in follicular dynamics can be affected by fat supplementation and may lead to a more fertile ovulation. This improvement may be due to alterations in metabolic hormones and growth hormone or hormonal clearance [9]. Thus, the fatty acid present in *D. scandens* Sw must be responsible for reduction of swelling and nourishing of the uterine wall.

**Table 4: Experimental chemical shifts and multiplicity of CHDS11 compared to literature.**


Proton	Experimental chemical shift $\delta$ and multiplicity	SDBS literature	Anatoli et al Literature
CH – 9,10,12,13	5.05 (t)	5.58-5.12	5.35 (m)
CH – 8,14	2.23 (t)	2.764	2.77 (t)
CH <sub>2</sub> – 2	1.97 (t)	2.33	2.33
CH <sub>2</sub> – 3	1.54 (d)	1.63	1.61
CH – 11	1.61 (s)	2.02	2.07 (q)
CH – 15	1.20 (m)	-	1.30 (m)
CH – 16	1.20 (m)	-	1.30 (m)
CH – 17	1.20 (m)	1.10–1.53	1.31(m)
CH <sub>2</sub> – 4,5,6,7	1.20 (m)	1.10 – 1.53	1.33(m)
CH <sub>3</sub> – 18	0.80 (qd)	0.9	0.9 (t)

Experimental NMR data showing the protons and chemical shifts in ppm

**Figure 1: CIS, CIS-9,12-OCTADECADIENOIC ACID, C<sub>18</sub>H<sub>32</sub>O<sub>2</sub> (linoleic acid)(CHDS11)**



**<sup>1</sup>H NMR spectrum of cis,cis-9,12-octadecadienoic acid**



# **<sup>1</sup>H NMR spectrum of CHDS12 as 1,2,3-propanetriyltris(cis,cis-9,12-octadecadienoate) in CDCl<sub>3</sub> compared to literature.**

Table 4: showed the experimental chemical shift and multiplicity of CHDS12. It was orange oil and gave an  $R_f$  value of 0.68.

The chemical shifts 5.28(m) indicates vinyl protons at C – 9, 12 in the compound. The chemical shift 5.19(t) shows protons in between trilinoleoate chains. The chemical shift 4.22 (dd) and 4.07 (dd) indicates the proton glyceride group. The chemical shift 2.72 (dq) shows methylene protons in between the double bonds. The chemical shift 2.24 (td) shows – CH<sub>2</sub>– group following the ester group. The three chemical shifts 1.99 (m), 1.54 (d) and 1.21 (m) are methylene group in between the vinyl carbon and the – CH<sub>2</sub>– group attached to the ester carbon. 0.81 (td) shows a methyl group. The chemical shifts as compared to literature confirm the presence of triglyceride, 1, 2, 3 – propanetriyltris (cis, cis – 9, 12 – octadecadienoate) is present in *D. scandens* Sw.

This compound 1,2,3-propanetriyltris(cis,cis-9,12-octadecadienoate) increases the blood permeability in the brain. The blood brain barrier forms a structural and functional barrier between the blood circulation and brain parenchyma, regulates the transport of molecules, and prevents blood cells accessing brain tissues. Triolein emulsion transiently increased vascular permeability with interstitial edema in the cat brain when administrated via the carotid artery [10]. Triglycerides are esters of glycerol combined with three chains of fatty acids. Elevated triglyceride are strong indicator of biliary function, fat metabolism, the function of the liver and hereditary. There is generally a sugar-handling issue with elevated triglyceride or adult onset diabetes. Decrease triglyceride suggests poor release of fatty acid,


endocrine hyperfunction, and immune problem [11]. Triglyceride insulates the body from extreme temperature changes. They absorb and transport vitamin A, D, E and K through the blood cell [12].

**Table 5: Experimental chemical shift of CHDS12 compared to literature**

Proton	Experimental Chemical shift $\delta$ and Multiplicity	SDBS Literature
CH=CH9,10,12,13	5.28 (m)	5.35
H-C-O-2	5.19 (t)	5.27
CH <sub>2</sub> O-1 <sup>1</sup>	4.22 (dd)	4.295
CH <sub>2</sub> O-1 <sup>2</sup>	4.07 (dd)	4.147
CH <sub>2</sub> -11	2.72 (dq)	2.769
CH <sub>2</sub> -2	2.24 (td)	2.315
CH <sub>2</sub> -8,14	1.99 (m)	2.049
CH <sub>2</sub> -3	1.54 (d)	1.61
CH <sub>2</sub> --7,15	1.21(m)	1.36
CH <sub>2</sub> -4,5,6,16,17	1.21 (m)	1.30
CH <sub>3</sub> -18	0.81 (td)	0.889


Experimental NMR data showing the protons and chemical shifts in ppm.





**Figure 2: 1,2,3-propanetriyltris(cis,cis-9,12-octadecadienoate)**

**<sup>1</sup>H NMR SPECTRUM OF 1,2,3-propanetriyltris(cis,cis-9,12-octadecadienoate)**



## CONCLUSIONS

The phytochemical analysis carried out showed the presence of steroids, glycosides, alkaloids, saponins, in n-hexane leaves crude extract, steroids, and saponins in the crude ethylacetate extract and steroids, saponins, alkaloids. The n-hexane crude extract was fractionated and two pure samples were obtained and were labeled CHDS11, CHDS12.

The  $^1\text{H}$  NMR spectral revealed that the sample labeled CHDS11 contained cis,cis-9,12-octadecadienoic acid, CHDS12 contained 1,2,3-propaneteryltris(cis,cis-9,12-octadecadienoate). The presence of these compounds has shown that the traditional use of the plant could be correct.

1.

- 292 1. Field, C.B., Behrenfeld, M.J., Randerson, J.T and Falkowski, P. (1998). Primary  
293 production of the biosphere: Integrating terrestrial and oceanic components. pp 237-238.
- 294 2. Okoli, R.I., Turay, A.A., Mensah, J.K. and Aigbe, A.O (2009). Phytochemical and  
295 Antimicrobial properties of four herbs from Edo State, Nigeria. Report and Opinion 1(5): 67  
296 – 73.
- 297 3.Suman, R. K., Venkateshwer, G., Samuel, G, and Gangadhar S. (2013), Phytochemical  
298 screening of some compounds from leaf plant extracts of *Holoptela integrifolia* and *Celestus*  
299 *emarginata* used by Gondu tribes at Adilabad District, Andhrapradesh, India. 2: 65 - 70.
- 300 4. Essiette, U.A., Bala, D.N and Agbakalu, J. A (2010). Pharmacognostic studies of the leaves  
301 and stem of *Diodia Scandens* Sw in Nigeria. Scholar research library, Archives of Applied  
302 Science Research. 2(5): 184 – 198.
- 303 5. Okafor, J. C (2013). Tropical Plants in Health Care Delivery in Nigeria Bodija Ibadan pp 7  
304 – 58.
- 305 6. Akpuaka, M.U (2009). Essentials of Natural Products Chemistry, Mason, Enugu pp. 34 –  
306 64.
- 307 7. Sofowora, A. (1993). Medicinal Plants and Traditional Medicine in Africa, 2nd Edn,  
308 Ibadan, Nigeria pp 150 – 153.
- 309 8. Yadva, R.N.S., Munin, A. (2011). Phytochemical analysis of some medicinal plants. J. of  
310 Phyt. 3(12): 10-14.
- 311 9. Charles, Staples., Ricardo, Mattos., Scott, Boken., Lynn, Sollenberger., william, Thatcher.,  
312 and Tom, Jenkins (2014). Feeding Fatty acids for fertility, Proceedings of the 13th Annual  
313 Florida Ruminant Nutrition Symposium, pp 71-85.
- 314 10.Barzo P., Marmarou A., Fatouros P., Corwin F., Dunbar J (1996). Magnetic resonance  
315 imaging-monitored acute blood-brain barrier changes in experimental traumatic brain injury.  
316 J. Neurosurg. 85(21) 1113–1121.
- 317 11. Teresa C. Hill (2012). The importance of Triglyceride/ HDL ratio. *Advocate* 970: 522-  
318 537.
- 319 12. Assmann, G., Schulte, E., (1992). The importance of triglyceride. Euro. J. of  
320 Epidemiology. 8(1): 99.
- 321 13. Anatoli, P.S., Elvino B., Raffaella G. and Anna L.S (2005). Metabolic profile of lettuce  
322 leaves by high-field NMR spectra, MRC pp 625-638.
- 323
- 324 14. Finar, I. L (2013). Stereochemistry and the Chemistry of Natural Products, Dorling  
325 Kindersley, New Delhi, India, pp 78-80.

- 326 15. Harbourne, J.B. (1997). Phytochemical Methods, Chapman and Hall Ltd, London, pp 1 –  
327 29.
- 328 16. Spectral data base for organic compounds from [http://sdb.s.aist.go.jp/sdb/cgi-bin/frame\\_top.cgi](http://sdb.s.aist.go.jp/sdb/cgi-bin/frame_top.cgi) 19228HSP-46-188, 20-12-2016.  
329

UNDER PEER REVIEW